

MAST30025 assignment 3

● Graded

Student

James La Fontaine

Total Points

37 / 46 pts

Question 1

Q1

7 / 7 pts

1.1 1(a)

2 / 2 pts

The rubric is hidden for this question.

1.2 1(b)

2 / 2 pts

The rubric is hidden for this question.

1.3 1(c)

■ 3 / 3 pts

The rubric is hidden for this question.



Please check the solution.

Question 2

Q2

11 / 11 pts

2.1 2(a)

2 / 2 pts

The rubric is hidden for this question.

2.2 2(b)

3 / 3 pts

The rubric is hidden for this question.

2.3 2(c)

2 / 2 pts

The rubric is hidden for this question.

2.4 2(d)

3 / 3 pts

The rubric is hidden for this question.

2.5 2(e)

1 / 1 pt

The rubric is hidden for this question.

Question 3

Q3

0 / 5 pts

The rubric is hidden for this question.

Question 4

Q4

8 / 11 pts

4.1 4(a)

2 / 3 pts

The rubric is hidden for this question.

2 Please check the solution.

4.2 4(b)

2 / 2 pts

The rubric is hidden for this question.

4.3 4(c)

2 / 2 pts

The rubric is hidden for this question.

4.4 4(d)

2 / 2 pts

The rubric is hidden for this question.

4.5 4(e)

0 / 2 pts

The rubric is hidden for this question.

4 Please check the solution.

Question 5

Q5

11 / 12 pts

5.1 5(a)

1 / 2 pts

The rubric is hidden for this question.

1 Please check the solution.

5.2 5(b)

2 / 2 pts

The rubric is hidden for this question.

5.3 5(c)

3 / 3 pts

The rubric is hidden for this question.

5.4 5(d)

2 / 2 pts

The rubric is hidden for this question.

5.5 5(e)

3 / 3 pts

The rubric is hidden for this question.

No questions assigned to the following page.

MAST30025 Linear Statistical Models
Assignment 3

Student Code: 1079860

May 2023

Questions assigned to the following page: [1.1](#) and [1.2](#)

Question 1

(a)

$$r(A^c A) \leq \min[r(A^c), r(A)] \leq r(A)$$

$$r(A) = r(AA^c A) \leq \min[r(A), r(A^c A)] \leq r(A^c A)$$

$$\implies r(A^C A) = r(A)$$

(b)

$$(I - A(A^T A)^c A^T)(I - A(A^T A)^c A^T)$$

$$= I^2 - IA(A^T A)^c A^T - A(A^T A)^c A^T I + A(A^T A)^c A^T A(A^T A)^c A^T$$

$$= I - 2(A(A^T A)^c A^T) + A(A^T A)^c A^T A(A^T A)^c A^T$$

$$= I - 2(A(A^T A)^c A^T) + [A(A^T A)^c A^T A] (A^T A)^c A^T$$

$$= I - 2(A(A^T A)^c A^T) + A(A^T A)^c A^T$$

$I - A(A^T A)^c A^T$ is idempotent, I and $A(A^T A)^c A^T$ are idempotent and symmetric

Question assigned to the following page: [1.3](#)

(c)

$$r(A(A^T A)^c A^T) \leq \min[r(A), r((A^T A)^c), r(A^T)] \leq r(A)$$

$$r(A) = r(A(A^T A)^c A^T A) \leq \min[r(A), r(A(A^T A)^c A^T)] \leq r(A(A^T A)^c A^T)$$

$$\implies r(A(A^T A)^c A^T) = r(A)$$

$A(A^T A)^c A^T$ is $n \times n$ so I must be $n \times n$

$$\implies r(I) = n$$

As $I - A(A^T A)^c A^T$ is symmetric and $n \times n$, it is diagonalised by P , so
 $r(I - A(A^T A)^c A^T) = r(P^T(I - A(A^T A)^c A^T)P) = r(I - D)$

Since $A(A^T A)^c A^T$ is idempotent, it has $r(A(A^T A)^c A^T) = r(A)$ 1s on the diagonal of its diagonal matrix D

$\implies r(I - D) = r(I) - r(D) = n - r(A)$ as the rank of a diagonal matrix = the number of non-zero entries on its diagonal

Question assigned to the following page: [2.1](#)

Question 2


(a)

[R code]

```
rot_df = data.frame(rotting = c(13, 11, 3, 26, 19, 24, 10, 4, 7, 15,
                                22, 18, 15, 2, 7, 20, 24, 8),
                    oxygen_level = c(rep(1, 6), rep(2, 6), rep(3, 6)),
                    temperature = c(rep(c(10, 16), 3, each=3)) )

rot_df$oxygen_level = factor(rot_df$oxygen_level)
rot_df$temperature = factor(rot_df$temperature)

with(rot_df, (interaction.plot(oxygen_level, temperature, rotting)))
```



It may be reasonable to assume that there is no interaction due to the lines being relatively parallel despite the low sample size, although its hard to be certain.

Question assigned to the following page: [2.2](#)

(b)

[R code]

```
> y = rot_df$rotting
> n = length(y)
> X = matrix(c(rep(1, n), rep(0, n*5)), n, 6)
> X[cbind(1:n, as.numeric(rot_df$oxygen_level)+1)] = 1
> X[cbind(1:n, as.numeric(rot_df$temperature)+4)] = 1
>
> library(Matrix)
>
> r=rankMatrix(X)[1]
>
> # find conditional inverse of XtX
>
> XtX = t(X) %*% X
>
> M = XtX[2:5, 2:5]
>
> det(M)
[1] 972
>
> XtXc = matrix(0, 6, 6)
>
> XtXc[2:5, 2:5] = t(solve(M))
>
> XtXc = t(XtXc)
>
> all.equal(XtX %*% XtXc %*% XtX, XtX)
[1] TRUE
>
> b = XtXc %*% t(X) %*% y
>
> s2 = sum((y - X %*% b)^2) / (n-r)
```

Question assigned to the following page: [2.2](#)

```

> X
      [,1] [,2] [,3] [,4] [,5] [,6]
[1,]      1      1      0      0      1      0
[2,]      1      1      0      0      1      0
[3,]      1      1      0      0      1      0
[4,]      1      1      0      0      0      1
[5,]      1      1      0      0      0      1
[6,]      1      1      0      0      0      1
[7,]      1      0      1      0      1      0
[8,]      1      0      1      0      1      0
[9,]      1      0      1      0      1      0
[10,]     1      0      1      0      0      1
[11,]     1      0      1      0      0      1
[12,]     1      0      1      0      0      1
[13,]     1      0      0      1      1      0
[14,]     1      0      0      1      1      0
[15,]     1      0      0      1      1      0
[16,]     1      0      0      1      0      1
[17,]     1      0      0      1      0      1
[18,]     1      0      0      1      0      1
>
> s2
[1] 26.12698

```

Questions assigned to the following page: [2.3](#), [2.4](#), and [2.5](#)

(c)

[R code]

```
> tt = c(0, 0, 0, 0, 1, -1) # temp10mean - temp16mean
>
> ta = qt(0.975, n-r)
>
> halfwidth = ta * sqrt(s2 * t(tt) %*% XtXc %*% tt)
>
> tt %*% b + c(-1, 1) * halfwidth
[1] -16.723555 -6.387556
```

95% confidence interval for temp10effect - temp16effect :

[-16.724, -6.388]

(d)

$H_0 : \tau_1 = \tau_2 = \tau_3 = 0$

[R code]

```
> C = matrix(c(0,0,1,1,-1,0,0,-1,0,0,0,0), 2, 6)
>
> all.equal(round(C %*% XtXc %*% t(X) %*% X, 3), C)
[1] TRUE
>
> numer = t(C %*% b) %*% solve(C %*% XtXc %*% t(C)) %*% C%*%b
>
> Fstat = (numer / 2) / s2
>
> pf(Fstat, 2, n-r, lower=F)
      [,1]
[1,] 0.4481124
```

$p\text{-value} = 0.448 \implies \text{Cannot reject } H_0 \text{ at } 5\% \text{ significance level}$

(e)

This would be a complete block design study with oxygen level as the factor of interest and temperature as the blocking factor.

Question assigned to the following page: [3](#)

Question 3

Question assigned to the following page: [4.1](#)

Question 4


(a)

[R code]

```
toycars = read.csv("toycars.csv")
```

```
toycars$car = factor(toycars$car)
```

```
plot(toycars$angle, toycars$distance, pch=array(toycars$car),  
col=as.numeric(toycars$car)+1, cex=1.5)
```



It seems that the distance travelled by the car increases as the angle increases. It would also seem that the type of car doesn't seem to have as much of an effect as the angle in this experiment, although we can still say that car 3 generally travels the least distance, while car 1 travels slightly further and car 2 travels the furthest.

Question assigned to the following page: [4.2](#)

(b)

[R code]

```
> imodel = lm(toycars$distance ~ toycars$car * toycars$angle, data=toycars)
>
> amodel = lm(toycars$distance ~ toycars$car + toycars$angle, data=toycars)
>
> anova(amodel, imodel)
Analysis of Variance Table

Model 1: toycars$distance ~ toycars$car + toycars$angle
Model 2: toycars$distance ~ toycars$car * toycars$angle
  Res.Df      RSS Df Sum of Sq    F Pr(>F)
1      23 0.105657
2      21 0.093271  2   0.012386 1.3944   0.27
```

No significant interaction present between the type of toy car and the angle

Question assigned to the following page: [4.3](#)

(c)

[R code]

```
> fullmodel = imodel
>
> drop1(fullmodel, scope = ~ ., test="F")
Single term deletions

Model:
toycars$distance ~ toycars$car * toycars$angle
              Df Sum of Sq      RSS       AIC  F value
Pr(>F)
<none>                        0.09327  -141.038
toycars$car                2    0.01979  0.11307  -139.842
2.2284    0.1325
toycars$angle              1    0.44593  0.53920   -95.664 100.4023 1.87e-09
toycars$car:toycars$angle  2    0.01239  0.10566  -141.672
1.3944    0.2700

<none>
toycars$car
toycars$angle ***
toycars$car:toycars$angle
-----
Signif. codes:  0   ***    0.001   **    0.01   *    0.05   .    0.1
>
> backmodel2 = lm(toycars$distance ~ toycars$car + toycars$angle, data=toycars)
>
> drop1(backmodel2, scope = ~ ., test="F")
Single term deletions

Model:
toycars$distance ~ toycars$car + toycars$angle
              Df Sum of Sq      RSS       AIC  F value    Pr(>F)
<none>                        0.10566  -141.672
toycars$car                2    0.16945  0.27511  -119.833   18.444 1.662e-05 ***
toycars$angle              1    1.65108  1.75673   -67.774 359.416 1.547e-15 ***
-----
Signif. codes:  0   ***    0.001   **    0.01   *    0.05   .    0.1
>
> # all tests are significant, we stop at backmodel2
```


Question assigned to the following page: [4.4](#)

(d)

$$H_0 : \tau_1 - \tau_3 = 0.05$$

[R code]

```
> linearHypothesis(backmodel2, c(0,0,-1, 0), 0.05)
Linear hypothesis test
```

```
Hypothesis:
- toycars$scar3 = 0.05
```

```
Model 1: restricted model
Model 2: toycars$distance ~ toycars$scar + toycars$angle
```

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	24	0.11033				
2	23	0.10566	1	0.0046722	1.0171	0.3237

Signif. codes:	0	***	0.001	**	0.01	*	0.05	.	0.1
----------------	---	-----	-------	----	------	---	------	---	-----

p-value = 0.3237 > 0.05, so we cannot reject H_0 at the 5% significance level

Question assigned to the following page: [4.5](#)

(e)

$$H_0 : \tau_2 = \tau_3$$

[R code]

```
> linearHypothesis(fullmodel, c(0,1,-1, 0, 0, 0), 0)
Linear hypothesis test
```

```
Hypothesis:
toycars$car2 - toycars$car3 = 0
```

```
Model 1: restricted model
Model 2: toycars$distance ~ toycars$car * toycars$angle
```

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	22	0.108767				
2	21	0.093271	1	0.015497	3.4891	0.07579 .

Signif. codes:	0	***	0.001	**	0.01	*	0.05	.	0.1
----------------	---	-----	-------	----	------	---	------	---	-----

p-value = 0.0758 > 0.05, so we cannot reject H_0 at the 5% significance level

Questions assigned to the following page: [5.1](#) and [5.2](#)

Question 5

(a)

μ can be regarded as a nuisance parameter since we are not interested in it specifically but it still accounts for some variation in our model

1

(b)

$$X_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

$$X_2 = \begin{bmatrix} 1 & 0 & x_{11} \\ 1 & 0 & x_{12} \\ 1 & 0 & x_{13} \\ 0 & 1 & x_{21} \\ 0 & 1 & x_{22} \\ 0 & 1 & x_{23} \end{bmatrix}$$

Question assigned to the following page: [5.3](#)

(c)

[R code]

```
> X1 = matrix(c(rep(1, 6)), 6, 1)
>
> H1 = X1 %*% ginv(t(X1) %*% X1) %*% t(X1)
>
> fractions(diag(6) - H1)
      [,1] [,2] [,3] [,4] [,5] [,6]
[1,]  5/6 -1/6 -1/6 -1/6 -1/6 -1/6
[2,] -1/6  5/6 -1/6 -1/6 -1/6 -1/6
[3,] -1/6 -1/6  5/6 -1/6 -1/6 -1/6
[4,] -1/6 -1/6 -1/6  5/6 -1/6 -1/6
[5,] -1/6 -1/6 -1/6 -1/6  5/6 -1/6
[6,] -1/6 -1/6 -1/6 -1/6 -1/6  5/6
```

$$\begin{aligned}
 X_{2|1} = [I - H_1]X_2 &= \begin{bmatrix} \frac{5}{6} & -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} \\ -\frac{1}{6} & \frac{5}{6} & -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} \\ -\frac{1}{6} & -\frac{1}{6} & \frac{5}{6} & -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} \\ -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} & \frac{5}{6} & -\frac{1}{6} & -\frac{1}{6} \\ -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} & \frac{5}{6} & -\frac{1}{6} \\ -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} & \frac{5}{6} \end{bmatrix} \begin{bmatrix} 1 & 0 & x_{11} \\ 1 & 0 & x_{12} \\ 1 & 0 & x_{13} \\ 0 & 1 & x_{21} \\ 0 & 1 & x_{22} \\ 0 & 1 & x_{23} \end{bmatrix} \\
 &= \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} & x_{11} - \bar{x} \\ \frac{1}{2} & -\frac{1}{2} & x_{12} - \bar{x} \\ \frac{1}{2} & -\frac{1}{2} & x_{13} - \bar{x} \\ -\frac{1}{2} & \frac{1}{2} & x_{21} - \bar{x} \\ -\frac{1}{2} & \frac{1}{2} & x_{22} - \bar{x} \\ -\frac{1}{2} & \frac{1}{2} & x_{23} - \bar{x} \end{bmatrix}
 \end{aligned}$$

Question assigned to the following page: [5.4](#)

(d)

$$\vec{b}_2 = (X_{2|1}^T X_{2|1})^c X_{2|1}^T \vec{y}$$

[R code]

```
> xs = c(2,4,8,7,6,4)
>
> y = c(4,2,10,8,8,12)
>
> xbar = mean(xs)
>
> X21 = matrix(c(0.5,0.5,0.5,-0.5,-0.5,-0.5,-0.5,-0.5,-0.5,0.5,0.5,0.5,
               xs[1]-xbar,xs[2]-xbar,xs[3]-xbar,xs[4]-xbar,xs[5]-xbar,
               xs[6]-xbar), 6,3)
>
> b2 = ginv(t(X21) %*% X21) %*% t(X21) %*% y
>
> b2
               [,1]
[1,] -1.6857143
[2,]  1.6857143
[3,]  0.6285714
```

$$\vec{b}_2 = \begin{bmatrix} -1.6857 \\ 1.6857 \\ 0.6286 \end{bmatrix}$$

Question assigned to the following page: [5.5](#)

(e)

$$\vec{b}_1 = (X_1^T X_1)^c (X_1^T \vec{y} - X_1^T X_2 \vec{b}_2)$$

[R code]

```
> b1 = ginv(t(X1) %*% X1) %*% (t(X1) %*% y - t(X1) %*% X2 %*% b2)
>
> b1
      [,1]
[1,] 4.085714
>
> X = cbind(X1, X2)
>
> b = rbind(b1, b2)
>
> t(X) %*% X %*% b
      [,1]
[1,] 44
[2,] 16
[3,] 28
[4,] 248
>
> t(X) %*% y
      [,1]
[1,] 44
[2,] 16
[3,] 28
[4,] 248
```

$$\vec{b}_1 = 4.0857$$

$$X^T X \vec{b} = X^T \vec{y}$$