

Assignment 5

● Graded

Student

James La Fontaine

Total Points

11 / 11 pts

Question 1

Question 1

11 / 11 pts

✓ + 1 pt 1a) M: Obtain an equation relating $\frac{du}{dx}$ and $\frac{dy}{dx}$.

✓ + 1 pt 1a) A: Substitute into original ODE and demonstrate algebraic steps with no errors to obtain $\frac{du}{dx} = \frac{\sqrt{1+u^2}}{x}$.

✓ + 1 pt 1b) M: Use separation of variables.

✓ + 1 pt 1b) A: $\operatorname{arcsinh}$ and \log appear as integrals.

✓ + 1 pt 1b) A: One of the answers equivalent to $\frac{A^2 x^2 - 1}{2Ax}$ or $\sinh(\log x + c)$.

✓ + 1 pt 1c) M: Multiply previous answer by x to write y in terms of x .

✓ + 1 pt 1b) or 1c) M: Use exponential form for \sinh or \log form for $\operatorname{arcsinh}$.

✓ + 1 pt 1c) J: Limit laws or other justification for calculating the limit.

✓ + 1 pt 1c) A: $y(x) = \frac{x^2 - e^2}{2e}$ or $y(x) = x \sinh(\log x - 1)$.

Note: An answer like $y(x) = x \sinh(\log x - 1)$ was accepted, but is often better simplified using the exponential form for \sinh .

✓ + 1 pt 1c) M: Use limit condition to find c .

✓ + 1 pt N: Notation is correct.

+ 0 pts No attempt or incorrect working



Great, good work!

Question assigned to the following page: [1](#)

Calculus 2 Written Assignment 5

1. $x \frac{dy}{dx} = y + \sqrt{x^2 + y^2}$ $x > 0$

a) $u = y/x$

$$\Rightarrow y = ux$$

$$\Rightarrow \frac{dy}{dx} = x \frac{du}{dx} + u \quad , \quad \text{product rule}$$

$$x \frac{dy}{dx} = y + \sqrt{x^2 + y^2}$$

$$\Rightarrow x \left(x \frac{du}{dx} + u \right) = ux + \sqrt{x^2 + u^2 x^2}$$

$$\Rightarrow x \frac{du}{dx} + u = u + \frac{\sqrt{x^2 + u^2 x^2}}{x} \quad , \quad x > 0$$

$$\Rightarrow \frac{du}{dx} = \frac{\sqrt{x^2(1+u^2)}}{x^2} \quad , \quad x > 0$$

$$\Rightarrow \frac{du}{dx} = \frac{\sqrt{1+u^2}}{x^2} \cdot x \quad , \quad x > 0$$

$$\Rightarrow \frac{du}{dx} = \frac{\sqrt{1+u^2}}{x}$$

Question assigned to the following page: [1](#)

b) $\frac{du}{dx} = \sqrt{1+u^2} \cdot \frac{1}{x}$, is separable
 - use sep. of variables

$$\frac{1}{\sqrt{1+u^2}} \frac{du}{dx} = \frac{1}{x} , \quad \sqrt{1+u^2} \geq 1$$

$$\int \frac{1}{\sqrt{1+u^2}} du = \int \frac{1}{x} dx$$

$$\operatorname{arcsinh}(u) = \log(x) + C , \quad x > 0$$

$$u = \sinh(\log(x) + C)$$

$$u = \frac{1}{2} (e^{\log(x)+C} - e^{-(\log(x)+C)})$$

$$u = \frac{1}{2} (e^{\log(x)+C} - \frac{1}{e^{\log(x)+C}})$$

$$u = \frac{1}{2} (e^C x - \frac{1}{e^C x}) = \frac{e^C x}{2} - \frac{1}{2e^C x}$$

$$u(x) = \frac{Ax}{2} - \frac{1}{2Ax} , \quad A = e^C$$

Question assigned to the following page: [1](#)

$$c) \frac{y}{x} = \frac{Ax}{2} - \frac{1}{2Ax}$$

$$y = \frac{Ax^2}{2} - \frac{1}{2A}$$

$$\lim_{x \rightarrow 0^+} y(x) = -\frac{e}{2}$$

$$\lim_{x \rightarrow 0^+} \left(\frac{Ax^2}{2} - \frac{1}{2A} \right) = -\frac{e}{2}$$

$$\lim_{x \rightarrow 0^+} \left(\frac{A}{2} x^2 \right) - \lim_{x \rightarrow 0^+} \left(\frac{1}{2A} \right) = -\frac{e}{2}, \text{ limit laws}$$

$$\frac{A}{2} \lim_{x \rightarrow 0^+} (x^2) - \frac{1}{2A} = -\frac{e}{2}, \text{ limit laws}$$

$$\frac{A}{2} \left(\lim_{x \rightarrow 0^+} x \right)^2 - \frac{1}{2A} = -\frac{e}{2}, \text{ limit laws}$$

$$\frac{A}{2} (0)^2 - \frac{1}{2A} = -\frac{e}{2}, \text{ limit laws}$$

$$-\frac{1}{2A} = -\frac{e}{2}$$

$$\frac{1}{A} = e$$

$$A = \frac{1}{e} = e^{-1}$$

$$\text{So } y(x) = \frac{x^2}{2e} - \frac{e}{2}$$

No questions assigned to the following page.

$$2. t \log(t) \frac{dr}{dt} + r = \frac{t}{(2t^2 - 9)^{3/2}}$$

Rewrite as:

$$\frac{dr}{dt} + \frac{1}{t \log(t)} r = \frac{t}{t \log(t) (2t^2 - 9)^{3/2}}, \quad t > 0, \log(t) \neq 0$$

$$\frac{dr}{dt} + \frac{1}{t \log(t)} r = \frac{1}{\log(t) (2t^2 - 9)^{3/2}}$$

Is linear with $P(x) = \frac{1}{t \log(t)}$

$$Q(x) = \frac{1}{\log(t) (2t^2 - 9)^{3/2}}$$

• Find an integrating factor:

$$I(x) = e^{\int \frac{1}{t \log(t)} dt}$$

Solve: $\int \frac{1}{t \log(t)} dt$

Let $u = \log(t)$

$$\frac{du}{dt} = \frac{1}{t}$$

Derivative
Substitution

$$\int \frac{du}{dt} \frac{1}{u} dt = \int \frac{1}{u} du = \log|u| + C$$

$$\Rightarrow I(x) = e^{(\log|\log t| + C)}$$

$$= |\log t| = \log t, \quad \text{only need 1 integrating factor}$$

No questions assigned to the following page.

• Multiply ODE by I

$$\log(t) \frac{dr}{dt} + \frac{1}{t} r = \frac{1}{(2t^2-9)^{3/2}}$$

$$\Rightarrow \frac{d}{dt} (\log(t) r) = \frac{1}{(2t^2-9)^{3/2}}$$

$$\Rightarrow \log(t) r = \int \frac{1}{(2t^2-9)^{3/2}} dt$$

$$\text{Solve: } \int \frac{1}{(2t^2-9)^{3/2}} dt$$

$$= \int \frac{1}{(2(t^2-\frac{9}{2}))^{3/2}} dt$$

$$= \int \frac{1}{2\sqrt{2}} \frac{1}{(t^2-\frac{9}{2})^{3/2}} dt$$

$$= \frac{1}{2\sqrt{2}} \int \frac{1}{(t^2-\frac{9}{2})^{3/2}} dt$$

$$\text{Let } t = \frac{3}{\sqrt{2}} \cosh \Theta$$

$$\Theta = \operatorname{arccosh}\left(\frac{\sqrt{2}t}{3}\right), \quad \text{Hyperbolic Substitution}$$

This is valid when

$$\frac{\sqrt{2}t}{3} \in \operatorname{dom}(\operatorname{arccosh}) \quad \text{and} \quad \Theta \in \operatorname{range}(\operatorname{arccosh})$$

$$\Rightarrow \frac{\sqrt{2}t}{3} \geq 1 \quad \text{and} \quad \Theta \geq 0$$

$$\Rightarrow t \geq \frac{3}{\sqrt{2}} \quad \text{and} \quad \Theta \geq 0$$

$$\text{Also need } (\sqrt{t^2-\frac{9}{2}})^3 \neq 0$$

$$\Rightarrow t > \frac{3}{\sqrt{2}} \Rightarrow \Theta > 0$$

No questions assigned to the following page.

$$t = \frac{3}{\sqrt{2}} \cosh \theta$$

$$\cosh \theta = \frac{\sqrt{2}t}{3}$$

$$\cdot \frac{dt}{d\theta} = \frac{3}{\sqrt{2}} \sinh \theta$$

$$\cdot \frac{1}{\left(\sqrt{t^2 - \frac{9}{2}}\right)^3} = \frac{1}{\left(\sqrt{\frac{9}{2} \cosh^2 \theta - \frac{9}{2}}\right)^3}$$

$$= \frac{1}{\left(\sqrt{\frac{9}{2} (\cosh^2 \theta - 1)}\right)^3} = \frac{1}{\left(\frac{\sqrt{3}}{2} \sqrt{\sinh^2 \theta}\right)^3}$$

$$= \frac{1}{\left(\frac{\sqrt{3}}{2} (\sinh \theta)\right)^3}, \quad \sinh \theta > 0 \text{ as } \theta > 0$$

$$= \frac{1}{\frac{3\sqrt{3}}{8} \sinh^3 \theta} = \frac{8}{3\sqrt{3} \sinh^3 \theta}$$

$$\text{Therefore } \frac{1}{2\sqrt{2}} \int \frac{1}{(t^2 - \frac{9}{2})^{3/2}} dt$$

$$= \frac{1}{2\sqrt{2}} \int \frac{8}{3\sqrt{3} \sinh^3 \theta} \cdot \frac{3}{\sqrt{2}} \sinh \theta d\theta$$

$$= \frac{1}{2\sqrt{2}} \int \frac{8}{\sqrt{6} \sinh^2 \theta} d\theta$$

$$= \frac{1}{2\sqrt{2}} \int \frac{8\sqrt{6}}{6 \sinh^2 \theta} d\theta$$

$$= \frac{8\sqrt{6}}{12\sqrt{2}} \int \frac{1}{\sinh^2 \theta} d\theta$$

$$= \frac{2\sqrt{3}}{3} \int \operatorname{cosech}^2 \theta d\theta$$

$$= -\frac{2\sqrt{3}}{3} [\coth \theta + C]$$

No questions assigned to the following page.

$$= -\frac{2\sqrt{3}}{3} \left[\frac{\cosh \theta}{\sinh \theta} + C \right]$$

$$= -\frac{2\sqrt{3}}{3} \left[\frac{\cosh(\operatorname{arccosh}(\frac{\sqrt{2}t}{3}))}{\sqrt{\cosh^2 \theta - 1}} + C \right]$$

$$= -\frac{2\sqrt{3}}{3} \left[\frac{\frac{\sqrt{2}t}{3}}{\sqrt{\frac{2}{9}t^2 - 1}} + C \right], \quad t > \frac{3}{\sqrt{2}}$$

$$= -\frac{2\sqrt{6}t}{9} \cdot \frac{1}{\sqrt{\frac{2}{9}t^2 - 1}} + d, \quad d = -\frac{2\sqrt{3}}{3} \cdot C$$

$$= -\frac{2\sqrt{6}t}{3\sqrt{2t^2 - 9}} + d$$

Therefore:

$$\log(t)r = -\frac{2\sqrt{6}t}{3\sqrt{2t^2 - 9}} + d$$

$$r = -\frac{2\sqrt{6}t}{3\log(t)\sqrt{2t^2 - 9}} + \frac{d}{\log(t)}$$

