

# (Hidden) assumptions of simple compartmental ODE models

Rebecca Borchering, PhD

U.S. Centers for Disease Control and Prevention (CDC)

University of Florida and MMED Alumnus


**ICI3D Faculty** 

**MMED 2023** 

### Goals

- Review the main uses of applied epidemiological modelling
- Introduce our conceptual framework for applied modelling
- Review commonly overlooked assumptions that are inherent in the structure of simple compartmental ODE models
- Discuss when these assumptions might be problematic, and when they may be desirable
- Begin to explore some alternative model structures that relax the assumptions





Related to the distribution and determinants of healthrelated states and events







Insight

- Improving understanding of the dynamics of health and disease
- Translation of results into decision-making and communication tools



Insight

**Estimation** 

• Improving measurement and interpretation of key health indicators at the population and individual levels



Insight

Estimation

Prediction

• Projection and forecasting of expected future trends



Insight

**Estimation** 

Prediction

**Planning** 

- Guiding study design and intervention roll-out
- Informing decisions through analysis and comparison of policy scenarios



Insight

**Estimation** 

Prediction

**Planning** 

Assessment

- Evaluating the impact of public health interventions
  - Assessing risk of future public health events



Insight

- Improving understanding of the dynamics of health and disease
- Translation of results into decision-making and communication tools

Estimation

• Improving measurement and interpretation of key health indicators at the population and individual levels

Prediction

Projection and forecasting of expected future trends

**Planning** 

- Guiding study design and intervention roll-out
- Informing decisions through analysis and comparison of policy scenarios

Assessment


- Evaluating the impact of public health interventions
  - Assessing risk of future public health events



### Goals

- Review the main uses of applied epidemiological modelling
- Introduce our conceptual framework for applied modelling
- Review commonly overlooked assumptions that are inherent in the structure of simple compartmental ODE models
- Discuss when these assumptions might be problematic, and when they may be desirable
- Begin to explore some alternative model structures that relax the assumptions






### Model Worlds

- A model world is an abstraction of the world that is simple and fully specified, which we construct to help us understand particular aspects of the real world
- A mathematical model is a formal description of the assumptions that define a model world
  - We know exactly what assumptions we've made, and we can follow those assumptions to their logical conclusions to address research questions




### The SIR Model World





### SIR: ODE Model



$$I = \frac{bI}{N}$$


$$\frac{dS}{dt} = -\frac{bSI}{N}$$

$$N=S+I+R$$

$$\frac{dI}{dt} = \frac{bSI}{N} - gI$$


$$\frac{dR}{dt} = gI$$

### SIR: Reed-Frost Model






### SIR: Stochastic Reed-Frost



### SIR: Chain Binomial Model



$$S_{t+\Delta t} = S_t - X$$

$$I_{t+\Delta t} = I_t + X - Y$$

$$R_{t+\Delta t} = R_t + Y$$


$$\mathbb{P}(X = x) = \binom{S_t}{x} p^x (1-p)^{S_t - x}$$

$$\mathbb{P}(Y = y) = \binom{I_t}{y} r^y (1-r)^{I_t - y}$$

$$\mathbb{P}(X=x) = \binom{S_t}{x} p^x (1-p)^{S_t - x}$$

$$\mathbb{P}(Y=y) = \binom{I_t}{y} r^y (1-r)^{I_t-y}$$

### The SIR Model Family



# A mathematical model is formal description of the assumptions that define a model world





#### **CONTINUOUS TREATMENT OF INDIVIDUALS**

(averages, proportions, or population densities)

#### **DISCRETE TREATMENT OF INDIVIDUALS**

#### **CONTINUOUS TIME**

- Ordinary differential equations
- Partial differential equations

#### **DISCRETE TIME**

Difference equations
 (eg, Reed-Frost type models)

#### **CONTINUOUS TIME**

Stochastic differential equations

#### **DISCRETE TIME**

• Stochastic difference equations

#### **CONTINUOUS TIME**

• Gillespie algorithm

#### **DISCRETE TIME**



#### **CONTINUOUS TREATMENT OF INDIVIDUALS**

**DISCRETE TREATMENT OF INDIVIDUALS** 

(averages, proportions, or population densities)

#### **CONTINUOUS TIME**

- Ordinary differential equations
- Partial differential equations

#### **DISCRETE TIME**

Difference equations
 (eg, Reed-Frost type models)

#### **CONTINUOUS TIME**

Stochastic differential equations

#### **DISCRETE TIME**

• Stochastic difference equations

#### **CONTINUOUS TIME**

Gillespie algorithm

#### **DISCRETE TIME**



#### CONTINUOUS TREATMENT OF INDIVIDUALS

(averages, proportions, or population densities)

DISCRETE TREATMENT OF INDIVIDUALS

#### **CONTINUOUS TIME**

- Ordinary differential equations
- Partial differential equations

#### **DISCRETE TIME**

 Difference equations (eg, Reed-Frost type models)

#### **CONTINUOUS TIME**

Stochastic differential equations

#### **DISCRETE TIME**

Stochastic difference equations

#### **CONTINUOUS TIME**

Gillespie algorithm

#### **DISCRETE TIME**



#### **CONTINUOUS TREATMENT OF INDIVIDUALS**

.

**DISCRETE TREATMENT OF INDIVIDUALS** 

(averages, proportions, or population densities)

#### **CONTINUOUS TIME**

- Ordinary differential equations
- Partial differential equations

#### **DISCRETE TIME**

Difference equations
 (eg, Reed-Frost type models)

#### **CONTINUOUS TIME**

Stochastic differential equations

#### **DISCRETE TIME**

Stochastic difference equations


#### **CONTINUOUS TIME**

Gillespie algorithm

#### **DISCRETE TIME**



What is a **compartmental** model?



### Goals

- Review the main uses of applied epidemiological modelling
- Introduce our conceptual framework for applied modelling
- Review commonly overlooked assumptions that are inherent in the structure of simple compartmental ODE models
- Discuss when these assumptions might be problematic, and when they may be desirable
- Begin to explore some alternative model structures that relax the assumptions




#### **CONTINUOUS TREATMENT OF INDIVIDUALS**

DISCRETE TREATMENT OF INDIVIDUALS


(averages, proportions, or population densities)

CONTINUOUS TIME




large population size

CONTINUOUS TIME





### Demographic stochasticity




#### **CONTINUOUS TREATMENT OF INDIVIDUALS**

DISCRETE TREATMENT OF INDIVIDUALS


(averages, proportions, or population densities)





large population size


#### CONTINUOUS TIME





### Environmental stochasticity






### Compartmental ODE models assume

- Large population size
- Deterministic progression
  - for a given set of initial conditions and parameter values,
     a deterministic model always gives the same outcome

**Benefit:** Simplicity and consequences of assumptions facilitate quick assessments of what model outcomes are possible, and when

### Compartmental ODE models assume

- Large population size
- Deterministic progression
  - for a given set of initial conditions and parameter values,
     a deterministic model always gives the same outcome






CONTINUOUS TREATMENT OF INDIVIDUALS

DISCRETE TREATMENT OF INDIVIDUALS

(averages, proportions, or population densities)

#### **CONTINUOUS TIME**

Ordinary differential equations




### Simple ODE models assume

- Time proceeds in a continuous manner
- Parameter values remain constant

#### **CONTINUOUS TIME**

Ordinary differential equations




$$\frac{dN_t}{dt} = bN_t - dN_t$$

### Simple ODE models assume

- Time proceeds in a continuous manner
- Parameter values remain constant

#### **CONTINUOUS TIME**

Ordinary differential equations




$$r = b - d$$

$$\frac{dN_t}{dt} = rN_t$$

$$N_{t+Dt} = N_t e^{rDt}$$

#### **CONTINUOUS TIME**

Ordinary differential equations



$$r = b - d$$

$$\frac{dN}{dt} = rN$$

#### **DISCRETE TIME**

Difference equations

$$I_{\mathrm{D}t} = e^{r\mathrm{D}t}$$

$$N_{t+Dt} = I_{Dt}N_t$$

 $N_{t+\mathrm{D}t} = N_t e^{r\mathrm{D}t}$ 


## Simple compartmental ODE models assume

- Homogeneity within compartments
- Large population size
- Deterministic progression
- Time proceeds in a continuous manner
- Parameter values remain constant
- Memory-less processes



### Simple ODE models assume

Memory-less processes




$$\frac{N_t}{N_0} = e^{-mt}$$

$$\frac{dN_t}{dt} = -mN_t$$

$$N_t = N_0 e^{-mt}$$
 least  $\tau$  least  $\tau$  of  $\tau$  of  $\tau$  of  $\tau$  least  $\tau$  least  $\tau$  of  $\tau$  of


### Realistic waiting times



Time since infection


## Simple compartmental ODE models assume

- Homogeneity within compartments
- Large population size
- Deterministic progression
- Time proceeds in a continuous manner
- Parameter values remain constant







Simple ODE models are important tools for building understanding




- Simple ODE models are important tools for building understanding
- It's important to recognize the assumptions built into these models
  - When populations are small, average behaviors can be misleading



- Simple ODE models are important tools for building understanding
- It's important to recognize the assumptions built into these models
  - When populations are small, average behaviors can be misleading
  - When rates vary, simple ODEs can fail to reproduce important (observed) dynamics



- Simple ODE models are important tools for building understanding
- It's important to recognize the assumptions built into these models
  - When populations are small, average behaviors can be misleading
  - When rates vary, simple ODEs can fail to reproduce important (observed) dynamics




Dushoff lecture on heterogeneity (Wed)



- Simple ODE models are important tools for building understanding
- It's important to recognize the assumptions built into these models
  - When populations are small, average behaviors can be misleading
  - When rates vary, simple ODEs can fail to reproduce important (observed) dynamics



- The applied epidemiological modelling process requires
  - abstraction
  - specification and implementation
  - gaining an understanding of the dynamics
  - interpretation







This presentation is made available through a Creative Commons Attribution-Noncommercial license. Details of the license and permitted uses are available at

http://creativecommons.org/licenses/by/3.0/



© 2014-2023 International Clinics on Infectious Disease Dynamics and Data

(Hidden) assumptions of simple compartmental ODE models. DOI: 10.6084/m9.figshare.5044606

#### Attribution:

**Juliet Pulliam & Rebecca Borchering** 

Clinic on the Meaningful Modeling of Epidemiological Data

Source URL:

https://figshare.com/articles/Mathematical Assumptions of Simple ODE models/5044606

For further information please contact <a href="mailto:admin@ici3d.org">admin@ici3d.org</a>.





