

Modern Physics Departmental Exam

James Natoli

January 2021

Problem 1

The muon is an unstable particle that spontaneously decays into an electron and two neutrinos. If the number of muons at $t = 0$ is N_0 , the number at time t is given by $N = N_0 e^{-\frac{t}{\tau}}$, where τ is the mean lifetime, equal to $2.2 \mu\text{s}$. Suppose the muons move at a speed of $0.95c$ and there are $5.0 \cdot 10^4$ muons at $t = 0$

- a) What is the observed lifetime of the muons?
- b) How many muons remain after traveling a distance of 3.0 km?

Solution

- a) This is a time dilation problem, just need to know the formula

$$t' = \gamma t$$

where $\gamma = \frac{1}{\sqrt{1-(v/c)^2}}$, t' is the dilated time, and t is the regular time

$$t' = 2.2 \times \frac{1}{\sqrt{1-(0.95)^2}}$$

$$t' = 7.0456 \mu\text{s}$$

To help check your answer, it should always be a longer time than the regular time

- b) The muons will see a shorter distance, due to length contraction

$$L' = \frac{L}{\gamma}$$

where γ is same as before, L' is the contracted length, and L is the regular length

$$L' = 3 \times 10^3 \cdot \sqrt{1-0.95^2}$$

$$L' = 936.75 \text{ m}$$

We need a time for the formula given in the prompt, $N = N_0 e^{-\frac{t}{\tau}}$, so we need to find the time it takes for the muons to travel L' at $0.95c$

$$\frac{L'}{v} = \text{time} = \frac{936.75}{0.95c} = 3.28 \cdot 10^{-6} \text{ s}$$

$$N(t) = N_0 e^{-\frac{t}{\tau}} = 5 \cdot 10^4 \cdot e^{-\frac{3.28 \cdot 10^{-6}}{2.2 \cdot 10^{-6}}}$$

$$N(3.28 \cdot 10^{-6}) = 1.1223 \cdot 10^4 \text{ muons}$$

Problem 2

X-ray photons of wavelength 0.02580 nm are incident on a target and the Compton-scattered photons are observed at 90°

- a) What is the wavelength of the scattered photons?
- b) What is the momentum of the incident photons? Of the scattered photons?
- c) What is the kinetic energy of the scattered electrons?
- d) What is the momentum (magnitude and direction) of the scattered electrons?

Solution

a)

Problem

a)

Solution

a)