If a basis is not specified in a question, it is assumed to be the standard one e.g. $|0\rangle$, $|1\rangle$ for single qubit states, and $|00\rangle$, $|01\rangle$, $|10\rangle$, $|11\rangle$, in that order, for two qubit states.

The Pauli operators X, Y and Z on \mathbb{C}^2 are defined by

$$X|0\rangle = |1\rangle;$$
 $X|1\rangle = |0\rangle$
 $Y|0\rangle = i|1\rangle;$ $Y|1\rangle = -i|0\rangle$
 $Z|0\rangle = |0\rangle;$ $Z|1\rangle = -|1\rangle.$

1. Pauli operators/Pauli matrices

- (a) Calculate $YZ|0\rangle$ and $YZ|1\rangle$, and hence show for any state $|\psi\rangle = \alpha|0\rangle + \beta|1\rangle$ that $YZ|\psi\rangle = iX|\psi\rangle$. This shows that YZ = iX (why?).
- (b) Calculate the Pauli matrices σ_x , σ_y and σ_z , the matrices associated to X, Y and Z in the standard basis. Show that $\sigma_y \sigma_z = i\sigma_x$. (Notice the correspondence to part (a)).

2. Dirac Form

- (a) Find the matrix of *Y* with respect to the standard basis and hence show that *Y* is self-adjoint and unitary.
- (b) Write *Y* in Dirac form in the standard basis.
- (c) Using the Dirac form for Y, calculate $Y|+\rangle$ and $Y|-\rangle$, where $|+\rangle$ and $|-\rangle$ are the +1 and -1 eigenstates of X. Confirm that you get the same result by calculating these states directly from the definition of Y.

3. Matrix versus operator notation

Consider the operator A and the state $|\phi\rangle$ on \mathbb{C}^2 defined by

$$A|0\rangle = a|0\rangle + b|1\rangle;$$
 and $|\phi\rangle = \alpha|0\rangle + \beta|1\rangle$
 $A|1\rangle = c|0\rangle + d|1\rangle;$

 $a, b, c, d, \alpha, \beta$ are constants.

- (a) Find the matrix *M* of *A* with respect to the standard basis.
- (b) Find the Dirac form for A (i.e. write A in the form $\sum A_{ij}|i\rangle\langle j|$) with respect to the standard basis.
- (c) Calculate the action of the matrix M on the column vector $\begin{pmatrix} \alpha \\ \beta \end{pmatrix}$.
- (d) Calculate $A|\phi\rangle$ directly using operator notation. Confirm that your two calculations agree.

4. States and operators on \mathbb{C}^3

Show that the following three states form an orthonormal basis for \mathbb{C}^3

$$|\psi_{0}\rangle = \frac{1}{\sqrt{3}} (|0\rangle + |1\rangle + |2\rangle);$$

$$|\psi_{1}\rangle = \frac{1}{\sqrt{3}} (|0\rangle + \omega|1\rangle + \omega^{2}|2\rangle);$$

$$|\psi_{2}\rangle = \frac{1}{\sqrt{3}} (|0\rangle + \omega^{2}|1\rangle + \omega|2\rangle),$$

where $|0\rangle$, $|1\rangle$ and $|2\rangle$ are orthonormal basis elements for \mathbb{C}^3 and $\omega=\exp(2\pi i/3)$.

Write down the matrix of the operator M that satisfies $M|a\rangle = |\psi_a\rangle$ for a = 0,1,2. What can you say about M?

5. Self-adjoint and unitary operators

Let the operator W on \mathbb{C}^2 be defined by

$$W|0\rangle = |0\rangle + |1\rangle;$$

 $W|1\rangle = |1\rangle.$

- (a) Find the matrix of W in the standard basis, find its adjoint, W^{\dagger} ; you should write down $W^{\dagger}|0\rangle$ and $W^{\dagger}|1\rangle$ explicitly. Show that W is neither self-adjoint nor unitary.
- (b) Let the operator T be defined by $T = WW^{\dagger}$. Calculate $T|0\rangle$ and $T|1\rangle$. Show that T is self-adjoint, but not unitary.

[©] University of Bristol 2025. This material is copyright of the University unless explicitly stated otherwise. It is provided exclusively for educational purposes at the University and is to be downloaded or copied for your private study only.