

Photochemical production of oxylipin infochemicals in West Antarctica: An approach using new tools for semi-untargeted lipidomics

Jamie Collins ^{1,2}, Helen F. Fredricks ¹, and Benjamin A.S. Van Mooy ¹

¹Dept. Marine Chemistry & Geochemistry and ²MIT/WHOI Joint Program in Oceanography, Woods Hole Oceanographic Institution

Hugh W. Ducklow

Lamont-Doherty Earth Observatory, Columbia University


Introduction

We combine **high-resolution mass spectrometry** & **semi-untargeted lipidomics** to observe simultaneous changes over time in a **broad suite of compounds** involved in the **photochemical production** of **oxylipins** in the **surface ocean**


Oxylipins are bioactive “**infochemicals**” produced by **oxidation of intact lipids**. Oxylipins have **antimicrobial properties** and act as **stress/defense signals** in many terrestrial plant and aquatic systems. They can be produced by:

- regulated, **biologically-mediated pathways** (e.g., lipoxygenase)
- **abiotic** sources of **reactive oxygen species**, such as **photooxidation** (the subject of this study)


Study locations



Photodegradation experiments



Liposomes were added to glass vials in a **matrix** of 0.2 μm **filtered natural seawater** — containing both **DOC** and **NO₃** as potential chromophores — and **incubated at 1 m depth** in large-volume **on-deck aquaria**




Sample collection & HPLC/ESI-MS method

After incubation (experiments) **or retrieval via CTD** (environmental samples), lipids & oxylipins were extracted in 9:1 DCM : MeOH; an **internal standard** was added


HPLC/ESI-MS analysis: Thermo Exactive Orbitrap with reversed-phase chromatography; 100-1500 *m/z* scan range w/ionization mode switching (adapted from ref. 1)

For each experiment and environmental data set, **sample data were then pooled** and **screening** was performed using the **lipidomics data analysis pipeline** described in the adjacent column

Lipidomics data analysis pipeline³




Results from a lipid photodegradation experiment



Evidence in **differences between treatments**⁴ for photooxidation of an intact parent lipid, production of oxidized intermediate, and accumulation of a “terminal” oxylipin


Loss of unoxidized parent compound

18:2, 18:2 ($\Delta^9,12$ -*cis*) PC (36:4 PC)





Production of intact intermediate

36:4 PC + 20 hydroperoxide⁵



Accumulation of degradation product

18:2 + 20 hydroperoxy (oxylipin) fatty acid⁵



Summary of results & significance

With **high-resolution mass spectrometry** and a new **lipidomics pipeline**, we:



- **quantify thousands of compounds** at multiple positions in the lipid photooxidation reaction chain, and
- **identify**, with high confidence, **~ 10³ oxylipins** in a typical data set

Whereas rates of photooxidation are typically quantified indirectly (via assays such as TBARS or MDA) or by measuring only the accumulation of terminal degradation products (e.g., by GC/MS), our method allows us to **quantify rates from both upstream & downstream of a biogeochemically significant process**

Acknowledgements

We thank Eugene Melamed, Krista Longnecker, Liz Kujawinski, Winn Johnson, Colleen Hansell, Phil Gschwend, Jeff Bowman, Justin Ossolinski, Sebastian Edwards, John Brinckerhoff, Oscar Schofield, Tina Haskins, Oliver Ho, Austin Mello, Shellie Bence, Naomi Shelton, Sebastian Vivanco, Meredith Helfrich, Sandy Aylesworth, Julie Jackson, Karen Lipke, Greg Roberts, the 2013–2014 Palmer Station crew, and the captains and crewmembers of the R/V *Laurence M. Gould* (LMG 1401) and R/V *Knoir* (KN207-1 and KN207-3). This research was supported by the Long Term Ecological Research (LTER) program of the National Science Foundation, the Gordon and Betty Moore Foundation, and a U.S. Environmental Protection Agency (EPA) STAR Graduate Fellowship to J.R.C. under Fellowship Assistance Agreement no. FP-91744301-0. The contents of this poster have not been formally reviewed by EPA. The views expressed in this poster are solely those of the authors, and EPA does not endorse any products or commercial services mentioned in this poster.

Insight from **lipidomics**: Tracking the **fates of multiple species** over time



Hummel et al. 2011, *Front. Plant Sci.* z. ² Melamud et al. 2010, *Anal. Chem.* 82 (23): 9818-9826. ³ Lipidomics analysis metrics reported for positive mode HPLC/ESI-MS data (36 samples) from the Oct. 9, 2013, experiment. ⁴ Asterisk denotes significance of treatment difference from control (ANOVA; $p < 0.05$); error bars: \pm SE. ⁵ Structures putative; determination of specific sites of oxidation is part of planned future analysis.