

MATH 324 Notes

Contents

1	General	3
1.1	Trigonometric Identities	3
1.2	Polar Coordinates	3
1.3	Polar Curves	3
1.4	Curve Parametrization	3
2	Section Fourteen	4
2.1	Chain Rule of Single Variable Functions	4
2.2	Chain Rule of Multivariable Functions	4
2.3	Chain Rule via Tree Diagram	4
2.4	Directional Derivative	5
3	Section Fifteen	6
3.1	Riemann Sum in \mathbb{R}^3	6
3.2	Double Integrals	6
3.3	Applications of Double Integrals: Lamina	7
3.4	Integrals and Probability	7
3.5	Joint Probability	7
3.6	Applications of Double Integrals: Joint Probability	7
3.7	Examples of Probability Density Functions	8
3.8	Applications of Double Integrals: Surface Area	8
3.9	Triple Integrals	9
3.10	Applications of Triple Integrals: Solids	9
3.11	Applications of Triple Integrals: Joint Probability	10
3.12	Cylindrical Coordinates for \mathbb{R}^3	10
3.13	Triple Integrals in Cylindrical Coordinates	10
3.14	Spherical Coordinates for \mathbb{R}^3	10
3.15	Triple Integrals in Spherical Coordinates	10
3.16	Change of Variables in Multiple Integrals	11
4	Section Sixteen	12
4.1	Vector Fields	12
4.2	Conservative Vector Fields	12
4.3	Fundamental Theorem of Line Integrals	12
4.4	Open Regions	13
4.5	Closed Regions	13
4.6	Connected Regions	13
4.7	Simply Connected Regions	13
4.8	Unifying Theorems	13
4.9	Boundary Curve Orientation in \mathbb{R}^2	13
4.10	Green's Theorem	14
4.11	Curl of a Vector Field	14
4.12	Divergence of a Vector Field	15
4.13	Curve Parametrization	15
4.14	Surface Parametrization	15
4.15	Surface Parametrization of Plane	16
4.16	Surface Parametrization of Cylinder	16
4.17	Surface Parametrization of Revolution	16

4.18 Surface Parametrization of Sphere	16
4.19 Surface Area of Parametrized Surface	16
4.20 Lamina of Parametrized Surface	16
4.21 Orientation of Surfaces	17
4.22 Tangent Plane of Parametrized Surface	17
4.23 Unit Normal Vector	17
4.24 Surface Integral of a Vector Field	18
4.25 Boundary Curve Orientation in \mathbb{R}^3	18
4.26 Stokes' Theorem	18
4.27 Gauss' Theorem (Divergence Theorem)	19

1 General

1.1 Trigonometric Identities

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\cos^2 \theta = \frac{1}{2}(1 + \cos 2\theta)$$

$$\sin^2 \theta = \frac{1}{2}(1 - \cos 2\theta)$$

1.2 Polar Coordinates



$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$r = \sqrt{x^2 + y^2}$$

$$\theta = \tan^{-1}\left(\frac{y}{x}\right)$$

1.3 Polar Curves



1.4 Curve Parametrization

$$\vec{r}(t) = \langle x(t), y(t), z(t) \rangle$$

$$\vec{r}'(t) = \left\langle \left(\frac{dx}{dt}\right)^2, \left(\frac{dy}{dt}\right)^2, \left(\frac{dz}{dt}\right)^2 \right\rangle$$

$$s = \int_a^b |\vec{r}'(t)| \, dt = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} \, dt$$

$$\frac{ds}{dt} = |\vec{r}'(t)| = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2}$$

$$ds = |\vec{r}'(t)| \, dt = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} \, dt$$

2 Section Fourteen

2.1 Chain Rule of Single Variable Functions

Given that $y = f(x)$, if $x = x(t)$ and $y = f(x(t))$ then

$$\text{if } x = x(t), y = f(x(t)) \xrightarrow{\text{chain rule}} \frac{dy}{dt} = \frac{dy}{dx} \times \frac{dx}{dt}$$


2.2 Chain Rule of Multivariable Functions

Given that $z = f(x, y)$, if $x = x(t)$, $y = y(t)$ and $z = f(x, y)$ then


$$\begin{aligned} \text{if } \begin{cases} x = x(t) \\ y = y(t) \end{cases} &\xrightarrow{\text{chain rule}} \frac{dz}{dt} = \frac{dz}{dx} + \frac{dz}{dy} \\ &= \frac{\partial z}{\partial x} \times \frac{dx}{dt} + \frac{\partial z}{\partial y} \times \frac{dy}{dt} \end{aligned}$$

2.3 Chain Rule via Tree Diagram

Multiply the partial derivatives along each branch, then sum up along all branches



$$\frac{dz}{dt} = \left[\frac{\partial z}{\partial x} \times \frac{dx}{dt} \right] + \left[\frac{\partial z}{\partial y} \times \frac{dy}{dt} \right]$$



$$\frac{dz}{dt} = \left[\frac{\partial z}{\partial x} \times \frac{dx}{du} + \frac{\partial z}{\partial x} \times \frac{dx}{dv} \right] + \left[\frac{\partial z}{\partial y} \times \frac{dy}{du} + \frac{\partial z}{\partial y} \times \frac{dy}{dv} \right]$$

2.4 Directional Derivative

$D_{\vec{u}}f(x, y)$ is the rate of change of z along the direction of the unit vector $\vec{u} = \langle a, b \rangle$

$$\begin{aligned} D_{\vec{u}}f(x, y) &= \nabla f(x, y) \cdot \vec{u} \\ &= \left\langle \frac{\partial f}{\partial x}(x, y), \frac{\partial f}{\partial y}(x, y) \right\rangle \cdot \vec{u} \end{aligned}$$

The gradient vector $\nabla f(x, y)$ maximizes the directional derivative of $f(x, y)$ and $\nabla f(x, y) \perp$ level curves

$$\begin{aligned} D_{\vec{u}}f(x, y) &= \nabla f(x, y) \cdot \vec{u} \longleftarrow \vec{u} \text{ is a unit vector} \\ &= |\nabla f(x, y)| |\vec{u}| \cos \theta \longleftarrow 0 \leq \theta \leq \pi \\ &= |\nabla f(x, y)| \cos \theta \begin{cases} \text{is maximized at } \theta = 0, & D_{\vec{u}}f(x, y) = \nabla f(x, y) \\ \text{is minimized at } \theta = \pi, & D_{\vec{u}}f(x, y) = -\nabla f(x, y) \\ \text{is 0 at } \theta = \frac{\pi}{2}, & D_{\vec{u}}f(x, y) = 0 \end{cases} \end{aligned}$$

Properties of the gradient vector

- $\nabla(f \pm g) = \nabla f \pm \nabla g$
- $\nabla(kf) = k \cdot \nabla f$
- $\nabla(fg) = g * \nabla f + f * \nabla g$
- $\nabla\left(\frac{f}{g}\right) = \frac{g * \nabla f - f * \nabla g}{g^2}$

3 Section Fifteen

3.1 Riemann Sum in \mathbb{R}^3

$$\partial x = \frac{b-a}{m}$$

$$\partial y = \frac{d-c}{n}$$

$$\partial A = \partial x \partial y$$

Sample point: (x_i^*, y_j^*)

$$V_{ij} \approx f(x_i^*, y_j^*) \cdot \partial A$$


$$V \approx \sum_{j=1}^n \sum_{i=1}^m V_{ij}$$

3.2 Double Integrals

Represents the volume that lies under a surface $f(x, y)$ and above the area D


Type I iterated integral

$$\iint_D f(x, y) \, dy \, dx$$




Type II iterated integral

$$\iint_D f(x, y) \, dx \, dy$$



Polar iterated integral

$$\iint_D f(x, y)(r) \, dr \, d\theta$$



3.3 Applications of Double Integrals: Lamina

Given a density function $\rho(x, y)$

$$m = \iint_D \rho(x, y) \, dA$$

$$\bar{x} = \frac{1}{m} M_y = \frac{1}{m} \iint_D x \rho(x, y) \, dA \longrightarrow M_y = \text{moment about } y\text{-axis}$$

$$\bar{y} = \frac{1}{m} M_x = \frac{1}{m} \iint_D y \rho(x, y) \, dA \longrightarrow M_x = \text{moment about } x\text{-axis}$$

$$I_x = \iint_D y^2 \rho(x, y) \, dA \longrightarrow I_x = \text{moment of inertia about } x\text{-axis}$$

$$I_y = \iint_D x^2 \rho(x, y) \, dA \longrightarrow I_y = \text{moment of inertia about } y\text{-axis}$$

$$\begin{aligned} I_o &= \iint_D r^2 \rho(x, y) \, dA \longrightarrow I_o = \text{moment of inertia about the origin} \\ &= I_x + I_y \end{aligned}$$

3.4 Integrals and Probability

Given a random variable x with probability density function $f(x)$

$$f(x) \geq 0 \longrightarrow P(X) \text{ cannot be smaller than } 0$$

$$\int_{-\infty}^{\infty} f(x) \, dx = 1 \longrightarrow \text{total probability is } 1$$

$$P(a \leq X \leq b) = \int_a^b f(x) \, dx$$

$$\bar{x} = \int_{-\infty}^{\infty} x f(x) \, dx$$

3.5 Joint Probability

The joint probability density function $f(x, y)$ represents the probability of both x and y occurring together

- x and y are independent if their joint probability density function is equal to the product of their individual probability density functions
- $f(x, y) = f_1(x)f_2(y) \Leftrightarrow P(x \cap y) = 0$

3.6 Applications of Double Integrals: Joint Probability

Given random variables x, y with joint probability density function $f(x, y)$

$$f(x, y) \geq 0 \longrightarrow P(X, Y) \text{ cannot be smaller than } 0$$


$$\iint_{\mathbb{R}^2} f(x, y) \, dA = 1 \longrightarrow \text{total probability is } 1$$

$$P((X, Y) \in D) = P(a \leq X \leq b, c \leq Y \leq d) = \iint_D f(x, y) \, dA$$

3.7 Examples of Probability Density Functions

Exponential probability density function


$$f(x) = \begin{cases} \frac{1}{\lambda} e^{-\frac{x}{\lambda}} & x \geq 0 \\ 0 & x < 0 \end{cases}$$



- Used when a random variable x has a mean value equal to λ

Normal distribution probability density function

$$f(x) = \frac{1}{\sqrt{\pi}} \underbrace{e^{-x^2}}_{\text{Gaussian function}}$$



- Used when a random variable x has a mean value equal to λ

3.8 Applications of Double Integrals: Surface Area

$$\vec{a} = \langle \partial x, 0, f_x(x_i, y_i) \partial x \rangle$$

$$\vec{b} = \langle 0, \partial y, f_y(x_i, y_i) \partial y \rangle$$

$$S_{ij} \approx |\vec{a} \times \vec{b}|$$

$$\approx \langle -f_x(x, y) \cdot \partial x \partial y, -f_y(x, y) \cdot \partial x \partial y, \partial x \partial y \rangle$$

$$\approx \langle -f_x(x, y), -f_y(x, y), 1 \rangle \partial x \partial y$$

$$S = \lim_{\substack{m \rightarrow \infty \\ n \rightarrow \infty}} \sum_{j=1}^n \sum_{i=1}^m S_{ij}$$

$$S = \iint_D \sqrt{[f_x(x, y)]^2 + [f_y(x, y)]^2 + 1} \, dA$$


$$= \iint_D (r) \sqrt{[f_x(x, y)]^2 + [f_y(x, y)]^2 + 1} \, dr d\theta$$

3.9 Triple Integrals

Represents the 4D object that lies under a surface $f(x, y, z)$ and above the 3D solid E


Type I triple integral

$$\iint_D \left(\int_{u_1}^{u_2} f(x, y, z) dz \right) dA$$




Type II triple integral

$$\iint_D \left(\int_{u_1}^{u_2} f(x, y, z) dx \right) dA$$



Type III triple integral

$$\iint_D \left(\int_{u_1}^{u_2} f(x, y, z) dy \right) dA$$



3.10 Applications of Triple Integrals: Solids

Given density function $\rho(x, y, z)$

$$m = \iiint_E \rho(x, y, z) dV$$

$$\bar{x} = \frac{1}{m} \iiint_E x \rho(x, y, z) dV$$

$$\bar{y} = \frac{1}{m} \iiint_E y \rho(x, y, z) dV$$

$$\bar{z} = \frac{1}{m} \iiint_E z \rho(x, y, z) dV$$

$$I_x = \iiint_E (y^2 + z^2) \rho(x, y, z) dV \longrightarrow I_x = \text{moment of inertia about } x\text{-axis}$$

$$I_y = \iiint_E (x^2 + z^2) \rho(x, y, z) dV \longrightarrow I_y = \text{moment of inertia about } y\text{-axis}$$

$$I_z = \iiint_E (x^2 + y^2) \rho(x, y, z) dV \longrightarrow I_z = \text{moment of inertia about } z\text{-axis}$$

3.11 Applications of Triple Integrals: Joint Probability

Given random variables x, y, z with joint probability density function $f(x, y, z)$

$$f(x, y, z) \geq 0 \longrightarrow P(X, Y, Z) \text{ cannot be smaller than } 0$$


$$\iiint_{\mathbb{R}^3} f(x, y, z) dV = 1 \longrightarrow \text{total probability is } 1$$

$$\begin{aligned} P((X, Y, Z) \in E) &= P(a \leq X \leq b, c \leq Y \leq d, e \leq Z \leq f) \\ &= \iiint_E f(x, y, z) dV \end{aligned}$$

3.12 Cylindrical Coordinates for \mathbb{R}^3

Points in \mathbb{R}^3 are located by (r, θ, z)

x, y converted to Polar coordinates



3.13 Triple Integrals in Cylindrical Coordinates

$$\iiint_E f(x, y, z) dV = \iiint_D \int_{u_1}^{u_2} f(r \cos \theta, r \sin \theta, z) \cdot r \, dz dr d\theta$$

3.14 Spherical Coordinates for \mathbb{R}^3

Points in \mathbb{R}^3 are located by (ρ, θ, ϕ)

$$r = OP' = \rho \sin \phi$$

$$x = r \cos \theta = \rho \sin \phi \cos \theta$$

$$y = r \sin \theta = \rho \sin \phi \sin \theta$$


$$z = \rho \cos \phi$$

$$\rho^2 = x^2 + y^2 + z^2$$

$$\rho^2 = r^2 + z^2$$

$$\tan \theta = \frac{y}{x}$$

$$\tan \phi = \frac{r}{z}$$



3.15 Triple Integrals in Spherical Coordinates

$$\iiint_E f(x, y, z) dV = \iiint_{g_1}^{g_2} f(\rho \sin \phi \cos \theta, \rho \sin \phi \sin \theta, \rho \cos \phi) \cdot \rho^2 \sin \phi \, d\rho d\theta d\phi$$

3.16 Change of Variables in Multiple Integrals

A change of variable is a transformation function $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$

$$\iint_{\substack{R \text{ in } xy \text{ plane}}} f(x, y) \, dA = \iint_{\substack{S \text{ in } uv \text{ plane}}} f(x(u, v), y(u, v)) \left| \det \left(\begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{bmatrix} \right) \right| \, dudv$$

$$\frac{\partial(x, y)}{\partial(u, v)} = \det \left(\begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{bmatrix} \right) \leftarrow \text{Jacobian of the transformation}$$

$$\iiint_{\substack{R \text{ in } xyz \text{ plane}}} f(x, y, z) \, dV = \iiint_{\substack{S \text{ in } uv t \text{ plane}}} f(x(u, v, t), y(u, v, t), z(u, v, t)) \left| \det \left(\begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial t} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial t} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial t} \end{bmatrix} \right) \right| \, dudvdt$$

$$\frac{\partial(x, y, z)}{\partial(u, v, t)} = \det \left(\begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial t} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial t} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial t} \end{bmatrix} \right) \leftarrow \text{Jacobian of the transformation}$$

4 Section Sixteen

4.1 Vector Fields

A vector field \vec{F} is a vector-valued function that assigns any point in the domain $D \subseteq \mathbb{R}^n$ an n dimensional vector

2D vector field

$$\vec{F}(x, y) = \langle P(x, y), Q(x, y) \rangle$$

3D vector field

$$\vec{F}(x, y, z) = \langle P(x, y, z), Q(x, y, z), R(x, y, z) \rangle$$

4.2 Conservative Vector Fields

A vector field \vec{F} is conservative if \vec{F} is the vector gradient of a function f . In this case we call the function f a potential function of the vector field \vec{F}

- Any vector field $\vec{F}(x, y) = \nabla f$ is conservative
- If D is an open connected region and \vec{F} is a vector field that is continuous in D , then \vec{F} is conservative if and only if the line integral of \vec{F} is path-independent in D

2D conservative vector field $\vec{F} = \langle P(x, y), Q(x, y) \rangle$

$$\vec{F}(x, y) \text{ is conservative if and only if } \frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$$

3D conservative vector field $\vec{F} = \langle P(x, y, z), Q(x, y, z), R(x, y, z) \rangle$

$$\vec{F}(x, y, z) \text{ is conservative if and only if } \frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x} \text{ and } \frac{\partial Q}{\partial z} = \frac{\partial R}{\partial y} \text{ and } \frac{\partial R}{\partial x} = \frac{\partial P}{\partial z}$$

4.3 Fundamental Theorem of Line Integrals

$$\int_C \nabla f \, d\vec{r} = \int_C \nabla f \cdot \vec{r}'(t) \, dt = \underbrace{f(\vec{r}(b))}_{\text{ending pt. B}} - \underbrace{f(\vec{r}(a))}_{\text{starting pt. A}}$$

$$\int_C \nabla f \, d\vec{r} = \int_C P \, dx + Q \, dy$$

- If \vec{F} is a conservative vector field, then the line integral of \vec{F} along any curve C is path-independent
- If \vec{F} is a conservative vector field, then the line integral of \vec{F} is equal to 0 along any closed curve

4.4 Open Regions

A region D in \mathbb{R}^n is open if for any point P in D , there exists a disk centered at P that lies entirely within the borders of D

- D is open if it does not contain any boundary points
- The region is defined by an inequality containing only $<$ and $>$
- i.e. $\{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 4\}$

4.5 Closed Regions

A region D in \mathbb{R}^n is closed if its complement $\mathbb{R}^n - D$ is open

- D is closed if it contains boundary points
- The region is defined by an inequality containing \leq or \geq
- i.e. $\{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 4\}$

4.6 Connected Regions

A region D in \mathbb{R}^n is connected if any two points in D can be joined by a path that lies in D

4.7 Simply Connected Regions

A region D in \mathbb{R}^n is simply-connected if D is connected, and the region enclosed by any closed curve C in D is contained entirely within D

- Does not contain a hole

4.8 Unifying Theorems

- If D is an open connected region and \vec{F} is a vector field that is continuous on D , then \vec{F} is conservative if and only if the line integral of \vec{F} is path-independent in D

4.9 Boundary Curve Orientation in \mathbb{R}^2

- The boundary curve(s) C of a 2D region D ($\partial D = C_1 \cup C_2 \dots$) is positively oriented if C is traversed in a way that the region D is always on the left side of C
- The boundary curve(s) C of a 2D region D ($\partial D = C_1 \cup C_2 \dots$) is negatively oriented if C is traversed in a way that the region D is always on the right side of C

4.10 Green's Theorem

Let C be a positively oriented simple closed piecewise-smooth curve in \mathbb{R}^2 that bounds the region D

Let $\vec{F}(x, y) = \langle P(x, y), Q(x, y) \rangle$ be a 2D vector field where $P(x, y)$ and $Q(x, y)$ both have continuous partial derivatives on an open region containing D

Then $\int_C \vec{F} d\vec{r} = \iint_D \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} dA$ (multiply by -1 if C is negatively oriented)

Uses:

- Alternative way to calculate line integrals
- Calculate the area of a region

4.11 Curl of a Vector Field

Given a 3D vector field $\vec{F} = \langle P(x, y, z), Q(x, y, z), R(x, y, z) \rangle$, the curl of \vec{F} is the vector field:

$$\begin{aligned} \text{rot} \vec{F} &= \nabla \times \vec{F} \\ &= \left\langle \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right\rangle \times \langle P(x, y, z), Q(x, y, z), R(x, y, z) \rangle \\ &= \left\langle \frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z}, \frac{\partial P}{\partial z} - \frac{\partial R}{\partial x}, \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right\rangle \end{aligned}$$

- The curl of \vec{F} measures the rotation of the vector field \vec{F}
 - Thinking of the velocity field as a fluid flow, $\text{rot} \vec{F}$ measures the rotation of an object caused by the fluid
- Magnitude of the curl of \vec{F} measures the intensity of rotation
- Direction of the curl of \vec{F} gives the axis of rotation by the right-hand grip rule
- To detect the rotational effect of a 2D vector field $\vec{F} = \langle P(x, y), Q(x, y) \rangle$, we may extend the 2D vector field to $\vec{F} = \langle P(x, y), Q(x, y), 0 \rangle$
- The curl of \vec{F} is zero if and only if \vec{F} is conservative

4.12 Divergence of a Vector Field

Given a vector field \vec{F} , the divergence of \vec{F} is the scalar function:

$$\begin{aligned}\operatorname{div} \vec{F} &= \nabla \cdot \vec{F} \\ &= \left\langle \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right\rangle \cdot \langle P(x, y, z), Q(x, y, z), R(x, y, z) \rangle \\ &= \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}\end{aligned}$$

- The divergence of \vec{F} measures the "outgoing-ness" of \vec{F}
 - Thinking of the velocity field as a fluid flow, $\operatorname{div} \vec{F}$ measures the net rate at which the fluid flows through a point
 - If $\operatorname{div} \vec{F} < 0$, there is more fluid going in than out
 - If $\operatorname{div} \vec{F} > 0$, there is more fluid going out than in
- Applies to 2D and 3D vector fields
- If $\operatorname{div} \vec{F} = 0$, then \vec{F} is incompressible
- $\operatorname{div}(\operatorname{curl} \vec{F})$ is always zero
 - Can be used to check if a curl is valid

4.13 Curve Parametrization

A curve C in \mathbb{R}^3 can be parametrized by a vector function of one parameter

$$C : \vec{r}(t) = \langle x(t), y(t), z(t) \rangle \quad t \in [a, b] \subseteq \mathbb{R}$$

- Consists of one parameter t
- C represents a 1D object in \mathbb{R}^3

4.14 Surface Parametrization

A surface S in \mathbb{R}^3 can be parametrized by a vector function of two parameters

$$S : \vec{r}(u, v) = \langle x(u, v), y(u, v), z(u, v) \rangle \quad (u, v) \in D \subseteq \mathbb{R}^2$$

- Consists of two parameters u, v
- S represents a 2D object in \mathbb{R}^3

4.15 Surface Parametrization of Plane

Plane through point $\vec{r}_0 = \langle x_0, y_0, z_0 \rangle$ and is spanned by vectors \vec{a} and \vec{b}

$$\vec{r}(u, v) = r_0 + u\vec{a} + v\vec{b}$$

4.16 Surface Parametrization of Cylinder

Cylinder centered at origin with radius r and axis of rotation z

$$\vec{r}(\theta, z) = \langle r \cos \theta, r \sin \theta, z \rangle$$

4.17 Surface Parametrization of Revolution

Revolution centered at origin with radius $f(z)$ and axis of rotation z

$$\vec{r}(\theta, z) = \langle f(z) \cos \theta, f(z) \sin \theta, z \rangle$$

4.18 Surface Parametrization of Sphere

Sphere centered at origin with radius ρ

$$\vec{r}(\phi, \theta) = \langle \rho \sin \phi \cos \theta, \rho \sin \phi \sin \theta, \rho \cos \phi \rangle$$

4.19 Surface Area of Parametrized Surface

Given $S : \vec{r}(u, v) = \langle x(u, v), y(u, v), z(u, v) \rangle$

$$dS = |\vec{r}_u \times \vec{r}_v| dA \quad \leftarrow \text{if } S \text{ is parametrized by } \vec{r}(u, v)$$

$$dS = \sqrt{f_x^2 + f_y^2 + 1} dA \quad \leftarrow \text{if } S \text{ is the graph } z = f(x, y)$$

$$\iint_S f(x, y, z) ds = \iint_D f(\vec{r}(u, v)) |\vec{r}_u \times \vec{r}_v| dA$$

$$\iint_S f(x, y, z) ds = \iint_D f(\vec{r}(u, v)) \sqrt{f_x^2 + f_y^2 + 1} dA$$

4.20 Lamina of Parametrized Surface

Given density function $\rho(x, y, z)$

$$m = \iint_S \rho(x, y, z) dS$$

$$\bar{x} = \frac{1}{m} \iint_S x \rho(x, y, z) dS$$

$$\bar{y} = \frac{1}{m} \iint_S y \rho(x, y, z) dS$$

$$\bar{z} = \frac{1}{m} \iint_S z \rho(x, y, z) dS$$

4.21 Orientation of Surfaces

A surface S is orientable if it has two sides

- If S is orientable, the tangent plane at any point on S has two unit normal vectors \vec{n}_1 and \vec{n}_2 where $\vec{n}_1 = -\vec{n}_2$
- The orientation of an orientable surface S is a choice between the two unit normal vectors of tangent plane at each point of the surface so that the choice varies continuously over S
- If S is a closed surface bounding solid E , S is positively oriented if \vec{n} is chosen to be pointing outward from E , and negatively oriented if \vec{n} is chosen to be pointing inward to E

4.22 Tangent Plane of Parametrized Surface

Given $S : \vec{r}(u, v) = \langle x(u, v), y(u, v), z(u, v) \rangle$

$$\left. \begin{aligned} \vec{r}_u &= \left\langle \frac{\partial x}{\partial u}, \frac{\partial y}{\partial u}, \frac{\partial z}{\partial u} \right\rangle \\ \vec{r}_v &= \left\langle \frac{\partial x}{\partial v}, \frac{\partial y}{\partial v}, \frac{\partial z}{\partial v} \right\rangle \end{aligned} \right\} \text{tangent vectors}$$

$$\vec{n} = \vec{r}_u \times \vec{r}_v \quad \leftarrow \text{normal vector } \vec{n} \text{ of tangent plane}$$

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n} \quad \leftarrow \text{equation of tangent plane, given known point } \vec{a}$$

4.23 Unit Normal Vector

If S is parametrized by $\vec{r}(u, v)$ then

$$\hat{n} = \pm \frac{\vec{r}_u \times \vec{r}_v}{|\vec{r}_u \times \vec{r}_v|} \quad \leftarrow \pm \text{ depends on the desired orientation}$$

If S is the graph of $z = g(x, y)$ then

$$\hat{n} = \pm \frac{\langle -g_x, -g_y, 1 \rangle}{\sqrt{g_x^2 + g_y^2 + 1}} \quad \leftarrow \pm \text{ depends on the desired orientation}$$

4.24 Surface Integral of a Vector Field

The surface integral of \vec{F} over an orientable surface S with unit normal vector \hat{n} is

$$\iint_S \vec{F} \cdot d\vec{S} = \iint_S \vec{F} \cdot \hat{n} \, dS \quad \leftarrow \begin{array}{l} \text{represents the flux of } \vec{F} \text{ across } S, \\ \text{the volume of fluid flowing through } S \text{ per time unit} \end{array}$$

If S is parametrized by $\vec{r}(u, v)$ then

$$\begin{aligned} \iint_S \vec{F} \cdot d\vec{S} &= \iint_D (\vec{F} \cdot \pm \frac{\vec{r}_u \times \vec{r}_v}{|\vec{r}_u \times \vec{r}_v|}) |\vec{r}_u \times \vec{r}_v| \, dA \\ &= \iint_D \vec{F} \cdot \pm (\vec{r}_u \times \vec{r}_v) \, dA \end{aligned}$$

If S is the graph of $z = g(x, y)$ then

$$\begin{aligned} \iint_S \vec{F} \cdot d\vec{S} &= \iint_D (\vec{F} \cdot \pm \frac{\langle -g_x, -g_y, 1 \rangle}{\sqrt{g_x^2 + g_y^2 + 1}}) \sqrt{g_x^2 + g_y^2 + 1} \, dA \\ &= \iint_D \vec{F} \cdot \pm \langle -g_x, -g_y, 1 \rangle \, dA \end{aligned}$$

4.25 Boundary Curve Orientation in \mathbb{R}^3

- The boundary curve(s) C of a surface S ($\partial S = C_1 \cup C_2 \dots$) is positively oriented if C is traversed in a way that the surface S is always on the left side of C from the perspective in the same direction of the normal vector of the surface
 - By the right hand rule, if the thumb is pointed towards \vec{n} , the fingers are curling along the orientation of C
- The boundary curve(s) C of a surface S ($\partial S = C_1 \cup C_2 \dots$) is negatively oriented if C is traversed in a way that the surface S is always on the right side of C from the perspective in the same direction of the normal vector of the surface

4.26 Stokes' Theorem

Let S be an orientable piecewise-smooth surface whose boundary C is a positively oriented simple closed piecewise-smooth curve

Let $\vec{F}(x, y, z) = \langle P(x, y, z), Q(x, y, z), R(x, y, z) \rangle$ be a 3D vector field where $P(x, y, z)$, $Q(x, y, z)$ and $R(x, y, z)$ each have continuous partial derivatives on an open region containing S

Then $\int_C \vec{F} \cdot d\vec{r} = \iint_S \text{curl} \vec{F} \cdot d\vec{S}$ (multiply by -1 if C is negatively oriented)

4.27 Gauss' Theorem (Divergence Theorem)

Let E be a 3D solid that is bounded by a closed surface S where S is positively oriented

Let $\vec{F}(x, y, z) = \langle P(x, y, z), Q(x, y, z), R(x, y, z) \rangle$ be a 3D vector field where $P(x, y, z)$, $Q(x, y, z)$ and $R(x, y, z)$ each have continuous partial derivatives on an open region containing E

$$\text{Then } \iint_S \vec{F} \cdot d\vec{S} = \iiint_E \text{div} \vec{F} \, dV$$