

MATH 402 Notes

Contents

1	Arithmetic in \mathbb{Z} Revisited	3
1.1	Division Theorem	3
1.2	Divisibility	3
1.3	Greatest Common Divisors	3
1.4	Euclid's Lemma	3
1.5	Diophantine Equations	3
1.6	Fundamental Theorem of Arithmetics	3
2	Congruence in \mathbb{Z} and Modular Arithmetic	4
2.1	Congruence	4
2.2	Congruence Class of $a \bmod n$	4
2.3	Additive Operations With Congruence Classes	4
2.4	Multiplicative Operations With Congruence Classes	4
2.5	Units in \mathbb{Z}_n	4
2.6	Zero Divisor in \mathbb{Z}_n	4
2.7	Multiplicative Inverse in \mathbb{Z}_n	4
3	Rings	5
3.1	Rings	5
3.2	Commutative Rings	5
3.3	Rings With Identity	5
3.4	Fields	5
3.5	Integral Domains	5
3.6	Units in Rings	5
3.7	Zero Divisor in Rings	5
3.8	Multiplicative Inverse in Rings	6
3.9	Subrings	6
3.10	Subring Theorem	6
3.11	Subring Set Theory	6
3.12	Finite Set $\mathbb{F}_p[\theta]$	6
3.13	Complex Set \mathbb{C}	6
3.14	Gaussian Integers $\mathbb{Z}[i]$	7
3.15	Matrices $M_2(\mathbb{F})$	7
3.16	Units in $M_2(\mathbb{F})$	7
3.17	Continuous Real-Valued Functions $C[0, 1]$	7
3.18	Ring Homomorphisms	8
3.19	Ring Isomorphisms	8
4	Arithmetic in $\mathbb{F}[x]$	9
4.1	Polynomials	9
4.2	Polynomial Rings	9
4.3	Finite Polynomial Rings	9
4.4	Division Theorem in Polynomials	10
4.5	Divisibility in Polynomials	10
4.6	Greatest Common Divisors in Polynomials	10
4.7	Bézout's Theorem for Polynomials	10
4.8	Polynomial Roots	10
4.9	Associates	10

4.10	Non-Trivial Factorization	10
4.11	Reducible Polynomials	11
4.12	Irreducible Polynomials	11
4.13	Theorems on Irreducible Polynomials	11
4.14	Rational Root Test	11
4.15	Gauss' Lemma	11
4.16	Polynomial Reduction $\text{mod } n$	12
4.17	Checking Irreducibility	12
4.18	Eisenstein's Criterion	12
4.19	Fundamental Theorem of Algebra	12
4.20	Polynomials in $\mathbb{R}[x]$	12
4.21	Roots of Unity	13
5	Congruence in $\mathbb{F}[x]$ and Congruence-Class Arithmetic	14
5.1	Polynomial Congruence	14
5.2	Polynomial Congruence Class of $f(x) \text{ mod } p(x)$	14
5.3	Polynomial Congruence Ring $\mathbb{F}[x]_{p(x)}$	14
5.4	Fields in $\mathbb{F}[x]_{p(x)}$	14
5.5	Finite Fields in $\mathbb{F}_p[x]_{p(x)}$	14
6	Ideals and Quotient Rings	15
6.1	Ideals	15
6.2	Maximal Ideals	15
6.3	Finitely Generated Ideals	15
6.4	Principal Ideals Generated by c	15
6.5	Principal Ideal Domains	15
6.6	Ideals in \mathbb{Z}	16
6.7	Ideals in $\mathbb{Z}[i]$	16
6.8	Ideal Lattice	16
6.9	Ideal Congruence	16
6.10	Ideal Congruence Class of $a \text{ mod } I$	17
6.11	Lagrange's Theorem for Finite Rings	17
6.12	Quotient Ring R/I	17
6.13	Kernels	17
6.14	Natural Homomorphisms	17
6.15	First Isomorphism Theorem	18
6.16	Product Decomposition Theorem	18
6.17	Product Decomposition Theorem for Polynomials	18
6.18	Additional Theorems	18
6.19	Prime Ideals	18
6.20	Maximal and Prime Ideals	18
7	Arithmetic in $\mathbb{Z}[i]$	19
7.1	Division Theorem in Gaussian Integers	19
7.2	Divisibility in Gaussian Integers	19
7.3	Greatest Common Divisors in Gaussian Integers	19
7.4	Bézout's Theorem for Gaussian Integers	19
7.5	Non-Trivial Factorization	19
7.6	Unique Factorization Theorem	19
7.7	Gaussian Primes	20
8	Appendix	21
8.1	Rings	21
8.2	Norm Functions	21
8.3	Quick Proofs	22

1 Arithmetic in \mathbb{Z} Revisited

1.1 Division Theorem

Given non-zero integers a and b where $b > 0$, there exists unique integers q and r such that $a = bq + r$ where $0 \leq r < b$

1.2 Divisibility

Given $a, b \in \mathbb{Z}$, $b \mid a$ means there exists $c \in \mathbb{Z}$ such that $a = bc$

1.3 Greatest Common Divisors

Given non-zero integers a and b , $\gcd(a, b) = (a, b)$ is the largest common divisor of a and b

- Finding $\gcd(a, b)$ by brute force

Let $\mathcal{D}_a = \{\text{divisors of } a\}$ and $\mathcal{D}_b = \{\text{divisors of } b\}$. Then $\mathcal{D}_a \cap \mathcal{D}_b = \{\text{common divisors of } a \text{ and } b\}$. Hence, $\gcd(a, b) = \max(\mathcal{D}_a \cap \mathcal{D}_b)$

- Finding $\gcd(a, b)$ by the Euclidean algorithm

Lemma 1.3.1. If $b \mid a$, then $\gcd(a, b) = |b|$

Lemma 1.3.2. If $x = yz + w$, then $\gcd(x, y) = \gcd(y, w)$

Given non-zero integers a and b , the recursive Euclidean Algorithm computes $\gcd(a, b)$

A full definition of the Euclidean Algorithm can be found in **MATH 300 Notes**, page 12

1.4 Euclid's Lemma

Assume a, b, c are non-zero integers. If $\gcd(a, b) = 1$ and $a \mid bc$, then $a \mid c$

1.5 Diophantine Equations

Any equation of the form $ax + by = c$ where $a, b, c \in \mathbb{Z}$ is called a linear Diophantine equation

- Has no solutions if $\gcd(a, b)$ does not divide c
- Has infinitely many solutions if $\gcd(a, b)$ divides c
- Given a solution where $x = x_0$, $y = y_0$, and $d = \gcd(a, b)$

$$x = x_0 + \frac{b}{d}k$$

$$y = y_0 - \frac{a}{d}k$$

1.6 Fundamental Theorem of Arithmetics

Given a positive integer $n > 1$ with two prime factorizations

$$n = p_1 p_2 \dots p_r$$

$$n = q_1 q_2 \dots q_s$$

Then $r = s$ and $\{p_1, p_2, \dots, p_r\} = \{q_1, q_2, \dots, q_s\}$. This means that the prime factorization of n is unique

2 Congruence in \mathbb{Z} and Modular Arithmetic

2.1 Congruence

Given two integers a and b and a modulus $m \geq 1$, a is congruent to $b \bmod m$ if and only if $m \mid a - b$. This is denoted as $a \equiv b \bmod m$

- Congruence is reflexive, symmetric and transitive

2.2 Congruence Class of $a \bmod n$

The congruence class of $a \bmod n$, denoted $[a]_n$, is the set of all integers that are congruent to $a \bmod n$

- $[a]_n = \{x \mid x \equiv a \bmod n\}$
- There are n congruence classes in \mathbb{Z}_n , $[0]_n [1]_n \dots [n-1]_n$
- Congruence classes are either equal or disjoint
- a is typically the least residue $\bmod n$

2.3 Additive Operations With Congruence Classes

- Behaves the same as integer addition
- $[a]_n + [b]_n = [a + b]_n$

2.4 Multiplicative Operations With Congruence Classes

- Behaves the same as integer multiplication
- $[a]_n \cdot [b]_n = [a \cdot b]_n$

2.5 Units in \mathbb{Z}_n

a is a unit in \mathbb{Z}_n if the equation $ax \equiv 1 \bmod n$ has a solution

- Has the associated linear Diophantine equation $ax + ny = 1$
- a is a unit in \mathbb{Z}_n if and only if $\gcd(a, n) = 1$

2.6 Zero Divisor in \mathbb{Z}_n

a is a zero divisor in \mathbb{Z}_n if $a \neq 0$ and the equation $ax \equiv 0 \bmod n$ has a non-zero solution for some $x \in \mathbb{Z}_n$

- \mathbb{Z}_n is a disjoint union of $\{0\} \cup \{\text{units}\} \cup \{\text{zero divisors}\}$
- If a is not 0 or a unit, then a is a zero divisor

2.7 Multiplicative Inverse in \mathbb{Z}_n

a is invertible in \mathbb{Z}_n if and only if $ax \equiv 1 \bmod n$ has integer solutions

- a is invertible if and only if $\gcd(a, n) = 1$
- x is the inverse of a , denoted a^{-1}
- Given p is prime, $a^{-1} \equiv a^{p-2} \bmod p$

3 Rings

3.1 Rings

A ring is a nonempty set R that can undergo two operations, usually written as addition and multiplication

- Additive operations satisfy the following axioms
 1. Closed under addition: if $a \in R$ and $b \in R$, then $a + b \in R$
 2. Associative: $a + (b + c) = (a + b) + c$
 3. Commutative: $a + b = b + c$
 4. Additive identity: there exists an element $0_R \in R$ such that $a + 0_R = a$ for all a
 5. Additive inverse: for each a , there exists an element $x \in R$ such that $a + x = 0_R$
- Multiplicative operations satisfy the following axioms
 6. Closed under multiplication: if $a \in R$ and $b \in R$, then $ab \in R$
 7. Associative: $a(bc) = (ab)c$
 8. Distributive: $a(b + c) = ab + ac$ and $(a + b)c = ac + bc$

Multiplicative operations are not necessarily commutative, i.e. $ab \neq ba$

Multiplicative operations do not necessarily have a multiplicative identity, i.e. $a1_R = 1_Ra = a$ for all a

3.2 Commutative Rings

A commutative ring is a ring R in which multiplication is commutative, i.e. $ab = ba$

3.3 Rings With Identity

A ring with identity is a ring R that contains one multiplicative identity, i.e. $a1_R = 1_Ra = a$ for all a

3.4 Fields

A field is a commutative ring with identity where all non-zero elements are units

- i.e. $\mathbb{C}, \mathbb{R}, \mathbb{Q}, \mathbb{Z}_p$
- All fields are integral domains

3.5 Integral Domains

An integral domain is a commutative ring with identity where there are no zero divisors

- Every finite integral domain is a field

3.6 Units in Rings

a is a unit in ring R if the equation $ax = xa = 1_R$ has a solution $x \in R$

3.7 Zero Divisor in Rings

a is a zero divisor in ring R if $a \neq 0$ and the equations $ax = 0_R$ or $xa = 0_R$ has a non-zero solution for some $x \in R$

3.8 Multiplicative Inverse in Rings

a is invertible in ring R if and only if $ax = xa = 1_R$ has solutions $x \in R$

- x is the inverse of a , denoted a^{-1}
- In a non-commutative ring, an inverse x of a that satisfies $ax = xa = 1_R$ is called a two-sided multiplicative inverse

3.9 Subrings

A subring is a nonempty subset S of a ring R that can undergo operations inherited from R

- i.e. $\mathbb{C}, \mathbb{R}, \mathbb{Q}, \mathbb{Z}$
- A subring must satisfy all the axioms of a ring
- Existence of a multiplicative identity is not an inherited property
 - The subring of a ring with identity does not have to contain an identity

3.10 Subring Theorem

Given S is a subset of R , S is a subring of R if and only if it satisfies the following axioms

- Closed under subtraction: if $a \in S$ and $b \in S$, then $a - b \in S$
- Closed under multiplication: if $a \in S$ and $b \in S$, then $ab \in S$

3.11 Subring Set Theory

Given that S and T are subrings of R

- $S \cap T$ is a subring of R
- $S \cup T$ is not a subring of R

3.12 Finite Set $\mathbb{F}_p[\theta]$

$\mathbb{F}_p[\theta]$ are the finite sets containing numbers of the form $a + b\theta$ where $a, b \in \mathbb{F}_p$

- \mathbb{F}_p represents the finite set $\{0, 1, \dots, p-1\}$ where p is prime
- The $\mathbb{F}_p[\theta]$ rings are a family of commutative rings with identity
- $\mathbb{F}_p[\theta]$ rings may or may not be fields

3.13 Complex Set \mathbb{C}

\mathbb{C} is the set of complex integers $a + bi$ where $a, b \in \mathbb{R}$

- The \mathbb{C} ring is a field and an integral domain
- $a + bi$ has the multiplicative inverse $\frac{a}{a^2 + b^2} - \frac{bi}{a^2 + b^2}$ where $a^2 + b^2 \neq 0$
- $x = a + bi$ has a complex conjugate of $\bar{x} = a - bi$

3.14 Gaussian Integers $\mathbb{Z}[i]$

$\mathbb{Z}[i]$ is the set of Gaussian integers $a + bi$ where $a, b \in \mathbb{Z}$

- The $\mathbb{Z}[i]$ ring is an integral domain but not a field
- $\mathbb{Z}[i]$ is a disjoint union of $\{0\} \cup \{\pm 1, \pm i\} \cup \{\text{everything else}\}$
- $\mathbb{Z}[i]$ is a subring of \mathbb{C}

3.15 Matrices $M_2(\mathbb{F})$

$M_2(\mathbb{F})$ is the set of 2×2 matrices $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ where $a, b, c, d \in \mathbb{F}$

- \mathbb{F} can be $\mathbb{C}, \mathbb{R}, \mathbb{Q}, \mathbb{Z}$ or \mathbb{F}_p
- The $M_2(\mathbb{F})$ ring is often non-commutative
- $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ has the two-sided multiplicative inverse $\frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$ where $ad - bc \neq 0$

3.16 Units in $M_2(\mathbb{F})$

A is a unit in ring $M_2(\mathbb{F})$ if its determinant $ad - bc \neq 0$ and its inverse is in $M_2(\mathbb{F})$

- The set of units $GL_2(\mathbb{F}) = \{A \in M_2(\mathbb{F}) \mid \det(A) \neq 0\}$ is the 2×2 general linear group over \mathbb{F}
- Multiplicative operations in $GL_2(\mathbb{F})$ satisfy the following axioms
 - Closed under multiplication: if $A \in GL_2(\mathbb{F})$ and $B \in GL_2(\mathbb{F})$, then $AB \in GL_2(\mathbb{F})$
 - * If you multiply two invertible matrices, then the product is also invertible
 - * AB has the two-sided multiplicative inverse $B^{-1}A^{-1}$
 - Associative: $A(BC) = (AB)C$
 - Multiplicative identity: $AI = IA = A$ for all A
 - * $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$
 - Multiplicative inverse: for each A , there exists an element $B \in GL_2(\mathbb{F})$ such that $AB = BA = I$
- $GL_2(\mathbb{F})$ is a subset of the ring $M_2(\mathbb{F})$ but it is not a subring

3.17 Continuous Real-Valued Functions $C[0, 1]$

$C[0, 1]$ is the set of continuous real-valued functions on $[0, 1]$

- The $C[0, 1]$ ring is a commutative ring with identity
- $C[0, 1]$ satisfies all the ring axioms
- Additive operations also satisfy the following axioms
 - Addition: $(f + g)(x) = f(x) + g(x)$
 - Closed under addition: if $f(x) \in C[0, 1]$ and $g(x) \in C[0, 1]$, then $f(x) + g(x) \in C[0, 1]$
 - Additive identity: $f(x) + o(x) = f(x)$, where $o(x) = 0$
- Multiplicative operations also satisfy the following axioms
 - Multiplication: $(f \cdot g)(x) = f(x) \cdot g(x)$
 - Closed under multiplication: if $f(x) \in C[0, 1]$ and $g(x) \in C[0, 1]$, then $f(x) \cdot g(x) \in C[0, 1]$
 - Multiplicative identity: $f(x) \cdot i(x) = f(x)$, where $i(x) = 1$

3.18 Ring Homomorphisms

Rings R and S are homomorphic if there exists a well-defined function $\lambda : R \rightarrow S$ such that

- $\lambda(a +_R b) = \lambda(a) +_S \lambda(b)$
- $\lambda(a \cdot_R b) = \lambda(a) \cdot_S \lambda(b)$

where $+_R, \cdot_R$ are operations defined in R and $+_S, \cdot_S$ are operations defined in S

- As a consequence of the two conditions:
 - Unit preserving: $\lambda(\text{unit}_R) = \text{unit}_S$
 - Multiplicative identity preserving: $\lambda(1_R) = 1_S$
 - Additive inverse preserving: $\lambda(-a) = -\lambda(a)$

3.19 Ring Isomorphisms

Rings R and S are isomorphic, denoted $R \cong S$, if there exists a well-defined bijective function $\phi : R \rightarrow S$ such that

- $\phi(a +_R b) = \phi(a) +_S \phi(b)$
- $\phi(a \cdot_R b) = \phi(a) \cdot_S \phi(b)$

where $+_R, \cdot_R$ are operations defined in R and $+_S, \cdot_S$ are operations defined in S

- The addition and multiplication tables of R and S match when translated via ϕ
- Isomorphic rings have the same size/cardinality
- Isomorphic rings have the same number of units

4 Arithmetic in $\mathbb{F}[x]$

4.1 Polynomials

$R[x]$ is the set of polynomials over ring R of the form $f(x) = a_0 + a_1x + \dots + a_nx^n$ where $a_i \in R$ and $n \geq 0$

- a_n is called the leading coefficient
- x is called the indeterminate
- If the largest coefficient $a_n = 1$, then f is called a monic polynomial
- If n is the largest number for which $a_n \neq 0$, then we say f has degree n , that is $\deg(f) = n$
 - The degree of the zero polynomial $f(x) = 0$ is not defined

4.2 Polynomial Rings

Given $p(x) = a_0 + a_1x + \dots + a_nx^n$ and $q(x) = b_0 + b_1x + \dots + b_mx^m$ in ring $R[x]$ where $m \leq n$

- Addition in $R[x]$
 - $p(x) + q(x) = c_0 + c_1x + \dots + c_nx^n$ where $c_i = a_i + b_i$
 - Additive identity is $0(x) = 0$
 - Standard algebraic addition of polynomials
 - Additive inverse is obtained by replacing all coefficients with their additive inverse in R
- Multiplication in $R[x]$
 - $p(x) \cdot q(x) = d_0 + d_1x + \dots + d_{n+m}x^{n+m}$ where $d_i = \sum_{k=0}^i a_k b_{i-k} = a_0 b_i + a_1 b_{i-1} + \dots + a_i b_0$
 - Multiplicative identity is $1(x) = 1$
 - Standard algebraic multiplication of polynomials

$R[x]$ is a commutative ring with multiplicative identity $f(x) = 1$

- R is a subring of $R[x]$
- If R is an integral domain, then $R[x]$ is an integral domain
- If \mathbb{F} is a field, then $\mathbb{F}[x]$ is an integral domain
- If \mathbb{F} is a field, then the units in $\mathbb{F}[x]$ are precisely the non-zero constant functions
- If \mathbb{F} is a field, then $\mathbb{F}[x] = \{0\} \cup \underbrace{\{0 \text{ degree}\}}_{\text{all units in } \mathbb{F}[x]} \cup \{1 \text{ degree}\} \cup \dots$

4.3 Finite Polynomial Rings

Given $\mathbb{F}_p[x]$ where $\mathbb{F}_p = \{0, 1, \dots, p-1\}$ and p is prime

- There are $(p-1)p^n$ possible polynomials of degree n
- $\mathbb{F}_p[x]$ is an infinite ring but has finite number of polynomials of degree n

4.4 Division Theorem in Polynomials

Given non-zero polynomials $f(x)$ and $g(x)$ in $\mathbb{F}[x]$ where \mathbb{F} is a field, there exists unique polynomials $q(x)$ and $r(x)$ such that $f(x) = g(x)q(x) + r(x)$ where either $r(x) = 0$ or $0 \leq \deg(r) < \deg(g)$

- Use long division to calculate $q(x)$ and $r(x)$

4.5 Divisibility in Polynomials

Given $f(x), g(x) \in \mathbb{F}[x]$, $g(x) \mid f(x)$ means there exists $h(x) \in \mathbb{F}[x]$ such that $f(x) = g(x)h(x)$

4.6 Greatest Common Divisors in Polynomials

Given non-zero polynomials $f(x)$ and $g(x)$, $\gcd(f(x), g(x)) = (f(x), g(x))$ is the monic polynomial of the largest degree that divides both $f(x)$ and $g(x)$

- Finding $\gcd(f(x), g(x))$ by the Euclidean algorithm

Lemma 4.6.1. *If $g(x) \mid f(x)$, then $\gcd(f(x), g(x)) = g^*(x)$ where $g^*(x)$ is the monicization of $g(x)$*

Lemma 4.6.2. *If $a(x) = b(x)c(x) + d(x)$, then $\gcd(a(x), b(x)) = \gcd(b(x), d(x))$*

Given non-zero polynomials $f(x)$ and $g(x)$, the recursive Euclidean Algorithm computes $\gcd(f(x), g(x))$

The Euclidean Algorithm for polynomials is similar to that for integers

4.7 Bézout's Theorem for Polynomials

If $f(x)$ and $g(x)$ are non-zero polynomials:

Then there exists polynomials $s(x)$ and $t(x)$ such that $\gcd(f(x), g(x)) = s(x)f(x) + t(x)g(x)$

Bézout's Theorem for polynomials is similar to that for integers

4.8 Polynomial Roots

$\alpha \in \mathbb{F}$ is a root of $f(x) \in \mathbb{F}[x]$ if and only if $f(\alpha) = a_0 + a_1\alpha + \dots + a_n\alpha^n = 0$

- α is a root of $f(x)$ if and only if $(x - \alpha) \mid f(x)$
- If $\deg(f) = n$ then $f(x)$ has at most n distinct roots

4.9 Associates

$f(x)$ is an associate of $g(x)$ in $\mathbb{F}[x]$ if and only if $f(x) = c \cdot g(x)$ for some unit $c \in \mathbb{F}$

- $\{\text{associates of } g(x)\} = \{c \cdot g(x) \mid c \text{ is unit}\}$

4.10 Non-Trivial Factorization

$p(x) \in \mathbb{F}[x]$ can be non-trivially factorized if there exists $f(x), g(x) \in \mathbb{F}[x]$ where

$$0 < \deg(f) < \deg(p)$$

$$0 < \deg(g) < \deg(p)$$

such that $p(x) = f(x)g(x)$

4.11 Reducible Polynomials

A non-zero non-unit polynomial $p(x)$ is reducible in field $\mathbb{F}[x]$ if and only if it can be non-trivially factorized

- $p(x)$ is reducible if and only if it can be non-trivially factored such that $p(x) = f(x)g(x)$, where $f(x)$ and $g(x)$ are polynomials of lesser degrees
- If a polynomial is not reducible, it is irreducible

4.12 Irreducible Polynomials

A non-zero non-unit polynomial $p(x)$ is irreducible in field $\mathbb{F}[x]$ if and only if its only divisors are its associates and the non-zero constant polynomial/units

- $p(x)$ is irreducible if and only if it cannot be non-trivially factored such that $p(x) = f(x)g(x)$, where $f(x)$ and $g(x)$ are polynomials of lesser degrees
- If a polynomial is not irreducible, it is reducible
- All polynomials of degree 1 are irreducible by definition

4.13 Theorems on Irreducible Polynomials

Given that $p(x) \in \mathbb{F}[x]$

- Every non-zero non-unit polynomial in $\mathbb{F}[x]$ is a product of irreducible polynomials
- There are infinitely many irreducible polynomials in $\mathbb{F}[x]$
- The factorization of a polynomial into irreducibles is unique
- If $p(x)$ is irreducible and $p(x) \mid b(x)c(x)$, then $p(x) \mid b(x)$ and $p(x) \mid c(x)$
- If $p(x)$ is irreducible and $p(x) = b(x)c(x)$, then either $b(x)$ or $c(x)$ is a non-zero constant polynomial/unit
- Polynomials of degree 1 are always irreducible
- If $\deg(p) = 2$ or 3 , then p is irreducible if and only if p has no roots in \mathbb{F}
- If p is irreducible and $\deg(p) > 1$ then p has no roots in \mathbb{F}
- If p has no roots in \mathbb{F} , this does not imply p is irreducible

4.14 Rational Root Test

If $\frac{r}{s} \in \mathbb{Q}$ is a root of $f(x) = a_0 + a_1x + \dots + a_nx^n$ where $f(x) \in \mathbb{Z}[x]$, then $r \mid a_0$ and $s \mid a_n$

- Rational root test narrows down the set of possible rational roots
- Check these possible roots manually to determine if they are actual roots

4.15 Gauss' Lemma

$p(x) \in \mathbb{Z}[x]$ is reducible in $\mathbb{Q}[x]$ if and only if it is reducible in $\mathbb{Z}[x]$

- Contrapositive: $p(x) \in \mathbb{Z}[x]$ is irreducible in $\mathbb{Z}[x]$ if and only if it is irreducible in $\mathbb{Q}[x]$

4.16 Polynomial Reduction mod n

$\bar{f}(x) = [f(x)]_n$ is the polynomial obtained by reducing all coefficients of $f(x)$ in $\mathbb{Z}[x]$ by mod n

- If $\bar{f}(x)$ is irreducible in $\mathbb{F}_p[x]$ for any prime p that does not divide the leading coefficient of $f(x)$, then $f(x)$ is irreducible in $\mathbb{Q}[x]$

4.17 Checking Irreducibility

The following algorithm checks if $f(x) \in \mathbb{F}_p[x]$ of degree n is irreducible

1. Plug in $0, 1, \dots, p-1$ into $f(x)$ to see if we have a root
2. Consider all polynomials of degree 2 and eliminate those that are reducible
3. Check if the irreducible polynomials of degree 2 divide $f(x)$ via long division
4. Repeat step 2 to step 3 with polynomials of degree $3, 4, \dots, \frac{n-1}{2}$

4.18 Eisenstein's Criterion

Suppose $f(x) = a_0 + a_1x + \dots + a_nx^n$ where $f(x) \in \mathbb{Z}[x]$, if there exists a prime p such that

- p divides each of a_0, a_1, \dots, a_{n-1}
- p does not divide a_n
- p^2 does not divide a_0

then $f(x)$ is irreducible in $\mathbb{Q}[x]$

4.19 Fundamental Theorem of Algebra

Every non-constant polynomial in $\mathbb{C}[x]$ has a root in \mathbb{C}

- The irreducible polynomials in $\mathbb{C}[x]$ are precisely the degree 1 polynomials

4.20 Polynomials in $\mathbb{R}[x]$

- All degree 1 polynomials in $\mathbb{R}[x]$ are irreducible
- Only degree 2 polynomials which have complex roots in $\mathbb{R}[x]$ are irreducible
 - Polynomial $ax^2 + bx + c$ in $\mathbb{R}[x]$ has negative discriminant $b^2 - 4ac$
- Complex roots in $\mathbb{R}[x]$ occur in conjugate pairs (i.e. $a \pm bi$ are roots)
 - Product of conjugate pairs is a real polynomial of degree 2

4.21 Roots of Unity

A complex number $z \in \mathbb{C}$ is called an n^{th} root of unity if z is a root of the polynomial $f(x) = x^n - 1$ such that $z^n = 1$

- The primitive n^{th} root of unity is given by the complex number $\omega = e^{\frac{2\pi i}{n}} = \cos \frac{2\pi}{n} + i \sin \frac{2\pi}{n}$
 - A root of unity is said to be primitive if it is not the power of another root of unity
- The n^{th} roots of unity are given by $\omega^k = e^{\frac{2k\pi i}{n}} = \cos \frac{2k\pi}{n} + i \sin \frac{2k\pi}{n}$ for $k = 0, 1, \dots, n-1$
- The n^{th} roots of unity represent the vertices of an n sided polygon inscribed in the unit circle
- The n^{th} roots of unity have the identity $1 + \omega + \omega^2 + \dots + \omega^{n-1} = 0$
- The complete factorization of $f(x)$ into irreducibles is given by $\prod_{k=0}^{n-1} (x - \omega^k)$
- Roots of unity also exist in $\mathbb{Q}[x]$, $\mathbb{R}[x]$ and $\mathbb{F}_p[x]$

5 Congruence in $\mathbb{F}[x]$ and Congruence-Class Arithmetic

5.1 Polynomial Congruence

Given two polynomials $f(x)$ and $g(x)$ and a modulus $p(x)$, $f(x)$ is congruent to $g(x)$ if and only if $p(x) \mid f(x) - g(x)$. This is denoted as $f(x) \equiv g(x) \pmod{p(x)}$

- $f(x)$ and $g(x)$ are congruent if they have the same remainder after long division by $p(x)$
- Polynomial congruence is reflexive, symmetric and transitive

5.2 Polynomial Congruence Class of $f(x) \pmod{p(x)}$

The polynomials congruence class of $f(x) \pmod{p(x)}$, denoted $[f(x)]_{p(x)}$, is the set of all polynomials that are congruent to $f(x) \pmod{p(x)}$

- $[f(x)]_{p(x)} = \{g(x) \mid g(x) \equiv f(x) \pmod{p(x)}\}$
- Polynomial congruence classes are either equal or disjoint
- $f(x)$ is typically the least residue $\pmod{p(x)}$

5.3 Polynomial Congruence Ring $\mathbb{F}[x]_{p(x)}$

$\mathbb{F}[x]_{p(x)}$ is the set of disjoint classes $[r(x)]_{p(x)}$ where $r(x)$ is are least residues $\pmod{p(x)}$ such that $r(x) = 0$ or $0 \leq \deg(r) < \deg(p)$

- $\mathbb{F}[x]_{p(x)} = \{r(x) \mid r(x) = 0 \text{ or } 0 \leq \deg(r) < \deg(p)\}$
 $= \{a_0 + a_1x + \dots + a_nx^n \mid a_0, a_1, \dots, a_n \in \mathbb{F} \text{ and } n = \deg(p) - 1\}$
- $[f(x)]_{p(x)} + [g(x)]_{p(x)} = [f(x) + g(x)]_{p(x)}$
- $[f(x)]_{p(x)} \cdot [g(x)]_{p(x)} = [f(x) \cdot g(x)]_{p(x)}$
- Additive identity is $O(x) = [0]_{p(x)}$
- Multiplicative identity is $I(x) = [1]_{p(x)}$

5.4 Fields in $\mathbb{F}[x]_{p(x)}$

$\mathbb{F}[x]_{p(x)}$ is a field if and only if $p(x)$ is irreducible in $\mathbb{F}[x]$ where \mathbb{F} is a field

- For all $f(x) \in \mathbb{F}[x]$, $\gcd(f(x), p(x)) = 1$

5.5 Finite Fields in $\mathbb{F}_p[x]_{p(x)}$

Given an irreducible polynomial in $\mathbb{F}_p[x]_{p(x)}$ of degree n , we can construct a finite field in $\mathbb{F}_p[x]_{p(x)}$ of order p^n

6 Ideals and Quotient Rings

6.1 Ideals

A subring I of ring R is an ideal in R if $ra \in I$ and $ar \in I$ for all $r \in R$ and $a \in I$

- If-condition can be simplified as $RI \subseteq I$ and $IR \subseteq I$
- A proper ideal I in R satisfies $I \subset R$
- All ideals are subrings
- Not all subrings are ideals

A subset I of a ring R is an ideal in R if and only if has the following properties

1. I is non-empty
2. If $a, b \in I$, then $a - b \in I$
3. If $r \in R$ and $a \in I$, then $ra \in I$ and $ar \in I$

6.2 Maximal Ideals

Let R be a commutative ring with identity. Then ideal M in R is maximal if $M \subset R$ and the only ideals containing M are M and R

- There does not exist an ideal J such that $M \subset J \subset R$
- i.e. M is as large as possible while being a proper subset of R

6.3 Finitely Generated Ideals

Let R be a commutative ring with identity and $c_1, c_2, \dots, c_n \in R$. Then $I = \{r_1c_1 + r_2c_2 + \dots + r_nc_n \mid r_1, r_2, \dots, r_n \in R\}$ is a finitely generated ideal in R

6.4 Principal Ideals Generated by c

Let R be a commutative ring with identity and $c \in R$. Then $I = \{rc \mid r \in R\}$ is the principal ideal generated by c , denoted (c)

- If $(m) \subseteq (n)$, then $n \mid m$
- Principal ideals are a special case of finitely generated ideals where $n = 1$

6.5 Principal Ideal Domains

A principal ideal domain (PID) is an integral domain in which every ideal is principal

- An integral domain is a commutative ring with identity with no zero divisors
- If \mathbb{F} is a field, then \mathbb{F} is a principal ideal domain
- i.e. $\mathbb{Z}, \mathbb{F}[x], \mathbb{Z}[i]$

6.6 Ideals in \mathbb{Z}

Every subring in \mathbb{Z} is an ideal in \mathbb{Z} , and every ideal in \mathbb{Z} is a principal ideal generated by some c

- For every subring I in \mathbb{Z} , there exists $c \in \mathbb{Z}$ such that $I = (c)$
- c is the smallest positive element in I
- The maximal ideals in \mathbb{Z} are (p) for prime integers p
- The maximal ideal in $\mathbb{Z}/(p)$ is (p) when p is prime
- $\mathbb{Z}/(p)$ is a field if and only if p is prime


6.7 Ideals in $\mathbb{Z}[i]$

Every ideal in $\mathbb{Z}[i]$ is a principal ideal generated by some $a + bi$

- There are $N(a + bi)$ distinct ideals in $\mathbb{Z}[i]/(a + bi)$
- The maximal ideals in $\mathbb{Z}[i]$ are $(a + bi)$ for Gaussian primes $a + bi$
- The maximal ideal in $\mathbb{Z}[i]$ is $(a + bi)$ when $a + bi$ is a Gaussian prime
- $\mathbb{Z}[i]/(a + bi)$ is a field if and only if $a + bi$ is a Gaussian prime

6.8 Ideal Lattice

Ideal lattices describe the containment relations between different ideals



The ideal lattice tells us that

- (1) contains (2) and (3)
- (2) and (3) contain (6)
- (6) contains (12)

6.9 Ideal Congruence

Given ideal I in ring R and two elements a and b in R , a is congruent to b modulo I if and only if $a - b \in I$ or $a + I = b + I$. This is denoted as $a \equiv b \pmod{I}$

- Ideal congruence is reflexive, symmetric and transitive

6.10 Ideal Congruence Class of $a \bmod I$

Given ideal I in ring R and element a in R , the ideal congruence class of $a \bmod I$, denoted $[a]_I$, is the set of all elements in R that are congruent to $a \bmod I$

- $[a]_I = \{b \mid b \equiv a \bmod I\} = \underbrace{a + I}_{\text{left coset of } I \text{ represented by } a}$
- Ideal congruence classes / left cosets are either equal or disjoint

6.11 Lagrange's Theorem for Finite Rings

If R is a finite ring and I is an ideal in R , then $|I| \mid |R|$

- The cardinality of the ideal is a divisor of the cardinality of the ring

6.12 Quotient Ring R/I

R/I , also denoted as R_I , is the set of disjoint left cosets $[a]_I = a + I$ where a are elements in R

- $R/I = \{a + I \mid a \in R\}$
- $(a + I) + (b + I) = (a + b) + I$
- $(a + I) \cdot (b + I) = (a \cdot b) + I$
- Additive identity is $0 + I = I = [0]$
- Multiplicative identity is $1 + I = [1]$

6.13 Kernels

The kernel of a ring homomorphism $f : R \rightarrow S$ is $\text{Ker}(f) = \{r \in R \mid f(r) = 0_S\}$

- $\text{Ker}(f)$ contains every element in the domain R that has 0 value in the co-domain S
- $\text{Ker}(f)$ is an ideal in R
 - Given $a, b \in \text{Ker}(f)$, $a - b \in \text{Ker}(f)$ since $f(a - b) = f(a) - f(b) = 0_S - 0_S = 0_S$
 - Given $r \in R$ and $a \in \text{Ker}(f)$, $ra \in \text{Ker}(f)$ since $f(ra) = f(r) \cdot f(a) = f(r) \cdot 0_S = 0_S$
- $\text{Ker}(f) = \{0_R\}$ if and only if
 - f is injective
 - R is isomorphic to $f(R)$

6.14 Natural Homomorphisms

A natural homomorphism from R to R/I is a map $\pi : R \rightarrow R/I$ given by $\pi(r) = r + I$

- π is a surjective homomorphism with kernel I
- Natural homomorphisms are a special case of surjective homomorphisms

6.15 First Isomorphism Theorem

Let $f : R \rightarrow S$ be a surjective homomorphism of rings with $K = \text{Ker}(f)$. Then there exists an isomorphic function \bar{f} between R/K and S

$$\begin{array}{ccc} R & \xrightarrow{f} & S \\ \pi \downarrow & \nearrow \bar{f} & \\ R/K & & \end{array}$$

- $f(r) = \bar{f}(\pi(r)) = \bar{f}(r + K)$

6.16 Product Decomposition Theorem

Let a, b be positive integers and $\gcd(a, b) = 1$. Then $\mathbb{F}/(ab)$ is isomorphic to $\mathbb{F}/(a) \times \mathbb{F}/(b)$

6.17 Product Decomposition Theorem for Polynomials

Let $f(x), g(x)$ be polynomials in $\mathbb{F}[x]$ and $\gcd(f(x), g(x)) = 1$. Then $\mathbb{F}[x]/(f(x)g(x))$ is isomorphic to $\mathbb{F}[x]/(f(x)) \times \mathbb{F}[x]/(g(x))$

6.18 Additional Theorems

- M is a maximal ideal if and only if \mathbb{F}/M is a field
- $\mathbb{F}[x]/(f(x))$ is a field if and only if $f(x)$ is irreducible in $\mathbb{F}[x]$
- $(n) \cap (m) = (\text{lcm}(n, m))$
- $(n) + (m) = (\gcd(n, m))$
- $(n)(m) = (nm)$

6.19 Prime Ideals

An ideal P in ring R is called prime if $bc \in P$ implies $b \in P$ or $c \in P$

- P is a prime ideal in ring R if and only if R/P is an integral domain
- Prime ideals in \mathbb{Z} are (p) where p is prime

6.20 Maximal and Prime Ideals

- If M is a maximal ideal, then M is a prime ideal
- Let \mathbb{F} be a field and I a non-zero ideal in $\mathbb{F}[x]$. The following are equivalent
 - I is a maximal ideal
 - I is a prime ideal
 - $I = (f(x))$ for some irreducible polynomial $f(x) \in \mathbb{F}[x]$

Similar theorem holds true for integers

7 Arithmetic in $\mathbb{Z}[i]$

7.1 Division Theorem in Gaussian Integers

Given non-zero Gaussian integers $a_1 + a_2i$ and $b_1 + b_2i$, there exists $q_1 + q_2i$ and $r_1 + r_2i$ such that $a_1 + a_2i = (b_1 + b_2i)(q_1 + q_2i) + (r_1 + r_2i)$ where $N(r_1 + r_2i) \leq N(b_1 + b_2i)$

- The quotient and remainder is not unique

7.2 Divisibility in Gaussian Integers

Given $a_1 + a_2i, b_1 + b_2i \in \mathbb{Z}[i]$, $b_1 + b_2i \mid a_1 + a_2i$ means there exists $c_1 + c_2i \in \mathbb{Z}[i]$ such that $a_1 + a_2i = (b_1 + b_2i)(c_1 + c_2i)$

7.3 Greatest Common Divisors in Gaussian Integers

Given non-zero Gaussian integers $a_1 + a_2i$ and $b_1 + b_2i$, $\gcd(a_1 + a_2i, b_1 + b_2i) = (a_1 + a_2i, b_1 + b_2i)$ is a common divisor of $a_1 + a_2i$ and $b_1 + b_2i$ with largest norm

- $a_1 + a_2i$ and $b_1 + b_2i$ are relatively prime if $(a_1 + a_2i, b_1 + b_2i)$ is a unit in $\mathbb{Z}[i]$
 - i.e. $(a_1 + a_2i, b_1 + b_2i) = \pm 1$ or $\pm i$
- The greatest common divisor is not unique
- $\gcd(a_1 + a_2i, b_1 + b_2i)$ can be found by the Euclidean algorithm

7.4 Bézout's Theorem for Gaussian Integers

If $a_1 + a_2i$ and $b_1 + b_2i$ are non-zero Gaussian integers

Then there exists Gaussian integers $s_1 + s_2i$ and $t_1 + t_2i$ such that

$$\gcd(a_1 + a_2i, b_1 + b_2i) = (s_1 + s_2i)(a_1 + a_2i) + (t_1 + t_2i)(b_1 + b_2i)$$

Bézout's Theorem for Gaussian integers is similar to that for integers

7.5 Non-Trivial Factorization

$a_1 + a_2i \in \mathbb{Z}[i]$ can be non-trivially factorized if there exists $b_1 + b_2i \in \mathbb{Z}[i]$ where

$$0 < N(b_1 + b_2i) < N(a_1 + a_2i)$$

such that $(b_1 + b_2i) \mid a_1 + a_2i$

7.6 Unique Factorization Theorem

If there are two factorizations of a Gaussian integer, then each component of the factorizations will equal each other, or differ by a factor of $-1, i$ or $-i$

7.7 Gaussian Primes

If a Gaussian integer $a_1 + a_2i$ with $N(a_1 + a_2i) > 1$ has only trivial factors, then it is a Gaussian prime

- The trivial factors are $\pm 1, \pm i, \pm(a_1 + a_2i), \pm(a_1 + a_2i)i$

If $a_1 + a_2i$ is a Gaussian integer and $N(a_1 + a_2i)$ is a prime integer, then $a_1 + a_2i$ is a Gaussian prime

- Note that the reverse implication does not hold

If $\pi \in \mathbb{Z}[i]$ is a Gaussian prime, then there exists a prime integer p such that $\pi \mid p$ in $\mathbb{Z}[i]$

- Gaussian primes are the factors of the prime integers in $\mathbb{Z}[i]$

Every Gaussian prime π has one of three forms:

- $\pi = \pm p$ or $\pm ip$ for some prime integer p where $p \equiv 3 \pmod{4}$
- π is part of an octet of factors $\pm a + \pm b$ or $\pm b + \pm a$ such that $p = a^2 + b^2$ and $p \equiv 1 \pmod{4}$
- π is one of $\pm 1, \pm i$

8 Appendix

8.1 Rings

Ring	Properties			
\mathbb{R}	infinite	commutative	has identity	field
\mathbb{Q}	infinite	commutative	has identity	field
\mathbb{E}	infinite	commutative	no identity	not field
\mathbb{Z}	infinite	commutative	has identity	not field
\mathbb{Z}_n	finite	commutative	has identity	not field
\mathbb{Z}_p	finite	commutative	has identity	field
\mathbb{C}	infinite	commutative	has identity	field
$\mathbb{Q}(\sqrt{2})$	infinite	commutative	has identity	field
$\mathbb{Z}_3[i]$	finite	commutative	has identity	field
$M_2(\mathbb{Z})$	infinite	not commutative	has identity	not field
$M_2(\mathbb{E})$	infinite	not commutative	no identity	not field
$M_2(\mathbb{Z}_n)$	finite	not commutative	has identity	not field
$M_2(\mathbb{Z}_p)$	finite	not commutative	has identity	not field
$\left\{ \begin{pmatrix} 0 & r \\ 0 & 0 \end{pmatrix} \mid r \in \mathbb{Z}_n \right\}$	finite	commutative	no identity	not field

8.2 Norm Functions

- $N(a) = a^2$
- $N(a + b\sqrt{-1}) = a^2 + b^2$
- $N(a + b\sqrt{-m}) = a^2 + b^2m$
- $N(a + b\sqrt{m}) = a^2 - b^2m \leftarrow \text{verification needed}$

\mathbb{E} denotes the ring containing all integers divisible by 2, i.e. $-2, 0, 4$

8.3 Quick Proofs

- Every field \mathbb{F} is an integral domain

Given $a, b \in \mathbb{F}$,

$$ab = 0 \rightarrow a^{-1}ab = a^{-1}0 \rightarrow b = 0$$

$$ab = 0 \rightarrow abb^{-1} = 0b^{-1} \rightarrow a = 0$$

Therefore, there are no zero divisors in \mathbb{F}

- Prove that S is a subring of R

Given $a, b \in S$

$$a \in R \text{ and } b \in R \text{ such that } S \subset R$$

$$a - b \in S \text{ such that } S \text{ is closed under subtraction}$$

$$ab \in S \text{ such that } S \text{ is closed under multiplication}$$

Therefore, S is a subring of R

- a is a unit / invertible in \mathbb{Z}_n if $\gcd(a, n) = 1$

Given $\gcd(a, n) = 1$

By Bézout's Theorem there exists $x, y \in \mathbb{Z}_n$ such that $ax + ny = 1$

Then $ny = 1 - ax$ and $n \mid 1 - ax$ such that $ax \equiv 1 \pmod{n}$

Therefore, there exists $x \in \mathbb{Z}_n$ such that $ax = 1$ and a is a unit / invertible

- Check if $\mathbb{F}[x]$ contains units

Given $A \in \mathbb{F}[x]$

$$N(AB) = N(A)N(B)$$

$$N(1) = 1$$

If A is a unit, then there exists $B \in \mathbb{F}[x]$ such that $AB = 1$

This is equivalent to $N(A)N(B) = N(1) = 1$

Therefore if A is a unit, then there exists $B \in \mathbb{F}[x]$ such that $N(A)N(B) = 1$