## 

Personal communication Ian J. Wright Personal communication John Morgan

personal communication Sean Gleason

Personal communication, C. James

## References

- Clifford, H. T. (2000). Dicotyledon seedling morphology as a correlate of seed-size h. trevor clifford. *Proceedings of the Royal Society of Queensland*, 109:39–48.
- Gallagher, R. V., Hughes, L., and Leishman, M. R. (2013). Species loss and gain in communities under future climate change: consequences for functional diversity. *Ecography*, 36(5):531–540.
- Gallagher, R. V. and Leishman, M. R. (2012). Contrasting patterns of trait-based community assembly in lianas and trees from temperate Australia. *Oikos*, (March):1–10.
- Kyle, G. and Leishman, M. R. (2009). Plant functional trait variation in relation to riparian geomorphology: the importance of disturbance. Austral Ecology, 34(7):793– 804.
- Lake, J. C. and Leishman, M. R. (2004). Invasion success of exotic plants in natural ecosystems: the role of disturbance, plant attributes and freedom from herbivores. *Biological Conservation*, 117(2):215–226.
- Stuart, S. A. (2011). Cold Comfort: Diversification and Adaptive Evolution across Latitudinal Gradients.
- Tng, D. Y. P., Jordan, G. J., and Bowman, D. M. J. S. (2013). Plant traits demonstrate

4 REFERENCES

that temperate and tropical giant eucalypt forests are ecologically convergent with rainforest not savanna.  $PLoS\ ONE,\ 8(12):1-13.$ 

Westoby, M. (1998). A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and Soil*, 199(2):213–227.