Four points and meeting regions

Mathematical overview | Draft and unpiloted | 3 October 2026

The mathematical core

Planar Radon theorem: Any four labeled points in the plane can be partitioned into two nonempty groups whose convex hulls intersect. Each label is used once. Coincident labels and collinear placements are allowed; intersection at a single point is enough.

The convex hull is the filled stretched-band region: a singleton is a point, two locations give a closed segment, and three noncollinear locations give a filled triangle. The guarantee is sharp: three noncollinear points have no successful split. There are seven unordered splits of four labels: four of type 1+3 and three of type 2+2.

If no three of four distinct points are collinear, exactly one split succeeds: either the crossing diagonals of a convex quadrilateral, or an interior point versus the triangle of the other three. Degenerate placements can have several successful splits. With four labels, the intersection can be a point or segment, but cannot contain a filled triangle.

What the activity develops

K–1: Make legal groups and point to an actual shared location. The existing adult launch models a point, a segment and a shaded triangle before the tasks.

Grades 2–3: Organize a complete list, distinguish duplicate splits, and construct arrangements with different successful choices.

Grades 4–5: Separate existence from uniqueness and explain why the guarantee survives collinear and coincident cases; contrast four-point existence with the three-point counterexample.

From trials to explanation

Trying many placements motivates the four-point conjecture. Checking all seven splits proves a claim about one placement only. A geometric case argument establishes the guarantee for every placement. The detailed tools and complete proof remain on printed pages 4 and 18.

Meeting regions facilitator guide

Week 22 | Planar convex partitions | Prepared 3 October 2026, revised 6 November 2026

Help children make two filled stretched-band regions from four labeled points, then ask whether every placement can be split successfully. The answer is yes. Three points do not give the same guarantee.

Status and scope. This is a separate adult-guide draft (F22-FAC-v7) for the current student v7 packets, renumbered after the 6 November 2026 kid-fit revision. It is unpiloted and belongs to an unscheduled activity-library slot. The Grades 2-3 and 4-5 opening pages include the E-J region key and a complete K-L-M-N split; the K-1 opening shows only the K-L-M-N split. Timings below are planning choices, not classroom evidence.

Use this guide at the table

Begin with materials and a short shared demonstration (guide pp. 2-3). Offer a few pages at a time. Keep the answer galleries, seven-split checklist, and proof cases with the adult until children need them. A child may enter another band by readiness.

Notation: AB | CD means one group contains A and B, and the other contains C and D. Each label occurs once. Reversing the two groups gives the same split. In the diagrams, the first group is solid blue and the second dashed rust; pale fill shows a triangle. A colored ring shows a singleton; an additional black ring marks a shared point. Gold marks a shared segment. Color is optional: solid/dashed and labels carry the meaning.

Student packet and guide finder

K-1: 6 student pages, F22-K-v7.
P1 on student p. 1 -> guide p. 5; P2 on p. 2 and P3 on p. 3 -> p. 6; P4 on p. 4 -> p. 7; P5 on p. 5 -> p. 8; P6 on p. 6 -> p. 9.

Grades 2-3: 10 student pages, F22-23-v7.
P1 on student pp. 1-2 -> guide p. 10; P2 on p. 3, P6 on p. 9 and P7 on p. 10 -> p. 11; P3 on pp. 4-5 -> p. 12; P4 on p. 6 and P5 on pp. 7-8 -> p. 13.

Grades 4-5: 11 student pages, F22-45-v7.
P1 on student pp. 1-2 -> guide p. 14; P2 on pp. 3-4 and P3 on p. 5 -> p. 15; P4 on p. 6 -> p. 16; P5 on pp. 7-9 -> p. 17; P6 on p. 10 -> p. 18; P7 and P8 on p. 11 -> p. 19.

Common mathematical tools are on p. 4. Extensions, fidelity limits, and pilot notes are on p. 20. Source verification and reproducibility notes are on p. 21.

Preparation and a concrete launch

Allow about 15 minutes of adult preparation before the session

Prepare three tables

At each table: two distinct A-D marker sets (eight movable markers), four Letter tracing sheets, two colors of pencil or crayon for each pair, a ruler, an eraser, ten blank sheets, and removable tape. Give the two groups different colors or solid/dashed hatching. Keep spare tracing paper nearby for the multi-answer problems and a fresh sheet for each game round.

Print single-sided on US Letter at 100 percent. Each full work board is 12.6 cm square. The 5.2 to 7.4 cm record boards preserve answers. K-1 P1 and P4 print D on each board, so children draw straight onto them. In Grades 2-3 and 4-5, P1's record page has one small board per cross, with D printed; the large board stays for moving the marker.

For the current eleven-child group (four K-1, four middle, three upper), one packet per child means 4 K-1, 4 middle, and 3 upper packets (97 student sheets total). A lower-paper option is one packet per pair plus blank/tracing paper: 2 of each band, 54 sheets. Adults should decide before printing; no child needs to finish every page.

Read the answer key for your starting tasks, try a filled triangle containing a singleton, and check that both labels remain visible when stacked. Adults can draw a tiny cross for a center. Coordinate arithmetic in this guide is only an exact adult audit; children need none.

A three-minute whole-group launch

First let children handle the markers. Then use the K, L, M, N points printed on the opening key. Say: "Split the four labels into two groups, using every label once. Draw the smallest stretched-band region for each group. Find a point that belongs to both." Once, stretch a real rubber band around three upright markers and lift it away: the region is everything inside the band.

Use the printed K-L-M-N rectangle: put K and M in one group, L and N in the other. Draw the two segments without moving the points or changing either group. Ring their crossing: this point belongs to both regions, although it is not one of the four labels. Let a child point to each group and the shared point before starting an independent split.

Use the printed E-J key: E gives its point; F and G give the closed segment, including its endpoints; H, I and J give the lightly filled triangle, including its edge and interior. The small unlabelled interior dot marks a location, not another group label. Keep this key visible while children build their own regions. The K-1 opening prints only the K-L-M-N split: at a K-1 table, draw a point, a segment and a shaded triangle with a dot inside as you say this.

Say explicitly. "Touching counts. Use the centers of the marks. Swapping the two group colors does not make a new split. On a line, draw one group just above the dots and the other just below." Two labels may intentionally share a location, but both labels must still be used. Give each pair two colors, one per group, so overlapping segments on K-1 P6, middle P3 and upper P2 stay visible.

Physical safety and accuracy. No pins or stretched rubber bands are needed. If used, adults manage them. A rubber loop around one peg is not a small disk, and a loop around two pegs is not a thin rectangle. These are ideal points and segments.

Flexible hour menu and prerequisite gates

Choose by observed readiness | No reading or arithmetic race

A possible hour

0-5 minutes: handle markers; make and change groups.
5-8: shared action launch.
8-25: work on a first fixed configuration and preserve a solution.
25-28: optional movement reset; children stand at four imaginary points and name two groups.
28-50: continue the chosen route, a construction, or a placement game.
50-57: share one surprising success or failed attempt.
57-60: save a picture and clear the table.

The group-work budget is about forty minutes. A productive problem may occupy most of it. Offer a route change if energy drops; do not impose the next representation merely because time has passed.

Three starting routes

Concrete meeting route: K-1 P1, then the games P2 and P3. P4 is optional: each place of D lies exactly on a line. P5 and P6 are return pages, and P6 (dots in a row) is optional for K-1. Adult reads aloud and records dictated groups. Stop after a child can build a legal split and identify an actual shared point.

Complete-search route: middle P1, the game P2, then P3, P4 or P5. The three-dot pages P6 and P7 can follow the game directly or continue another day. Stop after a complete list with a sensible reason, or a pair of constructions with no common winner.

Guarantee route: upper P1, P3, then P4 and P5 as needed before P6-P8. P2 is useful preparation for degenerate cases. Stop after a justified conjecture if a full case argument is premature; distinguish it from a proof.

Gates that matter

Entry: recognize four labels, place each in exactly one nonempty group, and identify a point on a segment or in a filled region. Reading can be supplied orally; counting only to four is needed.
Enumeration: keep two attempts separate, recognize a color-swapped duplicate, and compare a short list.
Universal explanation: understand "every arrangement," a counterexample, an exhaustive case split, and why a boundary case still counts. No coordinates, fractions, algebra, or formal convexity vocabulary are prerequisites.

Help only as needed. Wait for attempts first. Ask a diagnostic question, offer the first held hint, then pause again. The second hint can reveal a useful smaller search. Give a worked split only when the action itself is still unclear.

Common trouble. An outline-only triangle misses interior points; shade it. A thick stroke makes a near miss look successful; use centers and a ruler. An unused D makes an illegal split; physically sort all four markers. Six record spaces do not mean six answers.

Adult mathematical tools

Keep the catalog and proof categories off the student boards

What a region means

A group's convex hull is the smallest filled convex set containing its point locations. A point and a segment count as regions. With three collinear labels, the hull is the segment from the two outside locations, including the third. Repeated locations do not thicken the hull.

Why there are seven candidate splits

For four labeled points, group sizes are 1 + 3 or 2 + 2. There are four choices of singleton, and three ways to pair all labels. Thus there are seven unordered candidates. This is a completeness tool for the adult, not a required student procedure.

Singleton candidates: A | BCD, B | ACD, C | ABD, D | ABC.
Pair candidates: AB | CD, AC | BD, AD | BC.
Our answer tables write A in the first group, so B | ACD appears as ACD | B, for example.

Three quick tests

Point versus triangle: the singleton must lie in or on the other hull. Being near an edge is insufficient.
Segment versus segment: the closed segments must cross, touch, or overlap. Crossing the infinite supporting lines beyond their endpoints is insufficient.
All points on a line: replace each group by the interval from its leftmost to rightmost point. Two intervals meet exactly when they share a point.

Why a general-position example has one answer

For a convex quadrilateral, a singleton lies outside the triangle of the other three, and the two pairings of opposite boundary edges are disjoint. Only the pairing of opposite corners crosses. For a triangle with a fourth point strictly inside, only the inside singleton works: every outside vertex is outside the other triangle, and a segment from an outside vertex to the interior point cannot reach the opposite side.

That last segment stays in the triangle's interior after leaving its starting vertex. The opposite side is part of the boundary, so the segment cannot meet it. These observations check all four singleton and three pairing possibilities, without relying on a picture alone.

Existence and uniqueness are different

Four labels always permit at least one successful split, but there need not be just one. The full proof, including collinearity and coincidence, is on guide p. 18. A single successful example does not establish a guarantee. A single failed split does not establish an impossible arrangement.


The exact rational audit uses separate point-in-triangle and segment-intersection tests for all seven candidates. It does not import the student checker. Its finite stress test supplements the geometric proof; it cannot replace the proof.

K-1 Problem 1


Student p. 1 | Three different positions of D

Gate: use all four labels once and find a point shared by two filled regions. Read the task aloud. D is printed at a different place on each of the three boards, so children draw straight onto them.


Board 1: AD | BC



Board 2, D inside: ABC | D



Board 3: AC | BD



Exact answer: each of the three placements has exactly the one split shown. The outside placements form convex quadrilaterals; their opposite-corner pairs cross. At the middle placement, D is strictly inside triangle ABC, so D | ABC works even though D is on none of the triangle's sides.

Shared points. On the outside placements, mark the actual crossing of the two drawn segments. At the inside placement, mark D itself. The circled point belongs to both regions in each diagram.

Held hint 1. "Can a group have just one dot? What is that dot's region?" Use only if the child rules out 1 + 3 grouping.

Held hint 2. For an outside placement: "Could both regions be straight segments?" Let the child choose which labels to pair.

Reasoning and check. The general-position argument on guide p. 4 rules out all other candidates. The three successful sets are {AD | BC}, {ABC | D}, and {AC | BD}; color-swapped drawings are duplicates.

Show the one-dot move on board 2. Here every pairing of two letters against two fails, and the K-1 key shows only a two-against-two crossing. Once the child has tried the pairings, show the move: D alone in one group, A, B and C in the other. Shade the triangle and point to D inside it; D is the shared point. A child may say "they do not touch" when D is inside the outline: ask whether the whole filled triangle includes D.

Extension and stopping point. After two successful placements, ask a partner to choose a new D location. This leads directly to the game in P2. Stop when the child can demonstrate a shared point without adult rescue; uniqueness is adult background here.

K-1 Problems 2 and 3

Student pp. 2 and 3 | Placement games

Problem 2 with A, B, C fixed

Student page 2. Gate: find and record one successful split for a chosen D placement.

Answer: yes, the responding player can always succeed. The fixed ABC is a noncollinear triangle. If D is inside or on it, ABC | D works. Outside, the appropriate opposite-corner pairing may work; in an outside triangular-hull case an original vertex can be the inside point instead. A point on an extended side can also make an original vertex the middle of a collinear triple.


Do not tell children that "outside means pair diagonals" in every case. For example, if D is far enough beyond A, A may lie inside BCD, so A | BCD is the useful split. Use the exhaustive rule on guide p. 18 if the adult is unsure.

Problem 3 with every point movable


Student page 3. Gate: the same legal-split action, plus patience with a partner's placement. Taking turns means one child places all four markers and the partner splits them; then they swap.

Answer: yes again. No arrangement of four labeled planar points defeats every split. This remains true on one line and at repeated locations. Children can test a conjecture through play; they need not produce the full proof to participate.


Inside singleton



Crossing pairs



Boundary contact



Held hint 1. "Could one point already belong to the region made by the other three?" Ask only after a real attempt, and allow the child to choose the singleton.

Held hint 2. "Try making two straight regions." If labels coincide, ask what happens when those two labels are put in different groups.

Adult proof and game care. Use guide p. 18 for all arrangements. Take turns placing, grouping, and checking. A slow response is not evidence of an impossible placement. Allow deliberate coincidences with both labels visible; do not enforce an unannounced general-position rule. A D placed inside the triangle needs the one-dot move from P1's second board.


When to stop. Save one attempt that first looked impossible. Let its maker explain the eventual shared point. Return to a fixed board if placement freedom overwhelms the child.

K-1 Problem 4


Student p. 4 | Two meeting groupings at three boundary placements | Optional

Gate: tell a new grouping from a color swap. Each place of D has two boards and exactly two successful splits, so two correct boards per place are a complete answer.


D on AB: AB | CD




D on AB: ABC | D




D beyond B: AD | BC




D beyond B: ACD | B



D on AC: AC | BD



D on AC: ABC | D



Complete lists: between A and B: AB | CD and ABC | D (shared point D). Beyond B: AD | BC and ACD | B (shared point B). On AC: AC | BD and ABC | D (shared point D). Each has exactly two.

Why no others. The middle point of the collinear triple can stand alone opposite a triangle; or it can be paired with the off-line point opposite the two endpoints. Every other split separates the hulls. Check the seven adult candidates if a child asks whether the list is complete.

Held hints. First: "Which point is between two others?" Then: "Can that point be alone? Can it be in a two-dot group instead?" Keep each success on its own board.


Optional for K-1. Every place of D here lies exactly on a line: on side AB, on line AB beyond B, and on side AC. One of the two ways on each board works only because D is exactly on that line, and a hand-drawn side AC can pass a millimetre beside D. Offer the page to children who are ready, perhaps only the two horizontal places (the first four boards), and skip it otherwise; the games and P5 do not depend on it.

K-1 Problem 5


Student p. 5 | Which crosses for C work

Gate: with three labels, one group is a single dot and the other a segment. The task asks which printed crosses work; every position is an extension.


Middle cross: AB | C



Right cross: AC | B



Upper cross: impossible



Answer: circle the middle cross (AB | C, shared point C) and the right cross (AC | B, shared point B). The upper cross fails: no split of A, B and C meets there.
Every position, for an extension: C may be anywhere on the entire horizontal line through A and B, restricted to the square, including the portions left of A and right of B and coincidence with A or B. No point off that line works.

Exact locus: A = (2.5, 5.6), B = (8.8, 5.6). Thus C = (x, 5.6) with 0 <= x <= 12.6. Coordinates are adult checks measured from the board's lower-left corner; children can draw the full line with a ruler.

Why these are all the positions

For three distinct collinear locations, one is in the middle. Put that label alone, and the other two make a segment containing it. If C is left of A, use A | BC; if C is between A and B, use AB | C; if C is right of B, use AC | B.

If C = A, both A | BC and AB | C work. If C = B, both AC | B and AB | C work. Coincident labels stay distinct labels. If C is off line AB, the three points form a triangle. Each of the three splits isolates a vertex that is not on the opposite side, so every split fails.

Held hint 1. "Does C have to be the dot that is alone?" This challenges the common guess that only segment AB works.

Held hint 2. "Try C beyond A. Which of the three dots is in the middle now?" Then let the child test an off-line point and use the three singleton choices.

Extension and evidence. Ask a partner to put C just above the line, then exactly on it. Judge centers, not the thickness of the marker. Children who are ready can trace every place C could go: the whole horizontal line, not only the circled crosses.

K-1 Problem 6

Student p. 6 | Dots in a row, printed order A, D, C, B | Optional return page

Gate: treat each group on a line as a segment or point. The labels are deliberately out of alphabetical order. The task asks for two ways, on four boards.

AB | CD



AC | BD



ABC | D



ABD | C



Exactly four successes: AB | CD, AC | BD, ABC | D, and ABD | C. The first two share the full segment from D to C. The third shares only D, and the fourth shares only C. Any two of them answer the task.

On this board D = (4.5, 6.3) and C = (7.5, 6.3), so the common segment of the two pairing solutions has length 3 cm at actual size. Measuring it is not required.

A complete child-accessible explanation

A or B alone cannot reach the other group because each is an outside point. C or D alone lies in the segment spanned by the other three, so both singleton choices work. Of the three pairings, AB | CD and AC | BD have overlapping intervals. AD | BC leaves a gap between D and C, so it fails. That checks all seven possibilities.

Held hint 1. "If three dots lie on one straight line, what is their stretched-band region?" Let the child cover it with a ruler or a strip of tracing paper.

Held hint 2. "Could a middle dot be a group by itself? What happens if an end dot is by itself?" If needed, keep the two successful pairings alongside the two singleton answers.

Show what meeting means on a line. A group drawn just above the dots and a group drawn just below never touch on paper; they meet because their stretches overlap or one contains the other's dot. Run a finger along each stretch and show the shared part on the line itself. Mark the page optional for K-1 and skip it if this convention needs repeated explaining.


Extension. Swap two labels while leaving the locations fixed. The number stays four, but the written winning splits may change. A child can predict the new label strings and then check them.

Grades 2-3 Problem 1


Student pp. 1-2 | Four placements, including exact boundary contact

Gate: shade a filled triangle, draw closed segments, and mark a genuine common point. One success per cross is required; the boundary cross has two possible answers. Move the D marker on the large board; draw each placement on its own record board on p. 2.


Upper-right: AD | BC




Interior: ABC | D



On AB: AB | CD



Below AB: AB | CD



All successful sets by cross: upper-right {AD | BC}; interior {ABC | D}; on AB {AB | CD, ABC | D}; below AB {AB | CD}. At the on-AB cross, either answer meets at D. At the interior cross, the shared point is D. The remaining crosses meet at the drawn segment crossing.

Proof check. The off-boundary examples have no three collinear and use guide p. 4. The on-AB cross is D = (6, 2.5), exactly halfway along A = (2, 3) to B = (10, 2). It is a real contact case. D = (6, 1) is below AB, so the last crossing is not at D.

Held hints. First: "Can a point be a region?" Then, if needed: "What happens when you pair the dots across the outside shape?" Do not require an exhaustive list on this launch page.

Grades 2-3 Problems 2, 6 and 7

Student pp. 3, 9 and 10 | The four-point game and the three-point contrast

The three-dot pages can follow the game directly: seeing three dots fail is what makes "four always works" a surprise.

Problem 2 four-point challenge

Student page 3. Gate: build a legal split for a partner's example and distinguish a failed attempt from impossibility.

Answer: four always suffice. The responder has a successful split for every placement, including collinear and coincident ones. Use the complete proof on guide p. 18 as the adult key. Any four-point drawing claimed impossible can be checked against the seven candidate splits.


Held hints for P2. First: "Can one dot be inside the other region?" Then: "Could two segments cross or overlap?" If no three lie on a line, look at the outside boundary. Keep the full case argument back until children seek a guarantee. Lay a fresh sheet of tracing paper over the box for each round.

Share. One example that first looked impossible plus its successful split is a useful stopping point. Call "always works" a conjecture until the group has covered every kind of placement.


Problem 6 three points

Student page 9. Gate: reason through three singleton-versus-segment possibilities.


P6: no split




P7 middle: AB | C



P7 right: AC | B



P7 upper: no split



P6, printed configuration: impossible. The three candidates are A | BC, AB | C, and AC | B. Each leaves a triangle vertex off its opposite segment. This covers every legal split.

Problem 7 crosses for C

Student page 10. Gate: test each cross with all three splits.

Answer: circle the two crosses on the line through A and B: between A and B (AB | C, shared point C) and right of B (AC | B, shared point B). The upper and lower crosses are off that line, so every split fails there.

Every position, for an extension: the whole horizontal line through A and B inside the box (y = 5.6 in adult coordinates), including beyond both ends and at A or B. For three distinct collinear points isolate the middle one; at C = A or C = B separate the coincident labels. Off the line, each singleton lies away from its opposite segment, so no split succeeds.

Held hints for P6 and P7. First: "What are all three possible choices for the single dot?" Then, for P7: "Does C have to be alone?" Ask about the cross beyond B if only the middle cross is circled.

Grades 2-3 Problem 3

Student pp. 4-5 | The printed order is A, D, B, C

Gate: compare complete lists without color-swapped duplicates. The record page supplies six blank copies so children can organize their own search.

AB | CD



AC | BD



ABC | D



ACD | B



Exactly four: AB | CD and AC | BD share segment DB. ABC | D shares D. ACD | B shares B. The shared segment runs from (4.6, 6.3) to (7.8, 6.3), length 3.2 cm at actual size.

Exhaustiveness

The outside labels are A and C, so A | BCD and ABD | C fail. The two inside labels D and B may be isolated, giving ABC | D and ACD | B. Among pairings, AB | CD and AC | BD have overlapping intervals. AD | BC has a gap from D to B and fails. These are the four singleton choices and three pairings, with no duplicates.

Held hint 1. "How can you tell whether a drawing you already have is being counted again?" Let children invent a recording system before offering label strings.

Held hint 2. "What sizes can the two groups have?" If the child proposes 1 + 3 and 2 + 2, use those as their own way to organize the search, rather than handing over the seven-item list immediately.

Why line order matters. Do not reuse K-1 P6's label answer list. Both boards have four winners, but their two inside labels differ. Checking the geometry is more reliable than remembering a string.

Extension. Keep the line but change unequal gaps, then ask whether any successful split changes. As long as the left-to-right label order and distinctness remain the same, the success list stays the same: interval overlap depends on order, not distance.

Grades 2-3 Problems 4 and 5


Student pp. 6-8 | Shared segments and incompatible answer sets

Problem 4 a whole segment


Student page 6. Gate: distinguish a point, a segment, and a triangle with positive area.

Construction: on one horizontal line put A, B, C, D in that order at x = 2, 4, 8, 10 and y = 6. Use AD | BC. Their common region is all of BC, a positive-length segment.


P4: the whole segment BC



P5 first arrangement



P5 second arrangement



Filled triangle is impossible: a partition of four labels has group sizes 1 + 3 or 2 + 2. At least one group has at most two labels, so its hull is a point or a segment. The intersection is a subset of that hull and cannot contain a nondegenerate filled triangle. A narrow triangle still has positive area; line thickness cannot supply it.

Held hints for P4. First: "Could both regions lie on the same straight line?" Then for the impossibility: "How many labels can the smaller group contain?"

Problem 5 no working split in common

Student pages 7-8. Gate: compare complete success sets for two arrangements, not just two chosen successes.

First: A = (2,2), B = (10,2), C = (2,10), D = (4,4). Only ABC | D works. D is strictly inside ABC because $x > 2$, $y > 2$, and $x + y < 12$.
Second: A = (2,2), B = (10,2), C = (10,10), D = (2,10). Only AC | BD works, at (6,6).

The complete success sets {ABC | D} and {AC | BD} are disjoint. The uniqueness proof on guide p. 4 and the full seven-split table on p. 17 verify the stronger requirement. Simply displaying different successful splits would not by itself show that no other split works for both.


Held hints for P5. First: "Keep both arrangements visible. Can your successful split from one work in the other?" Then: "Could you make an arrangement with only one success?" Keep the geometric construction idea in reserve.

Grades 4-5 Problem 1


Student pp. 1-2 | Four exact placements

Gate: identify a shared point using filled hulls and ideal centers. A complete catalog is unnecessary here, but the adult should know where a second answer exists. Each placement is drawn on its own record board on p. 2.


Upper-right: AD | BC




Interior: ABC | D



On AB: AB | CD



Left: AC | BD



Complete success sets: {AD | BC}, {ABC | D}, {AB | CD, ABC | D}, {AC | BD}, in the order shown. At the interior point the shared point is D. At the boundary point both winners share D. The other two winners share the actual diagonal crossing.

Exact boundary check. A = (2,2), B = (10.5,3), and D = (6.25,2.5) is their midpoint. The left cross D = (1,6) is outside ABC and not on AC. General-position uniqueness and the boundary argument explain the counts 1, 1, 2, 1.

Held hints. First: "Which hull includes a dot without its being a corner?" Then: "Try both group-size patterns." Ask for an actual common point rather than a visual impression that the figures are close.

Grades 4-5 Problems 2 and 3

Student pp. 3-5 | Enumeration and a placement challenge

Problem 2 the sloping line

Student pages 3-4. Gate: recognize collinearity independent of page orientation; organize a complete list.

Printed order: A, C, B, D along the diagonal. The small gallery below is rotated flat to make the order easy to compare; no point order changes.

AB | CD



AD | BC



ABD | C



ACD | B



Exactly four: AB | CD and AD | BC share segment CB, from (5,6) to (8,9). ABD | C shares C; ACD | B shares B. The shared segment has length 3 times the square root of 2 cm; no length calculation is needed.

Completeness: endpoints A and D cannot stand alone successfully; middle points C and B can. Pairings AB | CD and AD | BC overlap, while AC | BD leaves a gap. Thus the other three candidates, A | BCD, ABC | D, and AC | BD, fail.

Held hints for P2. First: "Which locations are the two ends?" Then: "How many singleton choices and pairings are possible?" Do not sort the labels alphabetically along the line.

Problem 3 challenge an opponent

Student page 5. Gate: test, record, and distinguish a conjecture from a guarantee.

Answer: the opponent can always find a successful split. There is no defeating four-point arrangement. The complete constructive strategy is on guide p. 18; it works for arbitrary placements, not only the printed instances.

Held hints for P3. First: "Does any dot lie in the region of the other three?" Then: "If all four are outside corners, what happens to the opposite-corner pairs?" If labels line up or coincide, ask about the exact point or segment shared.


Keep the problem open. Do not distribute the adult case checklist yet. Save attempted counterexamples as evidence for P6. If a child finds a success for each challenge, invite the partner to change the arrangement in a genuinely different way.

Grades 4-5 Problem 4


Student p. 6 | Coincident A and D are still separate labels

Gate: distinguish a label from its location. Stacking A and D preserves both labels; it does not replace them by one item. The page has six small boards; their number does not give the count.


A | BCD




AB | CD



AC | BD



ABC | D



Exactly four for the printed board: all four splits shown, each sharing the location A = D = (3,3). They are precisely the ways to put A and D into different groups, up to swapping group names. Assign B and C independently to either side to obtain four choices.

Why the remaining three fail: in AD | BC the location A = D is off segment BC. In ABD | C, C is off segment AB. In ACD | B, B is off segment AC. The three distinct locations form a noncollinear triangle.

General answer: yes, a success always exists when any two labels coincide. Put one of that pair alone and the other with the remaining labels; both hulls contain their common location. The other points may lie anywhere or also coincide. The exact count need not remain four: if all four labels share one location, all seven splits work.


Held hints. First: "Can A and D belong to different groups even though they share a place?" Then: "Where would both regions have to contain a point?" Keep the mark centered and both label names readable.

Grades 4-5 Problem 5


Student pp. 7-9 | Revised requirement: different unique successes

Gate: distinguish exactly one successful split from having found only one. Both first arrangements must have one winner, and their winning splits must differ. A mere translated copy does not meet the revised condition.


First: only ABC | D



Second: only AC | BD



Third: four successes



Reproducible constructions: first A(2,2), B(10,2), C(2,10), D(4,4); second A(2,2), B(10,2), C(10,10), D(2,10); third A(2,6), B(4,6), C(8,6), D(10,6). All lie inside the supplied board. Children may make other correct examples.

All seven unordered splits (yes/no columns are exhaustive).

Split	First	Second	Third
A | BCD	no	no	no
AB | CD	no	no	no
AC | BD	no	yes	yes
AD | BC	no	no	yes
ABC | D	yes	no	no
ABD | C	no	no	yes
ACD | B	no	no	yes

For the first example D lies strictly inside ABC, and the common point is D. For the second, the two diagonals meet at (6,6). The uniqueness arguments on guide p. 4 rule out the other six splits in each. In the third, AC | BD and AD | BC share BC, ABD | C shares C, and ACD | B shares B.

Held hint 1. "How could you check that there is no second answer?" Ask for a systematic list before revealing a new geometry.

Held hint 2. "Can your two drawings have different group sizes in their only successful split?" This suggests a meaningful contrast without prescribing coordinates.


Assessment. A valid solution supplies three arrangements, checks the two different one-element success sets, and shows more than one winner for the third. For the printed request to explain the number, an exhaustive check is stronger than displaying only two winners.

Grades 4-5 Problem 6


Student p. 10 | A complete proof for every arrangement

Gate: a reason must cover unseen placements. The following cases are an adult proof key. Invite children to organize their own examples before sharing this organization.

Repeated location




Three on one line




1. Repeated location. If two labels share a location, put one alone and all other labels together. Both hulls contain that location. This also covers three or four coincident labels.

2. Distinct locations with a collinear triple. Of those three locations, one is between the other two. Isolate that middle label. The hull of the remaining three contains the segment joining the two endpoints, hence contains the middle point. The fourth location can be anywhere. This includes all four collinear.

Triangle outside boundary



Four outside corners



3. Distinct locations and no collinear triple. Their convex hull has either three or four corners. It cannot have one or two without collinearity, and there are only four points. With three corners, the remaining point is strictly inside their triangle; isolate it. With four corners, pair opposite corners. The diagonals meet inside the quadrilateral: opposite corners lie on opposite sides of either diagonal's supporting line, so the two closed diagonal segments cross.

Conclusion: every arrangement is in one of these cases; in each, every label is used exactly once in two nonempty groups, and an actual shared point is identified. Thus every four-label planar arrangement has a successful split. Boundary contact is enough; positive area is never promised.


Held hints: "What kinds of examples have we not covered?" Then, only if needed: "Could labels coincide? Could three line up? How many outside corners remain otherwise?" A list of many experiments remains evidence for a conjecture, not the universal proof.

Grades 4-5 Problems 7 and 8


Student p. 11 | Three dots fail; the smallest guaranteed number is four

Gate: separate "some arrangement works" from "every arrangement works." Use a counterexample to disprove a guarantee, and the previous proof to establish one.


A | BC fails



AB | C fails



AC | B fails



Problem 7: the printed three points have no successful split. Each of the three legal splits is drawn. A singleton is a triangle vertex, and the other hull is the opposite closed side. No vertex lies on its opposite side. These three possibilities exhaust the 1 + 2 group sizes.

Exact noncollinearity check: A = (2,3), B = (10.3,4), C = (4.8,10.1). The determinant (B - A) cross (C - A) equals 5613/100, which is positive and nonzero. The printed triangle is genuinely noncollinear. Children can use its visible geometry; determinants are adult audit only.

Problem 8: why four is the sharp threshold

Four: every arrangement works by the complete argument on guide p. 18.
Three: the printed noncollinear triangle is one arrangement that fails, so three cannot guarantee success.
Two: take two distinct locations. The only split puts one point in each group; their hulls are disjoint.
One: no partition into two nonempty groups exists.

Thus four is the smallest positive number guaranteeing a successful split in the plane. Some three-point arrangements do work, such as a collinear triple, and two coincident labels work too. Those examples do not establish a guarantee for every arrangement.

Held hint for P7. "How many possible ways can we split three labels into nonempty groups?" Let children draw all three before declaring impossibility.

Held hint for P8. "To show that three does not guarantee success, must every three-point arrangement fail?" One clear counterexample is enough. Then check two and one to justify the word smallest.

Extension. Ask whether five or more points always work. Choose four, make a successful split, then add each remaining label to either group. Hulls can only grow, so the old shared point remains. This is a short extension of the four-point proof.

Extensions and fidelity limits

Keep extensions optional and matched to prerequisites

Ready-to-use extensions with answers

How much overlap? Gate: compare dimension and group sizes. Ask whether four labels can ever give positive-area overlap. Answer: no, by the smaller-group argument in middle P4. Five can create a shared segment between a triangle and a segment; six can create positive-area overlap using two overlapping triangles. No claim of a universal guarantee is intended for these constructions.

One-dimensional threshold. Gate: middle points and the meaning of always. Restrict all positions to a line. Three labels always work: isolate the middle of three distinct locations, or separate coincident labels. Two distinct locations fail. The sharp threshold on a line is three.

Maximum number of successes. Gate: the seven-case catalog. Seven is an upper bound because there are only seven splits. It is attained when all four labels coincide. For four distinct collinear locations exactly four work, independent of spacing. General-position examples have exactly one.

Prescribed partition. Gate: construction and counterexample. Could one fixed label split work for every placement? No. The inside-triangle example has only ABC | D; the square example has only AC | BD. Their disjoint answer sets already disprove a universal fixed split.

What this session faithfully claims

The mathematical payoff is the planar four-point convex-partition theorem, including its sharp threshold. The exact planar proof, fixed-instance checks, and uniqueness examples support that claim. The sequence does not establish higher-dimensional Radon theory, Tverberg's theorem, algorithms for arbitrary point clouds, or fractional-coordinate methods as child prerequisites.

The guide diagrams are scaled answer diagrams, not full-size replacement student boards. Color, a visible line width, and rings are display conventions, not mathematical area. A shaded triangle always includes its boundary. Labels, rather than distinct locations, are partitioned.

The student revisions add record spaces and require different unique successes in upper P5. The 6 November 2026 kid-fit revision (v7) moved the games forward, lightened the K-1 pages, asks K-1 P5 and middle P7 which crosses for C work (every position is an adult extension), and split the three-dot questions from those that follow them. The earlier reviews assessed preceding drafts. This guide rechecked the final coordinates and new construction condition; review evidence is not a classroom pilot.

After the first use

Record which problems children actually used, where adults had to restate the action, whether filled hulls and singleton regions were understood, whether recording boards helped, whether K-1 children could manage the optional P4 and P6, and whether upper P6 reached a proof or a conjecture. Keep observations separate from guesses about causes. Revise pacing only after evidence; do not label this sequence classroom-tested merely because the mathematics was checked.

Sources and verification

Source roles, exact audit, and portable editable files

Verified mathematical source

Jean Gallier and Jocelyn Quaintance, Aspects of Convex Geometry, author-hosted version dated 9 October 2025, section 3.5, Theorem 3.10 and Figure 3.10, printed pp. 75-77. Verified from the theorem and proof body on 3 October 2026.
https://www.cis.upenn.edu/~jean/combtopol.pdf

The source gives Radon's theorem in arbitrary finite dimension and illustrates the planar four-point cases. It supplies adult mathematical context. This guide's elementary planar case proof, coincidence handling, all student-instance answers, and construction checks were worked out directly. The unverified Hug-Weil citation from older planning material is not used as evidence.

Teaching context actually consulted

Natasha Rozhkovskaya, Math Circles for Elementary School Students, "Introduction: Berkeley 2009," in the locally held MSRI/AMS book. That introduction describes adult assistance for each child and active parent involvement in the elementary circle. It supports the use of close adult help; it does not report a trial of this convexity activity.

Repository AGENTS.md, README.md, the Week 22 activity prompt, the general and mathematics reviews, and the final v3 sources and PDFs were read. These supply the actual group, student-page conventions, materials, and revision history. The proposed timing, held hints, route choices, and stopping points are facilitator design suggestions, not claims extracted from the source book.

What was checked independently

A separate exact-rational checker examines every unordered split of all 31 fixed, three-point, and construction configurations. It uses direct point-on-segment, point-in-triangle, and segment-intersection predicates rather than importing the student hull checker. It records exact shared-point or shared-segment endpoints. It also checks 6,561 labeled placements on a 3 by 3 test grid, including coincident points, as a stress test.

The new inside-triangle and square witnesses each have one successful split and those splits differ. All three collinear boards have four successes with their own label orders. The final student page counts are 6, 10, and 11 (v7). The universal existence claim rests on the complete proof on guide p. 18, not on the finite tests.

Editable source and rebuild

The companion facilitator-src folder contains build_guide.py (editable text and drawing source), check_math.py, geometry-results.json, student-input-manifest.json, checks.txt, README.md, and build.sh. Run ./build.sh from that folder with Python 3, ReportLab, pypdf, and Poppler installed. It writes the guide one level above and renders review images locally.

The final guide was rendered and every page visually inspected. The student PDFs were read and rendered for matching only; no student page or source was changed by this separate guide step. The 6 November 2026 kid-fit revision made the v7 packets; this guide (F22-FAC-v7) was renumbered to match. SHA-256 hashes in the input manifest identify the exact matched student PDFs.

Starting routes and return visits

Planning update of 4 October 2026. The timing menus are examples; a theme can continue over several meetings. Keep the full Grades 4-5 investigations and choose by readiness.

Use and return choices

Keep one adult at each table to read and record while children choose groups and placements. The middle return route, the three-dot pages P6-7, is marked on guide p. 3; they can also follow the game (P2) directly. Other return choices: K-1 can replay the games (P2 or P3), then add the three-label crosses question (P5) when ready; the dots-in-a-row page (P6) is optional for K-1. Grades 4-5 can use saved splits to examine coincident labels and multiple winners (P4-5), then make the all-arrangements, three-dot and minimum-number arguments (P6-8). Keep the two regions filled and the shared point explicit.