Mathematical overview: a distance between whole arrangements

The model. Equally spaced bins form an open line. The initial and target piles have the same number of interchangeable counters. Moving one counter across one neighboring gap costs one. Bins have unlimited capacity; counters may pass through occupied bins. A long delivery costs its number of gaps, and carrying a handful costs separately for each counter. The target remains fixed.

Every gap has unavoidable net traffic. Write initial and target counts as $a_0, \ldots, a_m$ and $b_0, \ldots, b_m$. Across the gap after bin i, right crossings minus left crossings must equal

$$S_i = \sum_{j=0}^{i} (a_j - b_j).$$

Consequently every plan costs at least $\sum_{i=0}^{m-1} |S_i|$. Children can say "two must cross left" without signed notation. Total traffic may exceed net traffic; counting only net crossings gives a lower bound, not an arbitrary plan's cost.

All gap bounds can be attained together. List initial counter positions $x_1 \le \cdots \le x_N$ and target slots $y_1 \le \cdots \le y_N$, retaining repeated positions. Match x_k to y_k. Opposite traffic across a gap would require an earlier initial counter to have a later target than a later initial counter, contradicting the ordering. Thus this plan costs exactly

$$\sum_k |x_k - y_k| = \sum_i |S_i|.$$

This proves optimality, not uniqueness. The nearest available *whole delivery* need not be part of an optimal plan: positions $0, 3$ to slots $2, 5$ give nearest-first cost $1 + 5 = 6$, while $0 \to 2$ and $3 \to 5$ cost $2 + 2 = 4$.

Minimum work is a metric. Zero work means identical piles; reversing moves gives symmetry; concatenating plans proves the triangle inequality. Equal average position does not mean equal piles: two counters at each of 0 and 4 and four counters at 2 both have position total 8, but their distance is 8. For probability measures of $N > 0$ counters each of mass $1/N$, Wasserstein-1 distance is our work divided by N.

A tree extension. Removing an edge splits a tree in two. The absolute initial-minus-target imbalance on one side forces that many crossings of the edge. Summing over edges gives the exact optimum: orient each nonzero required flow toward its deficit side; because the graph is acyclic, these flows can be decomposed into surplus-to-deficit paths. Cycles need a different argument because one removed edge need not separate the graph.

Readiness and discovery. Grades 2–3 can count small piles and moves and challenge a greedy rule. Grades 4–5 adds lower-bound certificates and later equal-center or branching-road questions. The student retains delivery choices even when an adult reads or records. Experiments propose a best route; matching an attaining route to an unavoidable-crossing bound proves it. No K–1 coverage, unequal-mass transport, road-capacity constraint or squared-distance objective is implied.

Preparation, shared launch and pacing

Status. Unpiloted review draft; physical printing, handling, preparation and timing are untested. The older-table packet has substantial continuations; finishing every page is not the goal.

K–1 route, adult-led. No K–1 packet is printed. Use the line mat (`week-99-materials.pdf`, page 1) with two or three counters; the adult sets each start and target and counts moves aloud with the child. First build Grades 2–3 Problem 1's two pairs on the mat: "Make it match. Can you use fewer moves?" Then gather two counters, one in bin 0 and one in bin 4, into bin 2. End with the game at 3 moves: one child puts two counters in the target row, and the other puts two counters in the moving lane so that matching takes exactly 3 moves. Key in plain words: $00 \to 24$ takes 6 moves and $03 \to 12$ takes 2; gathering 0 and 4 at 2 takes 4 moves; every two-counter target has a start exactly 3 moves away (start at the target and move counters three steps toward the farther end of the mat). That K–1 children keep counting moves, not hand motions, is a prediction.

Per pair. The line mat, page 1 of `week-99-materials.pdf`: six 30 mm bins 35 mm apart in a moving lane, a target row below it and a move-record strip. Add 8–12 counters up to 2 cm wide, which stack in a bin, in two colors if the target is set with counters. Keep the target outside the moving lane. Use only one recoverable record, such as writing each crossed gap and counting the entries afterward. Do not require simultaneous route, total and prefix tallies. Optional removable counter labels identify a planned pairing but never restrict which target a counter may fill. For later branching work, the star board is page 2 of the materials file: 35 mm circles for moving and a small star for the target picture. For upper Problem 4, print an extra copy of page 4 and cut its eight catalog cards before the session.

Staffing and preparation. Three older-table kits plus a demonstration mat; beyond printing, allow about 10 minutes to count counters and cut the upper Problem 4 cards, subject to rehearsal. Partners alternate mover and checker; the upper trio rotates a third recorder. An adult can read pictured piles or record a child's plan. The child should still choose where to deliver. If the adult must plan each move, reduce the number of counters and keep two genuinely different choices.

Complete launch. After brief free handling, use bins 0–3 of a line mat, with the target in its target row. The picture below is a non-task example. Move 0 to 1, then 2 to 3, preserving the target throughout.

	0	1	2	3
Start	1	0	1	0
After 0 to 1	0	1	1	0
After 2 to 3	0	1	0	1
Fixed target	0	1	0	1

The record is $0 \to 1$, $2 \to 3$: two counter-steps. Show that carrying one counter directly from 0 to 2 costs two even in one hand motion. A counter already in a useful place may stay there. Do not teach ordered matching or the gap formula before the children try their own deliveries.

A first visit. Five minutes movement, five minutes launch, then 20–25 minutes optimizing small pile changes and trying the nearest-first question. Use 10–15 minutes to defend one proposed optimum: "What must cross this gap, however clever the plan?" Save the full metric, equal-center catalog or branching road for another visit. First route: Problems 1–3 in both bands; return visits: Problems 4–6. One clear lower bound and a matching construction are already substantial mathematics.

Hints kept with adults. Cover the right side of a selected gap and compare the starting and target totals still visible. Keep zero piles in the picture. If counters cross the same gap in both directions, ask whether a different assignment could avoid that waste. A working plan supplies only an upper bound until an impossibility argument matches it. Equal-cost different plans are acceptable. Do not silently replace counter-steps with number of deliveries or number of hand motions.

Line deliveries: exact answers and matching lower bounds

Here a position row lists one bin number per counter, with repetitions. Thus 0055 means two counters in bin 0 and two in bin 5; it is not a four-digit integer. Match the first starting position to the first target, second to second, and so on. Each listed plan attains its gap bound. The gap row gives signed net crossings across gaps 0–1 through 4–5; negative means left. Use ordinary "must cross left/right" language with children.

Printed problem	Start	Target	Net gap crossings	Work
2–3 P1, first	00	24	$2, 2, 1, 1, 0$	6
2–3 P1, second	03	12	$1, 0, -1, 0, 0$	2
2–3 P2 first; 4–5 P1 first	0005	1234	$3, 2, 1, 0, -1$	7
2–3 P2 second; 4–5 P1 second	1144	0225	$-1, 1, -1, -1, 1$	5
2–3 P2, third	2222	0134	$-1, -2, 2, 1, 0$	6
4–5 P2, first	0055	2233	$2, 2, 0, -2, -2$	8
4–5 P2, second	1114	0235	$-1, 2, 1, 0, 1$	5
4–5 P2, third	0235	1144	$1, -1, 0, 1, -1$	4

Reading one proof. For the first $0005 \to 1234$ delivery, three counters must cross gap 0–1 to the right, two across 1–2, one across 2–3, none net across 3–4, and one across 4–5 to the left. At least $3 + 2 + 1 + 0 + 1 = 7$ counter-steps are required. Send the three counters at 0 to 1, 2 and 3, and the counter at 5 to 4: costs $1 + 2 + 3 + 1 = 7$. The lower and upper bounds agree.

Different plans. Different orders of the same moves give the same work. In every printed pair the best matching is unique once identical counters are ignored; different matchings can tie elsewhere, for example $01 \to 12$ by $0 \to 1$, $1 \to 2$ or by $0 \to 2$ with the counter at 1 staying. Inserting a needless out-and-back detour is a different plan but not another best plan.

A child-sized certificate. Pick one gap, cover the bins to the right, and compare the starting and target number of counters on the left. The difference cannot vanish unless that many net crossings occur. Repeat for enough gaps to account for the full claimed cost. A single-gap bound can be useful but need not prove the complete optimum. Keep the target visible and unchanged while counting.

Why the sorted plan attains all bounds. Suppose one paired counter crosses a gap rightward and another leftward. Their starting positions satisfy $x \le i < x'$, but their target positions satisfy $y > i \ge y'$. This reverses the target order, impossible in the sorted pairing. With no opposite traffic, total crossings at each gap equal its compulsory net number in absolute value. Summing counts each counter-step exactly once.

Greedy choices, invented starts and three arrangements

Both bands Problem 3, page 3: nearest delivery can fail

In the first pair, starts are 0 and 3; target slots are 2 and 5. The uniquely shortest delivery is $3 \to 2$, cost 1. It leaves $0 \to 5$, cost 5, for total 6. The other pairing costs $0 \to 2$ plus $3 \to 5$, or $2 + 2 = 4$. The latter meets the gap bound. A greedy choice that looks best locally need not leave a good remaining problem.

The Grades 2–3 second pair is starts 0,4 to slots 2,5. Nearest-first sends $4 \to 5$ and then $0 \to 2$, total 3, which is optimal. This contrast matters: the rule sometimes succeeds. The other pairing costs $4 \to 2$ plus $0 \to 5 = 7$.

The Grades 4–5 second pair is starts 0,4 to slots 3,5. Two shortest deliveries tie at cost 1. Choosing $4 \to 3$ leaves $0 \to 5$, total 6; choosing $4 \to 5$ leaves $0 \to 3$, total 4. Ties can therefore change the outcome. A delivery is committed to an unfilled target slot; moving one counter repeatedly or treating a temporary transit bin as a new supply changes the proposed rule.

Grades 2–3 Problem 4, page 4: work 4 and work 7

The fixed target is four counters at bin 2. Every initial counter must end at 2, so the exact work is the sum of its distance to 2. Examples of start position rows with work 4 are 0022, 1111 and 0224; examples with work 7 are 0003, 0014 and 0115. Their distances to bin 2 sum to the requested value. Children may find many others: the checker finds 16 work-4 starts and 19 work-7 starts, counting pile arrangements rather than labeled-counter identities. Exhausting those sets is not the printed task.

Grades 2–3 Problem 6, page 6: can a target defeat work 5?

No. Every target of four counters on bins 0–5 has an arrangement at distance exactly 5. If its position sum is s, moving all counters to 0 costs s and moving them all to 5 costs $20 - s$. One of these is at least 10. Starting at the target, move counters toward that endpoint for exactly five adjacent steps. All moves go in one direction, so the gap lower bound is exactly 5. Reverse those five moves to deliver from the constructed start to the target. The independent check also tests all 126 four-counter arrangements. The argument, rather than the finite sample alone, explains why the challenge always works.

Grades 4–5 Problem 6, page 6: a detour cannot beat the optimum

No: an optimal A-to-B plan followed by an optimal B-to-C plan is a valid A-to-C plan. Therefore $D(A, C) \leq D(A, B) + D(B, C)$. Equality example: four counters at 0, then four at 1, then four at 2 gives $4 + 4 = 8$. Strict example: four at 0, then four at 5, then four at 1 gives $20 + 16 > 4$. All three arrangements are different in both examples. The direct route means the least work between whole arrangements, not necessarily one hand movement. This is the triangle inequality for the new distance.

Grades 4–5 Problem 4: same center, different distributions

The eight movable cards are all four-counter arrangements on bins 0–4 whose position total is 8. A position total repeats a bin number once per counter: four counters at 2 contribute $2 + 2 + 2 + 2 = 8$. Before this unfamiliar convention, a non-task example with one counter at 1 and one at 3 gives total 4. Equal totals and equal numbers of counters mean equal average position, but they do not make the pile pictures identical.

Label the printed cards in reading order:

card	A	B	C	D	E	F	G	H
positions	0044	0134	0224	0233	1124	1133	1223	2222

Their exact minimum-work distances are below. Each entry is the sum of the four coordinate differences in absolute value, justified by the sorted-pairing theorem.

	A	B	C	D	E	F	G	H
A	0	2	4	4	4	4	6	8
B	2	0	2	2	2	2	4	6
C	4	2	0	2	2	4	2	4
D	4	2	2	0	4	2	2	4
E	4	2	2	4	0	2	2	4
F	4	2	4	2	2	0	2	4
G	6	4	2	2	2	2	0	2
H	8	6	4	4	4	4	2	0

The maximum is 8, attained only by A and H as an unordered card pair: two counters at 0 and two at 4 versus four at 2. A complete comparison of the 28 distinct pairs proves this finite maximum. The bound "each can gather at 2 in at most 8" would only give 16 by concatenation and is not enough to prove the claimed maximum.

Why no card is missing. Sort the four positions $a \le b \le c \le d$. If $a = 0$, then $b = 0, 1, 2$ give respectively 0044, 0134, and 0224 or 0233; $b \ge 3$ makes the sum too large. If $a = 1$, then $b = 1$ gives 1124 or 1133, and $b = 2$ gives 1223; $b \ge 3$ is too large. If $a = 2$, all entries must be 2. Larger a is impossible. This proves the eight-card inventory without treating physical card order as a different arrangement.

A short reason no pair needs more than 8. On every card, bin 0 and bins 0–1 each hold 0, 1 or 2 counters, and bins 0–2 and bins 0–3 each hold 2, 3 or 4. So the counts left of any of the four gaps differ by at most 2 between two cards, at most 2 counters must cross each gap, and no pair needs more than $4 \times 2 = 8$. Children can check the counts on the cards.

Pacing and support. The essential discovery is that two distributions with the same center can still require work. The full maximum and completeness proof are later depth. Pairs can share comparisons and pool a checkable record; each child need not do 28 calculations or transcribe every card. The supplied cards carry the cases so sorting and comparison remain the mathematics.

Branching roads, extensions and source limits

Both bands Problem 5, page 5

The star has center O and three leaves A,B,C. Each complete road costs one per counter; road length on paper is irrelevant. There is no shortcut directly between leaves.

The common first start has two at A and one at C; the target has two at B and one at O. Send A–O–B twice and C–O once: work 5. The A, B and C edges force respectively 2,2,1 crossings, proving optimality.

The Grades 2–3 second start has two at A and one at B; the target has one at A and two at C. Leave one at A, send A–O–C and B–O–C: work 4. Leaf-edge requirements are 1,1,2. An invented three-counter example of work 6 puts all three at A initially and all three at B in the target. All three traverse two edges, and each of those two leaf edges forces three crossings.

The Grades 4–5 second start has three at A and one at B; the target has one at A, two at C and one at O. Leave one at A, send two from A to C, and send B to O: $4 + 1 = 5$. The leaf-edge requirements are 2,1,2. Any other optimal plan is equally valid.

Why the star formula always works

Each leaf's imbalance must cross its only road. Send excess counters at leaves to O; then send counters from O to deficient leaves. Together with any initial counters already at O, there are enough because total initial and target mass agree. Remaining counters at O match its target automatically. No edge is used in opposite directions. Thus the sum of the absolute leaf imbalances is both a lower bound and attainable, for every equal-total start and target on this star.

For a general tree, the same argument can be applied by removing leaves successively, or by decomposing the uniquely required edge flows into surplus-to-deficit paths. A cycle has multiple routes: removing a single edge may leave the graph connected, so this exact cut argument does not apply unchanged. For example, a counter from 0 to 3 on a four-bin ring can use one road; on the open 0–1–2–3 line it needs three.

Later routes and connection boundaries

The entire pile arrangement is the point in this new geometry; minimum counter-work is its distance to another arrangement. Week 2's capacity-one swap puzzles are a genuine restricted relative, while these bins allow piles and passage through occupied bins. Labeled-object sorting, edge-disjoint routes and buying a cheapest network optimize different objects or costs. Week 89's running totals can support the gap certificate, without being the present question.

Peyré and Cuturi, *Computational Optimal Transport*, §2.6, Remarks 2.28 and 2.30, equations (2.33) and (2.37), give the ordered-position and cumulative-distribution formulas; §6.3 discusses graphs. The proofs here specialize those ideas to whole counters. No general higher-dimensional or unequal-mass solution is claimed. See `SOURCES.md` for exact links and provenance.

Independent checks cover the printed inputs, greedy alternatives, catalog, invented-distance challenge and star examples. General arguments distinguish proof from experimentation. The package QA records the final page inspection and clean ZIP rebuild; those are digital checks. Printing, material handling, pacing and classroom success remain untested. Record where children keep the planning decisions and where adult support becomes too directive.