

quality
organ been used as
nutritional interac-
d dietary protein
es reduce rates of
protein levels lead
through compensa-
growth rates. In
h-instar larvae of
ow protein levels
of 14 and 42°C,
tion of the fifth
perature range
actions between
(Kingsolver and
st that compens-
less effective in
es than in the
ay are commonly
ed by examining
4, and 5 separ-
king differences
(0). Mean growth
the first three
g the fifth instar
which laboratory
ed). Fifth instar
nsitive to high
in view of the
illar nutritional
et al. 2000). The
est insect which
ing leaf nitrogen
ures. Lindroth
s of temperature
al performance
e fourth instar,
over the larval
nation of poor
re can result in
g life cycles, as
ora *groenlandica*
opment time of
basking behav-
in more complex
rophic levels are

involved. Stamp (1990) measured growth of caterpillars (*M. sexta*) as influenced by interactions between temperature, nutrients, and a common phenolic, rutin. Numerous interactions were found, the most important being that the effects of diet dilution and rutin on larval growth rate were a function of temperature. More time was spent moulting at low temperatures and when rutin was included in the diet, resulting in lower growth rates. In temperate regions, interactions of temperature, food quality, and natural enemies are most likely

to affect gregarious spring-feeding caterpillars. In contrast, solitary behaviour reduces predation risk but forces caterpillars to forage in microhabitats that are suboptimal in terms of food quality and temperature (for review see Stamp 1993). Finally, the interplay between temperature and nutrition is critical in the context of global climate change. Reduced foliar nitrogen is a consistent response of plants grown in enriched CO₂ atmospheres (Ayres 1993; Lincoln et al. 1993; Watt et al. 1995). Insect performance may be altered by the

Figure 2.17 The thermal sensitivity of growth and feeding in *Manduca sexta* caterpillars, illustrated by performance curves measured over the temperature range 14–42°C for (a) mass-specific growth rate, (b) mass-specific consumption rate, (c) proteolytic digestion rate, (d) methionine absorption rate, and (e) mass-specific respiration rate.

