Lecture 17: Orthogonal iteration and QR iteration

Jamie Haddock

Table of contents

1 Orthogonal iteration and QR iteration **1**
1.1 Schur decomposition . . . 1
1.2 Orthogonal iteration . . . 3
1.2.1 Convergence to the Schur decomposition . . . 5
1.2.2 Convergence rate . . . 6
1.3 QR iteration . . . 7
1.3.1 Symmetric case . . . 10
1.3.2 Improvements to QR iteration . . . 11

1 Orthogonal iteration and QR iteration

Power iteration finds just one eigenpair at a time, and only the dominant one. Can we find *several* eigenvalues at once, or even all of them? In this lecture we run power iteration on a whole block of vectors simultaneously, and add one extra ingredient, orthogonalization, to keep the vectors from all collapsing onto the same dominant eigenvector. The result is the **QR iteration**, which (with some important improvements) is the basis of the standard algorithm for computing all the eigenvalues of a matrix.

Assumption:

Throughout this lecture, we assume that $\mathbf{A}$ is a diagonalizable $n \times n$ matrix whose eigenvalues satisfy

$$|\lambda_1| > |\lambda_2| > \cdots > |\lambda_n|.$$

We write $\mathbf{A} = \mathbf{V}\mathbf{D}\mathbf{V}^{-1}$, where $\mathbf{V} = \begin{bmatrix} \mathbf{v}_1 & \cdots & \mathbf{v}_n \end{bmatrix}$ holds the eigenvectors and $\mathbf{D} = \operatorname{diag}(\lambda_1, \ldots, \lambda_n)$.

For a real matrix $\mathbf{A}$ this assumption forces every eigenvalue to be real, since complex eigenvalues come in conjugate pairs and conjugates have equal modulus. So all of the computations below can be done in real arithmetic, with real orthogonal matrices in place of unitary ones and $\mathbf{Q}^* = \mathbf{Q}^\top$.

1.1 Schur decomposition

Power iteration produces an *eigenvector*. The natural target of the algorithms in this lecture is a different object, one that exists for every square matrix and is much better behaved numerically than the eigenvector matrix $\mathbf{V}$.

Definition: Schur decomposition

A **Schur decomposition** of an $n \times n$ matrix $\mathbf{A}$ is a factorization

$$\mathbf{A} = \mathbf{Q}\mathbf{T}\mathbf{Q}^*,$$

where $\mathbf{Q}$ is unitary ($\mathbf{Q}^*\mathbf{Q} = \mathbf{I}$) and $\mathbf{T}$ is upper triangular. The columns of $\mathbf{Q}$ are called **Schur vectors**.

Compare this with the eigendecomposition $\mathbf{A} = \mathbf{V}\mathbf{D}\mathbf{V}^{-1}$. We have traded a *diagonal* matrix for a merely *triangular* one, but in return $\mathbf{Q}$ has orthonormal columns (so $\kappa_2(\mathbf{Q}) = 1$), whereas $\mathbf{V}$ can be badly conditioned, or fail to exist at all if $\mathbf{A}$ is not diagonalizable.

Theorem:

Let $\mathbf{A}$ be an $n \times n$ matrix.

1. $\mathbf{A}$ has a Schur decomposition $\mathbf{A} = \mathbf{Q}\mathbf{T}\mathbf{Q}^*$.
2. The diagonal entries of $\mathbf{T}$ are the eigenvalues of $\mathbf{A}$.
3. For each $j = 1, \ldots, n$, let $\mathbf{Q}_j$ be the first j columns of $\mathbf{Q}$ and $\mathbf{T}_j$ the leading $j \times j$ block of $\mathbf{T}$. Then $\mathbf{A}\mathbf{Q}_j = \mathbf{Q}_j\mathbf{T}_j$. That is, $\text{range}(\mathbf{Q}_j)$ is an *invariant subspace* of $\mathbf{A}$: $\mathbf{x} \in \text{range}(\mathbf{Q}_j)$ implies $\mathbf{A}\mathbf{x} \in \text{range}(\mathbf{Q}_j)$.

Proof sketch: (Part 1: A has a Schur decomposition $\mathbf{A} = \mathbf{Q}\mathbf{T}\mathbf{Q}^*$.)

We use induction on n, and assume a matrix of size $n-1 \times n-1$ has a Schur decomposition.

Let $(\lambda, \mathbf{v})$ be an eigenpair of $\mathbf{A}$ with $\|\mathbf{v}\|_2 = 1$ (one exists over $\mathbb{C}$), and extend $\mathbf{v}$ to a unitary matrix $\mathbf{U} = \begin{bmatrix}\mathbf{v} & \mathbf{U}_2\end{bmatrix}$. Because $\mathbf{U}^*\mathbf{A}\mathbf{v} = \lambda\mathbf{U}^*\mathbf{v} = \lambda\mathbf{e}_1$,

$$\mathbf{U}^*\mathbf{A}\mathbf{U} = \begin{bmatrix}\lambda & \mathbf{w}^* \\ \mathbf{0} & \mathbf{A}_2\end{bmatrix}$$

for some vector $\mathbf{w}$ and some $(n-1) \times (n-1)$ matrix $\mathbf{A}_2$.

By induction, $\mathbf{A}_2 = \mathbf{Q}_2\mathbf{T}_2\mathbf{Q}_2^*$ with $\mathbf{Q}_2$ unitary and $\mathbf{T}_2$ upper triangular. Setting $\mathbf{Q} = \mathbf{U}\begin{bmatrix}1 & \\ & \mathbf{Q}_2\end{bmatrix}$ (a product of unitary matrices),

$$\mathbf{Q}^*\mathbf{A}\mathbf{Q} = \begin{bmatrix}\lambda & \mathbf{w}^*\mathbf{Q}_2 \\ \mathbf{0} & \mathbf{T}_2\end{bmatrix},$$

which is upper triangular.

Proof sketch: (Part 2: The diagonal entries of T are the eigenvalues of A.)

For part 2, $\mathbf{T} = \mathbf{Q}^*\mathbf{A}\mathbf{Q}$ is similar to $\mathbf{A}$, so it has the same eigenvalues, and the eigenvalues of a triangular matrix are its diagonal entries.

Proof sketch: (Part 3: For each $j = 1, \ldots, n$, let $\mathbf{Q}_j$ be the first j columns of $\mathbf{Q}$ and $\mathbf{T}_j$ the leading $j \times j$ block of $\mathbf{T}$. Then $\mathbf{A}\mathbf{Q}_j = \mathbf{Q}_j\mathbf{T}_j$. That is, $\text{range}(\mathbf{Q}_j)$ is an *invariant subspace* of $\mathbf{A}$: $\mathbf{x} \in \text{range}(\mathbf{Q}_j)$ implies $\mathbf{A}\mathbf{x} \in \text{range}(\mathbf{Q}_j)$.)

For part 3, take the first j columns of $\mathbf{A}\mathbf{Q} = \mathbf{Q}\mathbf{T}$. Since $\mathbf{T}$ is upper triangular, its first j columns are zero below row j, so the right-hand side is $\mathbf{Q}_j\mathbf{T}_j$.

Reading $\mathbf{A}\mathbf{Q} = \mathbf{Q}\mathbf{T}$ one column at a time,

$$\begin{aligned}
\mathbf{A}\mathbf{q}_1 &= t_{11}\mathbf{q}_1, \\
\mathbf{A}\mathbf{q}_2 &= t_{12}\mathbf{q}_1 + t_{22}\mathbf{q}_2, \\
&\vdots \\
\mathbf{A}\mathbf{q}_j &= t_{1j}\mathbf{q}_1 + \cdots + t_{jj}\mathbf{q}_j.
\end{aligned}$$

So the first Schur vector $\mathbf{q}_1$ is an eigenvector of $\mathbf{A}$, with eigenvalue t_{11}. The later Schur vectors are **not** eigenvectors in general: $\mathbf{q}_j$ is an eigenvector only "modulo" $\mathbf{q}_1, \ldots, \mathbf{q}_{j-1}$. But the eigenvalues all sit on the diagonal of $\mathbf{T}$, and that is what we need in order to compute them.

Goal: compute a Schur decomposition (or just its first few columns), using only matrix products and orthogonalization.

1.2 Orthogonal iteration

Power iteration applied to a vector $\mathbf{x}$ converges to the direction of $\mathbf{v}_1$, which is also the first Schur vector $\mathbf{q}_1$. How could we find the *second* Schur vector as well?

Idea: run power iteration on *two* vectors $\mathbf{x}$ and $\mathbf{y}$ at once. Unfortunately, if we just compute $\mathbf{A}^k\mathbf{x}$ and $\mathbf{A}^k\mathbf{y}$, then both converge to the direction of $\mathbf{v}_1$ (generically, both have a nonzero component along $\mathbf{v}_1$). We learn nothing about $\mathbf{q}_2$, and the two vectors become nearly parallel, a poor (nearly dependent) basis for the plane they span.

Fix: what *does* converge is the two-dimensional *span* of $\mathbf{A}^k\mathbf{x}$ and $\mathbf{A}^k\mathbf{y}$. It is drawn toward $\operatorname{span}\{\mathbf{v}_1, \mathbf{v}_2\}$, the invariant subspace belonging to the two dominant eigenvalues. So at every step we replace the pair of vectors by an **orthonormal basis** for the same span. Starting from $\begin{bmatrix}\mathbf{q}_1 & \mathbf{q}_2\end{bmatrix}$ with orthonormal columns, one step is

$$\mathbf{A}\begin{bmatrix}\mathbf{q}_1 & \mathbf{q}_2\end{bmatrix} = \begin{bmatrix}\mathbf{q}_1^+ & \mathbf{q}_2^+\end{bmatrix}\begin{bmatrix} r_{11} & r_{12} \\ 0 & r_{22}\end{bmatrix} \qquad \text{(a reduced QR factorization).}$$

Reading this column by column,

$$\mathbf{q}_1^+ = \frac{\mathbf{A}\mathbf{q}_1}{r_{11}}, \qquad \mathbf{q}_2^+ = \frac{\mathbf{A}\mathbf{q}_2 - r_{12}\mathbf{q}_1^+}{r_{22}}.$$

The first vector is just a normalized power iteration. The second is $\mathbf{A}\mathbf{q}_2$ with its component along $\mathbf{q}_1^+$ removed, then normalized (Gram–Schmidt!). The two vectors cannot collapse onto each other, since $\mathbf{q}_2^+ \perp \mathbf{q}_1^+$ by construction.

Where does this lead? Suppose $\mathbf{q}_1$ converges to an eigenvector and $\operatorname{span}\{\mathbf{q}_1, \mathbf{q}_2\}$ converges to $\operatorname{span}\{\mathbf{v}_1, \mathbf{v}_2\}$. Then in the limit

$$\mathbf{A}\mathbf{q}_1 = t_{11}\mathbf{q}_1, \qquad \mathbf{A}\mathbf{q}_2 = t_{12}\mathbf{q}_1 + t_{22}\mathbf{q}_2,$$

since $\mathbf{A}\mathbf{q}_2 \in \operatorname{span}\{\mathbf{v}_1, \mathbf{v}_2\} = \operatorname{span}\{\mathbf{q}_1, \mathbf{q}_2\}$. These are exactly the first two columns of a Schur decomposition!

Notice that $\mathbf{q}_2$ is a unit vector in $\operatorname{span}\{\mathbf{v}_1, \mathbf{v}_2\}$ that is orthogonal to $\mathbf{v}_1$. In general it is *not* $\mathbf{v}_2$.

The same idea works for any number $r \leq n$ of vectors.

Algorithm: Orthogonal iteration

Given matrix $\mathbf{A}$ and a number of columns r:

1. Choose $\mathbf{Q}_0 \in \mathbb{R}^{n \times r}$ with orthonormal columns (for instance, the $\mathbf{Q}$ factor of a random matrix).
2. For $k = 1, 2, \ldots$,
   a. Set $\mathbf{Z}_k = \mathbf{A}\mathbf{Q}_{k-1}$.
   b. Find a reduced QR factorization $\mathbf{Z}_k = \mathbf{Q}_k\mathbf{R}_k$, where $\mathbf{Q}_k$ is $n \times r$ with orthonormal columns and $\mathbf{R}_k$ is $r \times r$ upper triangular.

After each iteration, the $r \times r$ matrix $\mathbf{T}_k = \mathbf{Q}_k^*\mathbf{A}\mathbf{Q}_k$ provides eigenvalue estimates on its diagonal. For $r = 1$, this is exactly the power iteration (normalizing with the 2-norm instead of the ∞-norm).

Since a product of upper triangular matrices is upper triangular, the iteration satisfies

$$\mathbf{A}^k\mathbf{Q}_0 = \mathbf{Q}_k \underbrace{\mathbf{R}_k\mathbf{R}_{k-1}\cdots\mathbf{R}_1}_{\text{upper triangular}}.$$

(By induction: $\mathbf{A}^k\mathbf{Q}_0 = \mathbf{A}\mathbf{Q}_{k-1}(\mathbf{R}_{k-1}\cdots\mathbf{R}_1) = \mathbf{Q}_k\mathbf{R}_k\mathbf{R}_{k-1}\cdots\mathbf{R}_1$.)

In other words, for each $j \leq r$ the first j columns of $\mathbf{Q}_k$ are an orthonormal basis for the range of the first j columns of $\mathbf{A}^k\mathbf{Q}_0$. **Orthogonal iteration is power iteration on subspaces**.

```
  Project No packages added to or removed from `~/Documents/GitHub/Math-189M-NumericalLinearAlgebra/
 Manifest No packages added to or removed from `~/Documents/GitHub/Math-189M-NumericalLinearAlgebra/
```

```julia
%%julia
"""
    orthiter(A,r,numiter)

Perform `numiter` orthogonal iterations with the matrix `A`, tracking `r`
columns and starting from a random matrix with orthonormal columns. Returns an
`r×numiter` matrix whose `k`th column is the diagonal of `Q_k'*A*Q_k` (the
eigenvalue estimates after `k` iterations), and the final `n×r` matrix `Q`.
"""
function orthiter(A,r,numiter)
    n = size(A,1)
    Q = Matrix(qr(randn(n,r)).Q)     # n×r matrix with orthonormal columns
    d = zeros(r,numiter)
    for k in 1:numiter
        Z = A*Q
        Q = Matrix(qr(Z).Q)          # reduced QR: keep the n×r factor Q
        d[:,k] = diag(Q'*Z)          # diagonal of Q_k'*A*Q_k
    end
    return d,Q
end
```

```
orthiter (generic function with 1 method)
```

As in the power iteration, the only use of $\mathbf{A}$ is to compute the product $\mathbf{A}\mathbf{Q}$, which easily exploits the sparsity of $\mathbf{A}$.

1.2.1 Convergence to the Schur decomposition

Punchline:

Let $\mathbf{A} = \mathbf{V}\mathbf{D}\mathbf{V}^{-1}$, and let $1 \le r \le n$. Define $\mathbf{Q}^{\mathbf{V}}$ to be the $n \times r$ matrix produced by the QR decomposition of the matrix whose columns are the eigenvectors associated to the top r eigenvalues. The matrix $\mathbf{Q}_k^\top \mathbf{A}\mathbf{Q}_k$ converges (up to some sign changes) to

$$[\mathbf{Q}^{\mathbf{V}}]^\top \mathbf{A}\mathbf{Q}^{\mathbf{V}}$$

which is upper-triangular and has the top r eigenvalues $\lambda_1, \cdots, \lambda_r$ of $\mathbf{A}$ on the diagonal.

Let's test this on a 5×5 matrix with prescribed eigenvalues, built as $\mathbf{A} = \mathbf{V}\mathbf{D}\mathbf{V}^{-1}$ from a random matrix $\mathbf{V}$ of eigenvectors. We will track $r = 3$ columns.

```
%%julia
 = [1, 0.2, 0.08, 0.048, 0.0144]
n = length( )
V = randn(n,n)
A = V*diagm( )/V          # A = V D V ⁻¹ has eigenvalues
```

```
5×5 Matrix{Float64}:
 -5.52574   1.31592    -0.775713   -3.27216   24.6888
 -3.99519   0.994133   -0.579987   -2.36198   17.5075
 -1.62492   0.408387   -0.0841951  -0.910319   7.30608
 -0.156623  0.0620753   0.124872    0.041834   0.949583
 -1.31232   0.314361   -0.157009   -0.756372   5.91637
```

```
%%julia
r = 3
numiter = 50
d,Q = orthiter(A,r,numiter);
```

After the final iteration, the diagonal of $\mathbf{T}_k = \mathbf{Q}_k^* \mathbf{A}\mathbf{Q}_k$ (left column below) matches the three largest eigenvalues (right column) to many digits:

```
%%julia
[d[:,end]  [1:r]]
```

```
3×2 Matrix{Float64}:
 1.0   1.0
 0.2   0.2
 0.08  0.08
```


The matrix $\mathbf{T}_k$ is also nearly upper triangular, and $\mathbf{Q}_k$ nearly satisfies $\mathbf{A}\mathbf{Q}_k = \mathbf{Q}_k\mathbf{T}_k$, as the theorem predicts.

```
%%julia
T = Q'*A*Q
@show norm(tril(T,-1));      # size of the part below the diagonal
@show norm(A*Q - Q*T);       # how far range(Q) is from being an invariant subspace
```

```
norm(tril(T, -1)) = 1.452694205900598e-15
norm(A * Q - Q * T) = 3.947967500216431e-13
```

Here is how the error in each eigenvalue estimate behaves as the iteration proceeds.

```
%%julia
errs = max.(abs.(d .- [1:r]), eps())      # floor at machine epsilon so zeros can appear on a log axis
plot(1:numiter,errs',m=:o,label=["lambda_1" "lambda_2" "lambda_3"],
    title="Convergence of orthogonal iteration",
    xlabel="k",yaxis=("error in estimate",:log10,[1e-16,1]))
```



All three estimates converge linearly (straight lines on the semilog plot), but at *different* rates. The first eigenvalue is found fastest, and the third slowest. Let's find out why.

1.2.2 Convergence rate

For $j < n$, the distance between range($\mathbf{Q}_k[:, 1:j]$) and span$\{\mathbf{v}_1, \dots, \mathbf{v}_j\}$ is $O(\rho_j^k)$, where

$$\rho_j = \left| \frac{\lambda_{j+1}}{\lambda_j} \right|, \qquad j = 1, \dots, n-1.$$

The jth Schur vector $\mathbf{q}_j$ is determined by *both* range($\mathbf{Q}_k[:, 1:j]$) and range($\mathbf{Q}_k[:, 1:j-1]$). We expect it, and the eigenvalue estimate $(\mathbf{T}_k)_{jj}$, to converge linearly at the rate

$$\max\{\rho_{j-1}, \rho_j\} \qquad (\text{with } \rho_0 := 0).$$

```
%%julia
  = abs.( [2:end] ./  [1:end-1])        #  _1,…, _{n-1}
@show predicted = [max(j>1 ?  [j-1] : 0.0,  [j]) for j in 1:r];
@show observed = errs[:,16] ./ errs[:,15];
```

```
predicted = [max(if j > 1
             [j - 1]
         else
             0.0
         end, [j]) for j = 1:r] = [0.2, 0.39999999999999997, 0.6]
observed = errs[:, 16] ./ errs[:, 15] = [0.19984899469617173, 0.3999851043723837, 0.5632048277059104]
```

1.3 QR iteration

What if we want *all* of the Schur decomposition? Take $r = n$ and $\mathbf{Q}_0 = \mathbf{I}$ in orthogonal iteration. Then (assuming the leading principal submatrices of $\mathbf{V}^{-1}$ are nonsingular) the punchline says that $\mathbf{Q}_k$ converges to the unitary factor of a full Schur decomposition, and

$$\mathbf{T}_k = \mathbf{Q}_k^* \mathbf{A} \mathbf{Q}_k \ \rightarrow \ \mathbf{T}, \qquad \text{diag}(\mathbf{T}) = (\lambda_1, \ldots, \lambda_n).$$

It turns out that we can organize this computation so that we never form $\mathbf{A}\mathbf{Q}_{k-1}$ at all. We update the matrix $\mathbf{T}_k$ directly.

Algorithm: QR iteration

Given matrix $\mathbf{A}$:

1. Set $\mathbf{T}_0 = \mathbf{A}$.
2. For $k = 1, 2, \ldots,$
   a. Find a QR factorization $\mathbf{T}_{k-1} = \mathbf{U}_k \mathbf{R}_k$.
   b. Set $\mathbf{T}_k = \mathbf{R}_k \mathbf{U}_k$.

We factor, then multiply the factors back together *in the reverse order*. Why does this find eigenvalues?

Theorem:

Let $\mathbf{Q}_k = \mathbf{U}_1 \mathbf{U}_2 \cdots \mathbf{U}_k$. Then
1. $\mathbf{T}_k = \mathbf{U}_k^* \mathbf{T}_{k-1} \mathbf{U}_k = \mathbf{Q}_k^* \mathbf{A} \mathbf{Q}_k$. In particular, $\mathbf{T}_k$ is similar to $\mathbf{A}$.
2. $\mathbf{A}^k = \mathbf{Q}_k \left(\mathbf{R}_k \mathbf{R}_{k-1} \cdots \mathbf{R}_1\right)$.

Proof:

For part 1, $\mathbf{U}_k^* \mathbf{T}_{k-1} \mathbf{U}_k = \mathbf{U}_k^* (\mathbf{U}_k \mathbf{R}_k) \mathbf{U}_k = \mathbf{R}_k \mathbf{U}_k = \mathbf{T}_k$, and applying this repeatedly gives $\mathbf{T}_k = \mathbf{Q}_k^* \mathbf{A} \mathbf{Q}_k$.

For part 2, we use induction on k. The case $k = 1$ is $\mathbf{A} = \mathbf{U}_1 \mathbf{R}_1$. If $\mathbf{A}^{k-1} = \mathbf{Q}_{k-1} \mathbf{R}_{k-1} \cdots \mathbf{R}_1$, then by part 1 (in the form $\mathbf{A}\mathbf{Q}_{k-1} = \mathbf{Q}_{k-1} \mathbf{T}_{k-1}$),

$$\mathbf{A}^k = \mathbf{A}\mathbf{Q}_{k-1} \mathbf{R}_{k-1} \cdots \mathbf{R}_1 = \mathbf{Q}_{k-1} \mathbf{T}_{k-1} \mathbf{R}_{k-1} \cdots \mathbf{R}_1 = \mathbf{Q}_{k-1} \mathbf{U}_k \mathbf{R}_k \mathbf{R}_{k-1} \cdots \mathbf{R}_1 = \mathbf{Q}_k \mathbf{R}_k \cdots \mathbf{R}_1.$$

Part 2 says that $\mathbf{Q}_k$ is the $\mathbf{Q}$ factor in a QR factorization of $\mathbf{A}^k$. But orthogonal iteration with $r = n$ and $\mathbf{Q}_0 = \mathbf{I}$ produces iterates $\tilde{\mathbf{Q}}_k$ with $\mathbf{A}^k = \tilde{\mathbf{Q}}_k \tilde{\mathbf{R}}_k$, from the identity we found earlier. A QR factorization of an invertible matrix is unique up to the signs of the columns of $\mathbf{Q}$ (and rows of $\mathbf{R}$), so $\mathbf{Q}_k = \tilde{\mathbf{Q}}_k$ up to signs.

The QR iteration *is* orthogonal iteration in disguise (with $r = n$ and $\mathbf{Q}_0 = \mathbf{I}$), we have that the QR iterate $\mathbf{T}_k$ is exactly the matrix $\mathbf{Q}_k^* \mathbf{A} \mathbf{Q}_k$ from orthogonal iteration.

Therefore, under the same assumptions as before,

$$\mathbf{T}_k \to \mathbf{T} \text{ (upper triangular)}, \qquad \operatorname{diag}(\mathbf{T}_k) \to (\lambda_1, \ldots, \lambda_n), \qquad \mathbf{Q}_k \to \text{Schur vectors (up to signs)}.$$

The entries below the diagonal decay linearly, with $(\mathbf{T}_k)_{ij} \to 0$ at the rate $|\lambda_i/\lambda_j|^k$ for $i > j$. The slowest entries are on the first subdiagonal, with rates $|\lambda_{j+1}/\lambda_j| = \rho_j$.

```
%%julia
"""
    qriter(A,numiter)

Perform `numiter` iterations of the QR iteration with the matrix `A`. Returns
the final iterate `T_k`, the accumulated orthogonal matrix `Q_k = U_1 U_k`,
and a vector holding the norm of the strictly lower triangular part of `T_k`
after each iteration.
"""
function qriter(A,numiter)
    n = size(A,1)
    Tk = copy(A)
    Q = Matrix(1.0I,n,n)
    below = zeros(numiter)
    for k in 1:numiter
        U,R = qr(Tk)
        U = Matrix(U)
        Tk = R*U                        # factor, then multiply in reverse order
        Q = Q*U
        below[k] = norm(tril(Tk,-1))
    end
    return Tk,Q,below
end
```

```
qriter (generic function with 1 method)
```

We use a matrix with the same known eigenvalues as before.

```
%%julia
 = [1, 0.2, 0.08, 0.048, 0.0144]
n = length( )
V = randn(n,n)
A = V*diagm( )/V          # A = V D V ⁻¹ has eigenvalues
```

```
5×5 Matrix{Float64}:
  0.192741  -1.32032   -0.792326  -0.802939  -0.0838067
 -0.619556   6.47208    3.93886    4.30935    0.406667
  0.287389  -3.17412   -1.91404   -2.20384   -0.192338
  0.557902  -5.46934   -3.33561   -3.53152   -0.359904
 -0.072867   0.513122   0.293357   0.259357   0.123142
```

After 60 iterations, $\mathbf{T}_k$ looks upper triangular.

```
%%julia
Tk,Q,below = qriter(A,60);
round.(Tk,digits=6)
```

```
5×5 Matrix{Float64}:
  1.0  -0.642321   0.422371   1.44045   -12.0834
  0.0   0.2       -0.074502  -0.192891    1.69933
  0.0   0.0        0.08       0.01018    -0.177426
  0.0   0.0        0.0        0.048      -0.032402
 -0.0  -0.0       -0.0       -0.0         0.0144
```

The diagonal entries are the eigenvalues, in order of decreasing magnitude, just as predicted:

```
%%julia
[diag(Tk)  ]
```

```
5×2 Matrix{Float64}:
 1.0     1.0
 0.2     0.2
 0.08    0.08
 0.048   0.048
 0.0144  0.0144
```


Since $\mathbf{T}_k$ is similar to $\mathbf{A}$ by the orthogonal matrix $\mathbf{Q}_k = \mathbf{U}_1 \cdots \mathbf{U}_k$, we can also check part 1 of the theorem.

```
%%julia
@show norm(Q'*A*Q - Tk);
```

```
norm(Q' * A * Q - Tk) = 4.5535196783874965e-15
```

The part of $\mathbf{T}_k$ below the diagonal decays linearly. The slowest-converging entry is $(\mathbf{T}_k)_{43}$, which has rate $|\lambda_4/\lambda_3| = \rho_3$, the largest of the ratios ρ_j.

```
%%julia
plot(1:60,below,m=:o,title="QR iteration: part of T_k below the diagonal",
    xlabel="k",yaxis=("norm of strictly lower part",:log10,[1e-16,10]),legend=false)
```




```
%%julia
@show observed = below[31]/below[30];
@show theory = maximum(abs.( [2:end] ./  [1:end-1]));
```

```
observed = below[31] / below[30] = 0.6000000557042746
theory = maximum(abs.( [2:var"end"] ./  [1:var"end" - 1])) = 0.6
```

1.3.1 Symmetric case

Suppose $\mathbf{A}$ is real and symmetric. Then in a Schur decomposition $\mathbf{A} = \mathbf{Q}\mathbf{T}\mathbf{Q}^\top$,

$$\mathbf{T}^\top = (\mathbf{Q}^\top \mathbf{A} \mathbf{Q})^\top = \mathbf{Q}^\top \mathbf{A}^\top \mathbf{Q} = \mathbf{T}.$$

So $\mathbf{T}$ is both upper triangular and symmetric, and is therefore *diagonal*.

For a symmetric matrix, **the Schur decomposition and the eigendecomposition are the same**: $\mathbf{T} = \mathbf{D}$, and the Schur vectors are orthonormal eigenvectors. (This is the Spectral Theorem.) For the QR iteration this means that

- every iterate $\mathbf{T}_k = \mathbf{Q}_k^\top \mathbf{A} \mathbf{Q}_k$ is symmetric, and $\mathbf{T}_k$ converges to a *diagonal* matrix holding the eigenvalues;
- the accumulated matrix $\mathbf{Q}_k = \mathbf{U}_1 \cdots \mathbf{U}_k$ converges (up to the signs of its columns) to a matrix of orthonormal eigenvectors.

The eigenvalues converge even faster than for a nonsymmetric matrix. The diagonal entries of a symmetric matrix differ from its eigenvalues only by an amount that is quadratic in the size of the off-diagonal part. So if the off-diagonal part of $\mathbf{T}_k$ is $O(\rho^k)$, then the diagonal entries have error $O(\rho^{2k})$.

We construct a symmetric matrix with the same eigenvalues λ, using a random orthogonal matrix of eigenvectors.

```
%%julia
Q0 = Matrix(qr(randn(n,n)).Q)       # a random orthogonal matrix
S = Q0*diagm( )*Q0'
S = (S+S')/2                         # symmetrize, to remove any roundoff asymmetry
Sk,Q,below = qriter(S,60);
round.(Sk,digits=6)
```

```
5×5 Matrix{Float64}:
  1.0  -0.0  -0.0   -0.0    -0.0
 -0.0   0.2   0.0   -0.0    -0.0
 -0.0   0.0   0.08  -0.0    -0.0
  0.0  -0.0  -0.0    0.048   0.0
 -0.0   0.0  -0.0   -0.0     0.0144
```

The iterate is diagonal, and its diagonal agrees with the eigenvalues computed by Julia. Moreover, $\mathbf{Q}_k$ contains the eigenvectors: $\mathbf{S}\mathbf{Q}_k \approx \mathbf{Q}_k\mathbf{D}$.

```
%%julia
@show diag(Sk);
@show reverse(eigvals(Symmetric(S)));       # eigvals returns the eigenvalues in ascending order
@show norm(S*Q - Q*Diagonal(diag(Sk)));
```

```
diag(Sk) = [1.0, 0.2000000000000001, 0.08000000000000007, 0.048000000000000036, 0.01439999999999991]
reverse(eigvals(Symmetric(S))) = [1.0, 0.20000000000000015, 0.07999999999999997, 0.04800000000000001, 0
norm(S * Q - Q * Diagonal(diag(Sk))) = 8.666815562496249e-16
```

1.3.2 Improvements to QR iteration

The QR iteration as we have described it has two practical drawbacks:

1. **Cost.** Each iteration requires a QR factorization and a matrix product. For a dense $n \times n$ matrix each of these costs $O(n^3)$ flops, and we need many iterations.
2. **Slow convergence.** The linear rate $|\lambda_{j+1}/\lambda_j|$ can be arbitrarily close to 1. (If $|\lambda_{j+1}| = |\lambda_j|$ for some j, the iteration does not converge to a triangular matrix at all.)

Both can be fixed, and the two fixes together are the basis of the algorithms used in practice. (For example, LAPACK's routines for the nonsymmetric eigenvalue problem, which underlie `eigen` in Julia, are built on them.)

1. Put $\mathbf{A}$ first into upper Hessenberg form through Householder reflections – QR is much cheaper for a matrix of this form.
2. Speed convergence using a shift strategy like we saw in the inverse iteration method!

These methods are the workhorses for dense matrices of moderate size. For large sparse matrices, we will want methods that use only matrix-vector products, which we will see later in this course.