

Iterative Projection Methods

for noisy and corrupted systems of linear equations

Jamie Haddock


February 1, 2018

Graduate Group in Applied Mathematics
UC Davis

joint with Jesús De Loera and Deanna Needell
<https://arxiv.org/abs/1605.01418> (SISC 2017) and forthcoming articles

Setup

We are interested in solving **highly overdetermined systems of equations**, $A\mathbf{x} = \mathbf{b}$, where $A \in \mathbb{R}^{m \times n}$, $\mathbf{b} \in \mathbb{R}^m$ and $m \gg n$. Rows are denoted \mathbf{a}_i^T .



If $\{\mathbf{x} \in \mathbb{R}^n : A\mathbf{x} = \mathbf{b}\}$ is nonempty, these methods construct an **approximation** to an element:


1. Randomized Kaczmarz Method
2. Motzkin's Method(s)
3. Sampling Kaczmarz-Motzkin Methods (SKM)

Randomized Kaczmarz Method


Given $\mathbf{x}_0 \in \mathbb{R}^n$:

1. Choose $i_k \in [m]$ with probability $\frac{\|\mathbf{a}_{i_k}\|^2}{\|A\|_F^2}$.
2. Define $\mathbf{x}_k := \mathbf{x}_{k-1} + \frac{b_{i_k} - \mathbf{a}_{i_k}^T \mathbf{x}_{k-1}}{\|\mathbf{a}_{i_k}\|^2} \mathbf{a}_{i_k}$.
3. Repeat.


Kaczmarz Method




Kaczmarz Method



Kaczmarz Method



Kaczmarz Method



Theorem (Strohmer - Vershynin 2009)

Let \mathbf{x} be the solution to the consistent system of linear equations $A\mathbf{x} = \mathbf{b}$. Then the Random Kaczmarz method converges to \mathbf{x} linearly in expectation:


$$\mathbb{E} \|\mathbf{x}_k - \mathbf{x}\|_2^2 \leq \left(1 - \frac{1}{\|A\|_F^2 \|A^{-1}\|_2^2} \right)^k \|\mathbf{x}_0 - \mathbf{x}\|_2^2.$$

Motzkin's Relaxation Method(s)


Given $\mathbf{x}_0 \in \mathbb{R}^n$:

1. If \mathbf{x}_k is feasible, stop.
2. Choose $i_k \in [m]$ as $i_k := \operatorname{argmax}_{i \in [m]} |\mathbf{a}_i^T \mathbf{x}_{k-1} - b_i|$.
3. Define $\mathbf{x}_k := \mathbf{x}_{k-1} + \frac{b_{i_k} - \mathbf{a}_{i_k}^T \mathbf{x}_{k-1}}{\|\mathbf{a}_{i_k}\|^2} \mathbf{a}_{i_k}$.
4. Repeat.


Motzkin's Method



Motzkin's Method



Motzkin's Method



Theorem (Agmon 1954)


For a consistent, normalized system, $\|\mathbf{a}_i\| = 1$ for all $i = 1, \dots, m$, Motzkin's method converges linearly to the solution \mathbf{x} :


$$\|\mathbf{x}_k - \mathbf{x}\|^2 \leq \left(1 - \frac{1}{m\|A^{-1}\|^2}\right)^k \|\mathbf{x}_0 - \mathbf{x}\|^2.$$


Our Hybrid Method (SKM)

Given $\mathbf{x}_0 \in \mathbb{R}^n$:

1. Choose $\tau_k \subset [m]$ to be a sample of size β constraints chosen uniformly at random from among the rows of A .
2. From among these β rows, choose $i_k := \operatorname{argmax}_{i \in \tau_k} |\mathbf{a}_i^T \mathbf{x}_{k-1} - b_i|$.
3. Define $\mathbf{x}_k := \mathbf{x}_{k-1} + \frac{b_{i_k} - \mathbf{a}_{i_k}^T \mathbf{x}_{k-1}}{\|\mathbf{a}_{i_k}\|^2} \mathbf{a}_{i_k}$.
4. Repeat.








Theorem (De Loera - H. - Needell 2017)

For a consistent, normalized system the SKM method with samples of size β converges to the solution \mathbf{x} at least linearly in expectation: If s_{k-1} is the number of constraints satisfied by \mathbf{x}_{k-1} and

$V_{k-1} = \max\{m - s_{k-1}, m - \beta + 1\}$ then

$$\begin{aligned}\mathbb{E}\|\mathbf{x}_k - \mathbf{x}\|^2 &\leq \left(1 - \frac{1}{V_{k-1}\|A^{-1}\|^2}\right)\|\mathbf{x}_0 - \mathbf{x}\|^2 \\ &\leq \left(1 - \frac{1}{m\|A^{-1}\|^2}\right)^k \|\mathbf{x}_0 - \mathbf{x}\|^2.\end{aligned}$$

Convergence



$$\triangleright \text{RK: } \mathbb{E} \|\mathbf{x}_k - \mathbf{x}\|_2^2 \leq \left(1 - \frac{1}{\|A\|_F^2 \|A^{-1}\|_2^2} \right)^k \|\mathbf{x}_0 - \mathbf{x}\|_2^2.$$

Convergence Rates

$$\triangleright \text{RK: } \mathbb{E} \|\mathbf{x}_k - \mathbf{x}\|_2^2 \leq \left(1 - \frac{1}{\|A\|_F^2 \|A^{-1}\|_2^2} \right)^k \|\mathbf{x}_0 - \mathbf{x}\|_2^2.$$

$$\triangleright \text{MM: } \|\mathbf{x}_k - \mathbf{x}\|^2 \leq \left(1 - \frac{1}{m \|A^{-1}\|^2} \right)^k \|\mathbf{x}_0 - \mathbf{x}\|^2.$$

Convergence Rates

$$\triangleright \text{RK: } \mathbb{E} \|\mathbf{x}_k - \mathbf{x}\|_2^2 \leq \left(1 - \frac{1}{\|A\|_F^2 \|A^{-1}\|_2^2}\right)^k \|\mathbf{x}_0 - \mathbf{x}\|_2^2.$$

$$\triangleright \text{MM: } \|\mathbf{x}_k - \mathbf{x}\|^2 \leq \left(1 - \frac{1}{m \|A^{-1}\|^2}\right)^k \|\mathbf{x}_0 - \mathbf{x}\|^2.$$

$$\triangleright \text{SKM: } \mathbb{E} \|\mathbf{x}_k - \mathbf{x}\|^2 \leq \left(1 - \frac{1}{m \|A^{-1}\|^2}\right)^k \|\mathbf{x}_0 - \mathbf{x}\|^2.$$

Convergence Rates

▷ RK: $\mathbb{E} \|\mathbf{x}_k - \mathbf{x}\|_2^2 \leq \left(1 - \frac{1}{\|A\|_F^2 \|A^{-1}\|_2^2}\right)^k \|\mathbf{x}_0 - \mathbf{x}\|_2^2.$

▷ MM: $\|\mathbf{x}_k - \mathbf{x}\|^2 \leq \left(1 - \frac{1}{m \|A^{-1}\|^2}\right)^k \|\mathbf{x}_0 - \mathbf{x}\|^2.$

▷ SKM: $\mathbb{E} \|\mathbf{x}_k - \mathbf{x}\|^2 \leq \left(1 - \frac{1}{m \|A^{-1}\|^2}\right)^k \|\mathbf{x}_0 - \mathbf{x}\|^2.$

▷ Why are these all the same?

An Accelerated Convergence Rate

Theorem (H. - Needell 2018+)



Let \mathbf{x} denote the solution of the consistent, normalized system $A\mathbf{x} = \mathbf{b}$.
Motskin's method exhibits the (possibly highly accelerated) convergence rate:

$$\|\mathbf{x}_T - \mathbf{x}\|^2 \leq \prod_{k=0}^{T-1} \left(1 - \frac{1}{4\gamma_k \|A^{-1}\|^2} \right) \cdot \|\mathbf{x}_0 - \mathbf{x}\|^2$$



Here γ_k bounds the dynamic range of the k th residual, $\gamma_k := \frac{\|A\mathbf{x}_k - A\mathbf{x}\|^2}{\|A\mathbf{x}_k - A\mathbf{x}\|_\infty^2}$.

▷ improvement over previous result when $4\gamma_k < m$

γ_k : Gaussian systems




γ_k : Gaussian systems




$$\gamma_k \lesssim \frac{m}{\log m}$$

Gaussian Convergence




Is this the right problem?



▷ noisy

Is this the right problem?



▷ noisy

▷ corrupted



Noisy Convergence Results

Theorem (Needell 2010)

Let A have full column rank, denote the desired solution to the system $A\mathbf{x} = \mathbf{b}$ by \mathbf{x} , and define the error term $\mathbf{e} = A\mathbf{x} - \mathbf{b}$. Then RK iterates satisfy

$$\mathbb{E}\|\mathbf{x}_k - \mathbf{x}\|^2 \leq \left(1 - \frac{1}{\|A\|_F^2 \|A^{-1}\|^2}\right)^k \|\mathbf{x}_0 - \mathbf{x}\|^2 + \|A\|_F^2 \|A^{-1}\|^2 \|\mathbf{e}\|_\infty^2$$

Noisy Convergence Results

Theorem (Needell 2010)

Let A have full column rank, denote the desired solution to the system $A\mathbf{x} = \mathbf{b}$ by \mathbf{x} , and define the error term $\mathbf{e} = A\mathbf{x} - \mathbf{b}$. Then RK iterates satisfy



$$\mathbb{E}\|\mathbf{x}_k - \mathbf{x}\|^2 \leq \left(1 - \frac{1}{\|A\|_F^2 \|A^{-1}\|^2}\right)^k \|\mathbf{x}_0 - \mathbf{x}\|^2 + \|A\|_F^2 \|A^{-1}\|^2 \|\mathbf{e}\|_\infty^2$$

Theorem (H. - Needell 2018+)


Let \mathbf{x} denote the desired solution of the system $A\mathbf{x} = \mathbf{b}$ and define the error term $\mathbf{e} = \mathbf{b} - A\mathbf{x}$. If Motzkin's method is run with stopping criterion $\|A\mathbf{x}_k - \mathbf{b}\|_\infty \leq 4\|\mathbf{e}\|_\infty$, then the iterates satisfy

$$\|\mathbf{x}_T - \mathbf{x}\|^2 \leq \prod_{k=0}^{T-1} \left(1 - \frac{1}{4\gamma_k \|A^{-1}\|^2}\right) \cdot \|\mathbf{x}_0 - \mathbf{x}\|^2 + 2m \|A^{-1}\|^2 \|\mathbf{e}\|_\infty^2$$

Noisy Convergence



What about corruption?



Problem

	Problem:	$A\mathbf{x} = \mathbf{b} + \mathbf{e}$
(Corrupted)	Error (e):	sparse, arbitrarily large entries
	Solution (\mathbf{x}^*):	$\mathbf{x}^* \in \{\mathbf{x} : A\mathbf{x} = \mathbf{b}\}$

Problem

	Problem:	$A\mathbf{x} = \mathbf{b} + \mathbf{e}$
(Corrupted)	Error (e):	sparse, arbitrarily large entries
	Solution (\mathbf{x}^*):	$\mathbf{x}^* \in \{\mathbf{x} : A\mathbf{x} = \mathbf{b}\}$

Applications: logic programming, error correction in telecommunications


Problem

	Problem:	$A\mathbf{x} = \mathbf{b} + \mathbf{e}$
(Corrupted)	Error (e):	sparse, arbitrarily large entries
	Solution (\mathbf{x}^*):	$\mathbf{x}^* \in \{\mathbf{x} : A\mathbf{x} = \mathbf{b}\}$

Applications: logic programming, error correction in telecommunications

	Problem:	$A\mathbf{x} = \mathbf{b} + \mathbf{e}$
(Noisy)	Error (e):	small, evenly distributed entries
	Solution (\mathbf{x}_{LS}):	$\mathbf{x}_{LS} \in \operatorname{argmin} \ A\mathbf{x} - \mathbf{b} - \mathbf{e}\ ^2$

Why not least-squares?



MAX-FS: Given $A\mathbf{x} = \mathbf{b}$, determine the largest feasible subsystem.

MAX-FS: Given $A\mathbf{x} = \mathbf{b}$, determine the largest feasible subsystem.

- ▷ MAX-FS is NP-hard even when restricted to homogenous systems with coefficients in $\{-1, 0, 1\}$ (Amaldi - Kann 1995)

MAX-FS: Given $A\mathbf{x} = \mathbf{b}$, determine the largest feasible subsystem.

- ▷ MAX-FS is NP-hard even when restricted to homogenous systems with coefficients in $\{-1, 0, 1\}$ (Amaldi - Kann 1995)
- ▷ no PTAS unless $P = NP$

Goal: Use RK to detect the corrupted equations with high probability.

Goal: Use RK to detect the corrupted equations with high probability.


Lemma

Let $\epsilon^ = \min_{i \in \text{supp}(\mathbf{e})} |\mathbf{A}\mathbf{x}^* - \mathbf{b}|_i = |e_i|$ and suppose $|\text{supp}(\mathbf{e})| = s$. If $\|\mathbf{a}_i\| = 1$ for $i \in [m]$ and $\|\mathbf{x} - \mathbf{x}^*\| < \frac{1}{2}\epsilon^*$ we have that the $d \leq s$ indices of largest magnitude residual entries are contained in $\text{supp}(\mathbf{e})$. That is, we have $D \subset \text{supp}(\mathbf{e})$, where*

$$D = \underset{D \subset [A], |D|=d}{\operatorname{argmax}} \sum_{i \in D} |\mathbf{A}\mathbf{x} - \mathbf{b}|_i.$$


Proposed Method

Goal: Use RK to detect the corrupted equations with high probability.



Proposed Method

Goal: Use RK to detect the corrupted equations with high probability.




We call $\epsilon^*/2$ the *detection horizon*.

Method 1 Windowed Kaczmarz

- 1: **procedure** WK(A, \mathbf{b}, k, W, d)
 - 2: $S = \emptyset$
 - 3: **for** $i = 1, 2, \dots, W$ **do**
 - 4: $\mathbf{x}_k^i = k$ th iterate produced by RK with $\mathbf{x}_0 = \mathbf{0}$, A , \mathbf{b} .
 - 5: $D = d$ indices of the largest entries of the residual, $|A\mathbf{x}_k^i - \mathbf{b}|$.
 - 6: $S = S \cup D$
 - 7: **return** \mathbf{x} , where $A_{Sc}\mathbf{x} = \mathbf{b}_{Sc}$
-


Example

$\text{WK}(A, \mathbf{b}, k = 2, W = 3, d = 1)$: $j = 1, i = 1, S = \emptyset$




Example

$\text{WK}(A, \mathbf{b}, k = 2, W = 3, d = 1)$: $j = 1, i = 1, S = \emptyset$




Example

$\text{WK}(A, \mathbf{b}, k = 2, W = 3, d = 1): j = 2, i = 1, S = \{7\}$




Example

$\text{WK}(A, \mathbf{b}, k = 2, W = 3, d = 1)$: $j = 1, i = 2, S = \{7\}$




Example

$\text{WK}(A, \mathbf{b}, k = 2, W = 3, d = 1): j = 1, i = 2, S = \{7\}$




Example

$\text{WK}(A, \mathbf{b}, k = 2, W = 3, d = 1): j = 2, i = 2, S = \{7, 5\}$




Example

$\text{WK}(A, \mathbf{b}, k = 2, W = 3, d = 1): j = 1, i = 3, S = \{7, 5\}$




Example

$\text{WK}(A, b, k = 2, W = 3, d = 1)$: $j = 1, i = 3, S = \{7, 5\}$




Example

$\text{WK}(A, \mathbf{b}, k = 2, W = 3, d = 1)$: $j = 2$, $i = 3$, $S = \{7, 5, 6\}$



Example

Solve $A_{5c}\mathbf{x} = \mathbf{b}_{5c}$.



Lemma

Let $\epsilon^* = \min_{i \in \text{supp}(\mathbf{e})} \|\mathbf{A}\mathbf{x}^* - \mathbf{b}\|_i = |e_i|$ and suppose $|\text{supp}(\mathbf{e})| = s$. Assume that $\|\mathbf{a}_i\| = 1$ for all $i \in [m]$ and let $0 < \delta < 1$. Define

$$k^* = \left\lceil \frac{\log \left(\frac{\delta(\epsilon^*)^2}{4\|\mathbf{x}^*\|^2} \right)}{\log \left(1 - \frac{\sigma_{\min}^2(\mathbf{A}_{\text{supp}(\mathbf{e})^C})}{m-s} \right)} \right\rceil.$$

Then in window i of the Windowed Kaczmarz method, the iterate produced by the RK iterations, $\mathbf{x}_{k^*}^i$ satisfies

$$\mathbb{P} \left[\|\mathbf{x}_{k^*}^i - \mathbf{x}^*\| \leq \frac{1}{2}\epsilon^* \right] \geq p := (1 - \delta) \left(\frac{m-s}{m} \right)^{k^*}.$$


Theorem (H. - Needell 2018+)

Assume that $\|\mathbf{a}_i\| = 1$ for all $i \in [m]$ and let $0 < \delta < 1$. Suppose $d \geq s = |\text{supp}(\mathbf{e})|$, $W \leq \lfloor \frac{m-n}{d} \rfloor$ and k^* is as given in the previous lemma. Then the Windowed Kaczmarz method on A, \mathbf{b} will detect the corrupted equations ($\text{supp}(\mathbf{e}) \subset S$) and the remaining equations given by $A_{[m]-S}, \mathbf{b}_{[m]-S}$ will have solution \mathbf{x}^* with probability at least


$$p_W := 1 - \left[1 - (1 - \delta) \left(\frac{m-s}{m} \right)^{k^*} \right]^W.$$

Theoretical Guarantee Values (Gaussian $A \in \mathbb{R}^{50000 \times 100}$)


$$p_W := 1 - \left[1 - (1 - \delta) \left(\frac{m-s}{m} \right)^{k^*} \right]^W$$




Experimental Values (Gaussian $A \in \mathbb{R}^{50000 \times 100}$)




Experimental Values (Gaussian $A \in \mathbb{R}^{50000 \times 100}$)



Experimental Values (Gaussian $A \in \mathbb{R}^{50000 \times 100}$)



Experimental Values (Gaussian $A \in \mathbb{R}^{50000 \times 100}$)



Conclusions and Future Work

- Motzkin's method is accelerated even in the presence of noise
- RK methods may be used to detect corruption
- identify useful bounds on γ_k for other useful systems
- reduce dependence on artificial parameters in corruption detection bounds