

# 1    **Single-cell sequencing reveals unexpected genetic diversity among *Bodo* spp. flagellates** 2    **and their bacterial endosymbionts**

3    Sally D. Warring<sup>1</sup>, Jamie McGowan<sup>1</sup>, Estelle S. Kiliass<sup>2</sup>, James Lipscombe<sup>1</sup>, Elisabet Alacid<sup>2</sup>,  
4    Tom Barker<sup>1</sup>, Leah Catchpole<sup>1</sup>, Karim Gharbi<sup>1</sup>, Seanna McTaggart<sup>1</sup>, Thomas A. Richards<sup>2</sup>,  
5    David Swarbreck<sup>1</sup>, Neil Hall<sup>1,3</sup>

6

7    <sup>1</sup> *Earlham Institute, Norwich Research Park, Norwich, United Kingdom*, <sup>2</sup> *Department of*  
8    *Biology, University of Oxford, Oxford, United Kingdom*, <sup>3</sup> *School of Biological Sciences,*  
9    *University of East Anglia, Norwich, United Kingdom*

10

## 11    **Abstract**

12            *Bodo* is a cosmopolitan genus of free living bacterivorous single-celled flagellates in  
13    the class Kinetoplastea. Members of genus *Bodo* are considered the closest free-living  
14    relatives to the parasitic lineages *Trypanosoma* and *Leishmania*, the causative agents of the  
15    human diseases sleeping sickness, Chagas disease, and leishmaniasis. Currently, a single  
16    genome exists for the one formally described species in the genus, *Bodo saltans*. Previous  
17    studies on *B. saltans* have shown that it is dependent on an endosymbiotic bacterium from  
18    the order Holosporales, “*Candidatus* Bodocaeidibacter vickermanii”. Using single cell-  
19    sequencing, we isolated, sequenced, and assembled genomes for seven uncultured *Bodo*  
20    spp. cells from a single freshwater sample from Royal Leamington Spa, UK. By using  
21    comparative genomics, we show that these seven cells represent three potentially novel  
22    *Bodo* species and exhibit unexpected levels of diversity at the genome level. Our results  
23    indicate that Small Subunit (SSU) rDNA sequencing, often used to classify *Bodo* flagellates, is

insufficient for determining species delimitation in this genus. In addition, we recovered a  
 Holosporales bacterium genome from all seven *Bodo* spp. cells. Surprisingly, these seven  
 bacterial endosymbionts also represent three potentially novel species and one novel genus  
 of Holosporales bacteria. This diversity would be indistinguishable in routinely-used SSU  
 ribosomal DNA (rDNA) metabarcoding or bulk sequencing pipelines, thus demonstrating the  
 utility of using single-cell sequencing to reveal the level of genomic diversity within lineages  
 of microbial eukaryotes and their cobionts.

31

## 32 **Keywords**

Single-cell sequencing, Kinetoplastea, *Bodo saltans*, protist, Holosporales, environmental  
 sequencing

35

## 36 **Background**

*Bodo* is a genus of heterotrophic free-living bi-flagellated protists common in fresh  
 and brackish water and soil. They are members of the Kinetoplastea, a class of parasitic and  
 free-living protists distinguished by the presence of a large mass of mitochondrial DNA  
 known as kinetoplast DNA, or kDNA (1). Class Kinetoplastea is divided into two subclasses,  
 the Prokinetoplastina and the Metakinetoplastina, the latter containing four orders,  
 Eubodonida, Parabodonida, Neobodonida and the Trypanosomatida (2,3). Order  
 Trypanosomatida includes the parasites responsible for human diseases such as sleeping  
 sickness, Chagas' disease, and leishmaniasis. Phylogenies based on Small Subunit (SSU)  
 ribosomal DNA (rDNA) and protein sequences place order Eubodonida, and its one genus,

46 *Bodo*, as the sister clade to the Trypanosomatida, and *Bodo* is considered the closest free-  
47 living lineage to the parasitic Trypanosomatida lineages (4–6).

48 Traditionally, *Bodo* species were identified and distinguished by morphology, e.g. cell  
49 size and shape, length of the flagella, position of the nucleus and kinetoplast, and  
50 ultrastructural features visible by light and electron microscopy (7,8). Later, phylogenies  
51 based on molecular data showed the genus *Bodo* was paraphyletic and its members were  
52 distributed into three of the aforementioned four orders, with only one clade containing  
53 three species, *B. saltans*, *B. edax*, and *B. uncinatus* remaining genus *Bodo* (2,4,9). Further, it  
54 is suggested that *B. edax* and *B. uncinatus* are not true species and are likely isolates of *B.*  
55 *saltans* (2,10). Despite the important phylogenic position of genus *Bodo*, *B. saltans* remains  
56 the only formally described species in the genus, and a single genome from bulk culture  
57 currently exists for *B. saltans* strain Lake Konstanz (11).

58 Holosporales are alphaproteobacterial that are widespread obligate endosymbionts  
59 of eukaryotes, particularly protists (12,13). Holosporales form complex associations with  
60 their eukaryotic hosts, including infectious parasitic species (14), some that confer  
61 competitive advantages to their hosts (15), and some that may increase host fitness under  
62 certain conditions (16,17). Recent work has shown that *B. saltans* Lake Konstanz harbours  
63 an endosymbiotic bacterium from the order Holosporales, “*Candidatus* (Ca.)  
64 *Bodocaedibacter vickermanii*” (18). *B. saltans* appears dependent on its endosymbiont as  
65 antibiotic treatment results in rapid cell death, and it is hypothesized that this dependence  
66 is due to three putative addictive toxin/antitoxin systems encoded in the endosymbionts  
67 genome (18). These systems render the endosymbiont essential to its host as they encode a

68 long-lived toxin molecule alongside an antitoxin with a shorter half-life, and loss of the  
69 antitoxin results in the activation of the toxin and host cell death (19).

70 Here, we use single-cell sequencing to assemble genomes from seven uncultured  
71 cells from one environmental sample that were identified as belonging to genus *Bodo*.  
72 Genomic comparisons show that the seven genomes form three clades, potentially  
73 representing three novel *Bodo* species that diverge significantly from *B. saltans* and each  
74 other. In addition, we recover a Holosporales bacterial genome from each raw assembly and  
75 show that these putative endosymbionts also represent three potentially novel species,  
76 forming three clades that appear congruent with the phylogeny of the hosts. These results  
77 highlight the utility of using single-cell sequencing and comparative genomics to reveal the  
78 diversity within and between uncultured populations of microbial eukaryotes and their  
79 cobionts.

80

## 81 **Results**

### 82 **Single-cell sequencing and genome assembly**

83 Seven *Bodo* spp. cells were identified on two 96-well plates containing single cells  
84 isolated by Fluorescence-Activated Cell Sorting (FACS) from an environmental sample. After  
85 DNA amplification and short-read sequencing, between 20,738,413 – 48,141,972 paired-end  
86 reads were generated for each cell (Table 1). The cells were named after the well in which  
87 they were deposited and are hereafter referred to as A8, A10, B2, B7, F10, G10 and H10.  
88 Each raw assembly was curated by binning and taxonomic assignment to remove scaffolds  
89 of bacterial and non-target origin, and the curated eukaryotic assemblies range in size from

29,029,105 base pairs (bp) to 36,294,242 bp compared to the bulk-culture assembly for *B. saltans* lake Konstanz at 39,862,120 bp (Table 1). All assemblies are fragmented, each in more than 2,619 contigs with contig N50s ranging from 22,220 bp to 31,876 bp (Table 1). All assemblies range from 54.84 % to 56.02 % GC, a higher % GC than *B. saltans* lake Konstanz at 51.79 % (Table 1). Protein annotations of each genome resulted in 11,121-13,166 protein annotations per single cell genome, 30-40 % less than *B. saltans* lake Konstanz at 18,190 proteins (Table 1). Completeness analysis of the protein sets using BUSCO v5 with the Euglenozoa odb10 database gave 72.3 %-80.8 % complete BUSCOs per assembly, compared to *B. saltans* lake Konstanz at 88.5 % complete BUSCOs (Table 1).

99

| Cell Identifier   | raw paired-end reads | Assembly statistics |             |        |       | No. proteins annotated | BUSCO - Euglenozoa odb10 |     |   |    |    |
|-------------------|----------------------|---------------------|-------------|--------|-------|------------------------|--------------------------|-----|---|----|----|
|                   |                      | Length              | No. contigs | N50    | GC %  |                        | Complete %               | S   | D | F  | M  |
| A8                | 20,814,814           | 30,867,202          | 2,619       | 26,848 | 54.94 | 11,505                 | 72.3                     | 72  | 0 | 5  | 22 |
| A10               | 25,608,225           | 31,501,931          | 2,828       | 25,310 | 54.9  | 11,858                 | 76.2                     | 75  | 1 | 3  | 21 |
| B2                | 20,738,413           | 36,294,242          | 3,436       | 26,706 | 55.02 | 13,166                 | 80.8                     | 80  | 1 | 6  | 13 |
| B7                | 48,141,972           | 30,770,982          | 3,056       | 22,521 | 56.02 | 11,736                 | 77.0                     | 76  | 1 | 4  | 19 |
| F10               | 21,853,221           | 29,029,105          | 2,665       | 24,738 | 56.02 | 11,121                 | 75.4                     | 75  | 0 | 6  | 18 |
| G10               | 33,945,903           | 33,462,613          | 3,452       | 22,220 | 55    | 12,168                 | 74.6                     | 73  | 2 | 5  | 20 |
| H10               | 21,728,838           | 34,908,636          | 3,605       | 22,252 | 54.97 | 12,867                 | 80.0                     | 79  | 2 | 5  | 15 |
|                   |                      |                     |             |        |       |                        |                          |     |   |    |    |
| <i>B. saltans</i> |                      | 39,862,120          | 2,256       | 31,876 | 51.79 | 18,190                 | 88.5                     | 112 | 3 | 12 | 3  |

100

**Table 1 - *Bodo* spp. sequencing and assembly statistics.** Number of paired-end reads generated for each single cell assembly, and general assembly and annotation statistics. The assembly statistics for *B. saltans* lake Konstanz genome are shown as a comparison. BUSCO completeness for protein sets, S = complete single copy, D = complete duplicated, F = fragmented, M = missing. Euglenozoa odb10 includes 130 BUSCOs.

## The seven *Bodo* cells form three clades

BLASTn searches against the GeneBank Nucleotide (nt) database using the SSU rDNA sequences from each assembly returned three *Bodo* spp. as top hits (Table 2). For A8 and A10, the top hit is to ATCC isolate *Bodo edax* (ATCC30903), originally isolated from the Czech Republic (5). For B7 and F10, the best hit is to *B. saltans* strain HFCC309 isolated from a eutrophic pond in Germany (20). For B2, G10 and H10, the top hit is to a *B. saltans* isolate from soil in Malaysia (10). In all cases, the query sequences are more than 99% identical to their top hits. Figure 1A shows the pairwise identities of the SSU rDNA nucleotide sequences from the seven single cell assemblies. A8 and A10 are 100% identical. B7 and F10 are 99.82% identical, with one mismatch between the two sequences, a one nucleotide indel of a G residue in a string of G residues, however, due to the fragmented nature of the assemblies the SSU rDNA sequence from F10 is truncated, limiting full comparison of these sequences (Table 2 and Additional File 1). B2 and H10 are 100% identical, with G10 99.92 % identical to B2 and H10 with a one nucleotide polymorphism, a C to T transition in G10. All three SSU rDNA sequences from B2, H10 and G10 are truncated, missing ~700 nucleotides from their 5' ends (Table 2 and Additional File 1). A8/A10 differ from B2/G10/H10 by one nucleotide, an A/G transition, while B7/F10 differ more substantially from all the others, with pairwise identities ~98% (Figure 1A).

A Maximum-Likelihood (ML) tree of SSU rDNA sequences of *Bodo* and related species available in GeneBank places our seven cells onto two clades, all within the *B. saltans* lineage (Figure 1B). A8/A10 are sister to B2/G10/H10 and are part of a radiation that includes *B. edax*, *B. uncinatus* and three other *B. saltans* isolates. B7/F10 are sister to this clade, along with two other *B. saltans* isolates. *B. saltans* lake Konstanz branches with

other *B. saltans* isolates on a separate clade that diverges closer to the root of the tree (Figure 1B). The seven single cell *Bodo* genomes have SSU rDNA pairwise identities ranging between 94.7 and 95.7 % with *B. saltans* lake Konstanz and appear closer to the SSU rDNA sequence from *B. edax*, with pairwise identities ranging between 98.0 and 99.6 % identity (Figure 1B).

| Assembly | SSU length | Top hit Blastn subject accession | Top hit Blastn subject organism | Top hit Blastn query coverage | Top hit Blastn pairwise identity | Top hit Blastn bitscore |
|----------|------------|----------------------------------|---------------------------------|-------------------------------|----------------------------------|-------------------------|
| A8       | 2,147      | AY028451                         | <i>Bodo edax</i>                | 99.81%                        | 99.60%                           | 3829.84                 |
| A10      | 2,147      | AY028451                         | <i>Bodo edax</i>                | 99.81%                        | 99.60%                           | 3829.84                 |
| B7       | 2,147      | DQ207572                         | <i>Bodo saltans</i>             | 98.04%                        | 99.30%                           | 3724.34                 |
| F10      | 1,124      | DQ207572                         | <i>Bodo saltans</i>             | 97.86%                        | 99.40%                           | 1951.63                 |
| B2       | 1,393      | AY490226                         | <i>Bodo saltans</i>             | 99.86%                        | 99.70%                           | 2492.64                 |
| G10      | 1,263      | AY490226                         | <i>Bodo saltans</i>             | 100%                          | 99.80%                           | 2266.32                 |
| H10      | 1,409      | AY490226                         | <i>Bodo saltans</i>             | 99.86%                        | 99.80%                           | 2521.49                 |

**Table 2 – BLAST results for SSU rDNA sequences.** Top hits resulting from a BLASTn search of the nt database for the SSU rDNA sequence recovered from each single cell assembly.

### Extensive genomic diversity among the *Bodo* spp.

We next built a Maximum Likelihood (ML) tree from 488 single-copy orthogroups shared by *B. saltans* lake Konstanz and our seven single cell *Bodo* assemblies, using *Perkinsella* sp. as an outgroup (Figure 2A). Here, the seven single-cell genomes split into three clades with A8/A10 forming a sister clade to B2/G10/H10 and B7/F10 forming another clade branching closer to the root of the tree. *B. saltans* lake Konstanz forms its own clade closest to the root of the tree. Next, we calculated average pairwise amino acid identities (AAI) for all eight *Bodo* assemblies, with each pairwise comparison averaging AAI over a

minimum of 7,419 and a maximum of 10,181 reciprocal best hits (Additional File 2). The AAI within each clade is high (>97%) but varies substantially between the clades (Figure 2B). *B. saltans* lake Konstanz is equally distant from all seven single cell assemblies, with pairwise AAI between 51.3-51.7 % in all comparisons. B7/F10 are equally distant from A8/A10 and B7/G10/H10 with pairwise AAI between 62.1-62.6%. A8/A10 and B7/G10/H10 appear closer, with pairwise AAI between 84.8-85.0%.

We also compared the Average Nucleotide Identity (ANI) and the Aligned Fraction (AF) of each *Bodo* spp. genome using the tool skani (21) (Figure 2C). Here, a result of 0.00 for ANI and/or AF suggests that sequence similarity at the nucleotide level is too low for pairwise comparison using this method (21), e.g. *B. saltans* lake Konstanz is too distant to any of the single-cell *Bodo* genomes to compare using this metric, with values of 0.00 in every pairwise comparison. However, ANI and AF are relatively high within each of the three single-cell *Bodo* clades (>98 % ANI and > 80 % AF). Between the three clades B7/F10 is too distant from the others to compare, while A8/A10 and B2/G10/H10 show an ANI of ~84%, but an AF of ~14-16% indicating that only a small portion of the genome is close enough for comparison using ANI. The AF values shown in Figure 2C are the average of two AF values calculated for each pairwise comparison. A matrix with both AF values calculated for each pairwise comparison is in Additional File 3.

Next, we compared how many orthogroups are shared among the single-cell assemblies and *B. saltans* lake Konstanz (Figure 2D). The majority of orthogroups, 55.5% are shared by all eight assemblies. The two clades that branch most closely in the phylogenetic trees, A8/A10 and B2/G10/H10 share the highest percentage of orthogroups, 83.4%. B7/F10 share 66.1% with A8/A10 and 68% with B2/G10/H10. *B. saltans* lake Konstanz shares 62.8 %



with B7/F10. 61.4 % with A8/A10 and 63.7% with B7/G10/H10. Each clade has a small percentage of orthogroups that are unique, with *B. saltans* lake Konstanz having the most unique orthogroups.

To compare the functional complement of each genome, we mapped each protein set to the Pfam database. In each case, less than 50% of proteins in each annotation contain one or more Pfam domains (Figure 2E). We tabulated the Pfams found in each protein annotation and ran a Principal Components Analysis (PCA) on that table. The PCA separates the genomes into three clusters congruent with the phylogeny. Principle Component (PC) 1 separates *B. saltans* lake Konstanz from all others. While PC 2 separates B2 and F10 from A8/A10 and B7/G10/H10, which cluster together (Figure 2F, Additional File 4).

Taken together, the phylogeny, AAI, ANI, AF and Pfam clustering indicate that the seven single cell *Bodo* assemblies form three clades: clade A8/A10, clade B7/F10 and clade B2/G10/H10. All three clades are potentially novel *Bodo* species, and all appear equally distant from *B. saltans* lake Konstanz.

### **All *Bodo* spp. harbour Holosporales bacterial endosymbionts**

Previous studies show that *B. saltans* lake Konstanz is dependent on an endosymbiotic alphaproteobacterium from the order Holosporales, “*Ca. B. vickermanii*” (18). We went back to our binned raw genome assemblies and were able to identify a single bin in five of the genomes, 2 bins in B2 and 4 bins in B7 that were > 80 % complete according to CheckM (Additional File 5). CheckM analysis of these genomes showed that all were at least 89 % complete, and all but one had contamination levels < 5% (Table 3).

| Assembly                      |             |             |        |       | CheckM         |               |                      | Annotation |           |          |
|-------------------------------|-------------|-------------|--------|-------|----------------|---------------|----------------------|------------|-----------|----------|
| Cell                          | Length (bp) | No. contigs | N50    | GC %  | Completeness % | Contamination | strain heterogeneity | no. genes  | no. tRNAs | rDNAs    |
| A8                            | 1,591,871   | 98          | 28,883 | 37.82 | 90.32          | 0             | 0                    | 1,591      | 39 (20)   | 23s, 16s |
| A10                           | 1,758,263   | 139         | 26,374 | 37.27 | 90.32          | 0             | 0                    | 1,591      | 41 (19)   | 0        |
| B2                            | 2,174,152   | 145         | 31,250 | 34.85 | 93.55          | 26.34         | 10.34                | 2,011      | 55 (19)   | 0        |
| G10                           | 1,604,499   | 119         | 27,931 | 38.14 | 89.01          | 2.2           | 100                  | 1,363      | 39 (20)   | 23s, 16s |
| H10                           | 1,693,415   | 108         | 27,858 | 37.65 | 94.62          | 0             | 0                    | 1,463      | 39 (17)   | 0        |
| F10                           | 1,279,012   | 83          | 33,152 | 41.61 | 94.62          | 0             | 0                    | 1,234      | 35 (18)   | 0        |
| B7                            | 1,292,737   | 53          | 55,440 | 41.89 | 95.6           | 0             | 0                    | 1,220      | 38 (19)   | 0        |
|                               |             |             |        |       |                |               |                      |            |           |          |
| " <i>Ca. B. vickermanii</i> " | 1,391,311   | 1           | NA     | 40.61 | 94.62          | 0             | 0                    | 1,214      |           |          |

**Table 3 – Assembly statistics, CheckM results for Holosporales genomes.** General assembly and annotation statistics for each Holosporales genomes. No. tRNAs shows the total number of tRNAs, and the number of different amino acids in brackets. Values from "*Ca. B. vickermanii*" are shown where appropriate for comparison.

GTDB-TK classified one bin from B7 and the bin in F10 in the order Holosporales, family "*Ca. Paracaedibacteraceae*", genus UBA6184, as the same genus as "*Ca. B. vickermanii*" (Additional File 5). Six other bin, including a single bin from B2 were classified in order *Paracaedibacterales*, family UBA11393, genus JAGOTX01, which currently contains a single bacterial metagenome assembled genome (MAG) isolated from wastewater (22) (Additional File 5). The additional bins from B2 and B7 were not classified as alphaproteobacterial and were not analysed further (Additional File 5). Order *Paracaedibacterales* is considered a heterotypic synonym of order Holosporales (12). Recently it has been proposed that due to phylogenetic nesting of order Holosporales, it be down-ranked to suborder *Holosporineae*, which would include the families *Holosporaceae*,

“*Caedimonidacea*” and “*Ca. Paracaedibacteracea*” (13,23). Under this ranking, all our genomes would fall within suborder *Holosporineae*, two falling in family “*Ca. Paracaedibacteracea*” and the rest as unclassified *Holosporineae*.

To better place the bodo Holosporales, we built a ML tree from 24 single-copy orthogroups shared among Holosporales genomes and MAGs available in GeneBank and using two alphaproteobacterial as an outgroup (Figure 3A). This tree places the seven genomes into two lineages. F10/B7 branch on a clade with “*Ca. B. vickermanii*” and two MAGs classified as family “*Ca. Paracaedibacteracea*” assembled from metagenomes of wastewater samples (24,25). A8/A10 and B2/H10/G10 form another clade with three MAGs assembled from metagenomes of water samples, including wastewater (22,26,27). “*Ca. Finniella inopinata*”, an endosymbiont of the amoebaflagellate *Viridiraptor invadens* branches at the base of this clade (23,28). We calculated pairwise AAI for all the genomes in the tree (Figure 3A, Additional File 6). The pairs B7 and F10, A8 and A10, and G10 and H10 all have a pairwise AAI greater than or equal to 95 %, considered a cutoff for species delimitation in the literature (29,30), indicating that they are most likely the same species. B2 is more distant from G10 and H10 with AAI values of 58.8 and 59.62, respectively (Additional file 6). However, CheckM results show that this genome contains contamination, which could be interfering with these analyses. The AAI boundary for genus delimitation varies depending on the taxa under consideration (31–34). Here, we used a cutoff of 55%, as was used in Midha et al. (2021) when describing “*Ca. B. vickermanii*”. This cutoff place clade B7/F10 as a novel species in the same genus as “*Ca. B. vickermanii*”, while clade A8/A10 and G10/H10 form two novel species in a novel genus.

We also compared ANI and AF values for all pairwise comparisons between the *Bodo* spp. endosymbionts and “*Ca. B. vickermanii*” (Figure 3B). ANI and AF are high within clades (ANI >97%, AF >85%), but, between the clades, only A8/A10 and H10/G10 are close enough to compare, with ANI values of 85.42-86.31 % and AF at 27.73-29.52 %, indicating that like their hosts, these genomes share high sequence similarity only over a small proportion of their genomes. The B2 genome shows high ANI (> 99%) with the genomes from H10 and G10, but low AF (<25%) indicating that this genome is only similar to those genomes over a small proportion of the genome. However, contamination in this genome may mean that this result is an artifact. The AF values shown in Figure 3B are the average of two AF values calculated for each pairwise comparison. A matrix with both AF values calculated for each pairwise comparison is in Additional File 7.

Finally, we compared the number of shared orthogroups between the Holosporales genomes and “*Ca. B. vickermanii*” (Figure 3C). 23.2% of orthogroups are shared by all genomes. The genomes that are closest in the phylogeny, A8/10 and B2/G10/H10, share the highest proportion of orthogroups at 52.6 %. Genomes B2/G10/H10 have the highest proportion of unique orthogroups at 12.35, however, the contamination in the B2 genome is likely to be inflating this number.

In summary, *B. saltans* and each novel *Bodo* clade presented in this paper harbour a unique species of Holosporales endosymbiont. These seven endosymbionts split into two genera (genus “*Ca. Bodocaedibacter*” and a novel genus) that appear congruent with the phylogeny of the host and further support the conclusion that these are three novel *Bodo* species.

## **The *Bodo* Holosporales differ in their metabolic capacity.**

We compared the secretion systems and metabolic modules present in the bacterial genomes and compared them to “*Ca. B. vickermanii*” (Figure 4). Genus “*Ca. Bodocaedibacter*” (B7/F10) and the novel genus described in this study (A8/A10 and B2/G10/H10) encode a SEC-SRP and a Type VI secretion system (18). In addition, the novel genus encodes two out the four proteins of the TAT secretion system, while a partial Type IV conjugal transfer pilus assembly protein systems is spottily distributed across the taxa.

All seven genomes have a partially complete ( $\geq 66\%$  complete) pathway for lysine biosynthesis, which is absent in all seven hosts ( $\leq 22\%$  complete) (Figure 4). In addition, the novel genus encodes several metabolic pathways absent in genus “*Ca. Bodocaedibacter*”, notably including heme biosynthesis (Figure 4). Heme is an essential nutrient, and to date, no complete heme pathway has been described for a Kinetoplastea member, which either require an exogenous source of heme, or contain endosymbiotic bacteria that produce it (35). Figure 4 only shows KEGG modules that were at least 50% complete in one or more genomes. A full list of all the KEGG modules, including those at less than 50% completeness is presented in Additional File 8.

Midha et al. (2021) reported that “*Ca. B. vickermanii*” encodes three putative polymorphic toxin/antitoxin systems which may be responsible for its host’s dependency (18). Polymorphic toxin/antitoxin systems are typically composed of a large multi-domain protein containing an N-terminal secretion signal followed by a toxin domain, a protective immunity protein, and multiple cassettes encoding alternative toxic domains and associated immunity proteins (19). The toxin genes can contain a homologous repeat region that enables recombination between the full-length toxin gene and the alternative toxins. We

identified three genes belonging to one putative toxin/antitoxin system in both B7 and F10, but none in the other five genomes from the novel genus. These genes show some of the characteristics of a polymorphic toxin/antitoxin system including a homologous region. However, the putative large multidomain protein in B7 does not contain an N-terminal signal peptide, and the one putative orphan module in both B7 and F10 contains a toxin but no cognate anti-toxin (Additional File 9). Therefore, the locus' identity and function as a polymorphic toxin/antitoxin system is tentative. In addition, the incompleteness of the single cell genomes presented here raises the possibility that key proteins and/or pathways are missing from these assemblies due to under sampling.

## **Genus *Bodo* encodes many unique protein families, but core metabolism is generally conserved with parasitic Kinetoplastea**

We next compared orthogroups distributed across the Kinetoplastea lineage (Figure 5). For each kinetoplastid genome we tallied the total number of orthogroups in that genome, and the number of orthogroups unique to that genome (Figure 5). Species on long branches or located at the base of lineages have the highest number of unique orthogroups, with *B. saltans* lake Konstanz having the most unique orthogroups. We next calculated the total number of orthogroups and the number of unique orthogroups in three kinetoplastid genera, *Bodo*, *Leishmania*, and *Trypanosoma* (Figure 5). Consistent with previous studies showing gene number reduction in parasitic lineages compared to *B. saltans* (11), genus *Bodo* has the greatest number of orthogroups, and the highest proportion of unique orthogroups, 53%, compared to 35% and 5.5 % in *Trypanosoma* and *Leishmania*, respectively.

Finally, we compared the completeness of metabolic pathways present in *Bodo* spp. and kinetoplastid species available in the Kyoto Encyclopedia of Genes and Genomes (KEGG) database (Figure 6). The four *Bodo* spp. are nearly identical in their metabolic capacity, with almost no KEGG modules missing from one lineage and present in another (Figure 6). Exceptions are Arginine biosynthesis and Riboflavin biosynthesis, which are missing in *B. saltans* lake Konstanz and partially present (25-80% complete) in the other species. The four *Bodo* spp. also share most of their metabolic capacity with the parasitic lineages, as previously reported (11). However, we do observe some pathways that are complete or nearly complete ( $\geq 80\%$  complete) in the *Bodo* lineage and absent or nearly absent ( $\leq 20\%$  complete) in *Trypanosoma* and *Leishmania*. Namely, Tryptophan metabolism, Tyrosine degradation, Molybdenum cofactor biosynthesis, and Pyrimidine degradation (uracil  $\Rightarrow$  beta-alanine, thymine  $\Rightarrow$  3-aminoisobutanoate). Conversely, the Glyoxylate cycle is partial (60 % complete) in the *Trypanosoma* and *Leishmania*, and near absent in all *Bodo* spp., consistent with previous findings in *B. saltans* (36), while Nucleotide sugar biosynthesis (galactose  $\Rightarrow$  UDP-galactose) is partially complete (50 %) in all *Leishmania* and *T. cruzi*, but absent in all *Bodo* spp. Figure 6 only shows modules that are at least 60% complete in at least one or more genomes. The full list of all KEGG modules identified in each genome, including those less than 50% is in Additional File 10.

## Discussion

In this study, we show that single-cell sequencing reveals unexpected levels of diversity among uncultured candidate *Bodo* spp. cells from a single environmental sample. Metabarcoding, particularly of SSU rDNA, and metagenomics have become the norm for

assessing the diversity of microbial eukaryotes in environmental samples. For protists, traditionally these methods involve clustering reads and/or amplicons into Operational Taxonomic units (OTUs) based on similarity thresholds, usually at around 97-99% identity, or into Amplicon Sequence Variants (ASVs) produced by the DADA2 pipeline (37). The seven single cells presented here all have a pairwise SSU rDNA identity greater than 99% and would therefore have been clustered into one OTU in some metabarcoding pipelines. Instead, by generating single-cell genomes, we show that these seven cells form three distinct clades, with each clade potentially representing a novel species. ANI has been used to compare distances within and between closely-related species of eukaryotic microbes including microsporidia, yeast and some protists, with an interspecies ANI cutoff of 95% considered appropriate (38–40). Here, we found that genomes within each of the three putative *Bodo* species have an ANI > 98%, supporting the conclusion that each clade represents a *Bodo* species. However, ANI is not useful when comparing more dissimilar genomes, where AAI becomes a more useful metric (41,42). While more commonly used to compare prokaryotic genomes, AAI has also been applied to eukaryotes. A comparison of 1,196 Human and Mouse protein sequences showed an AAI of 85% (43), while recent studies in fungal lineages found AAI values for members of the same species were often >97% and demonstrated an estimated a genus boundary threshold of ~70-75% AAI for the family Hypoxylaceae (Ascomycota) (44,45). While further work is needed to explore appropriate species and genus cutoffs for AAI and ANI within and between protist lineages, our results showing that AAI within each *Bodo* species was > 97% suggest that the species thresholds of 95-98% used in bacteria and fungi may be appropriate here too.

All three novel *Bodo* species presented here are more similar by SSU rDNA identity to *B. edax* ATCC strain 30903 than they are to *B. saltans* lake Konstanz. However, as no



other molecular sequence data exists from this organism, we are unable to determine if any or all are indeed *B. edax*. Callahan et al. (2002) mentions that *B. edax* contains endosymbiotic bacteria and differs morphologically from *B. saltans* by lacking mastigonemes on the anterior flagellum (5), yet several authors have proposed that *B. edax* be reclassified as an isolate of *B. saltans* (2,10). Given the extensive genetic diversity observed among the *Bodo* spp. presented in this paper, more work is needed to determine if *B. edax* is a true species. Importantly, this study suggests that the SSU rDNA locus is insufficient for species delimitation within genus *Bodo* as it does not reflect the genomic diversity within the group.

Further supporting our conclusion that the three *Bodo* clades represent three species is the finding that each putative species harbors a unique species of Holosporlaes bacterium. The three novel bacterial species split into two genera, one species belonging to genus “*Ca. Bodovickermani*, and the other two species forming a novel genus. The phylogeny of the host *Bodo* spp. appears congruent with the phylogeny of the endosymbionts, however more sampling covering greater taxonomic distribution is needed to test if this congruence indicates co-evolution, or is the result of other processes (46,47). Infectivity has been demonstrated in the *Holosporaceae* family of Holosporales, endosymbionts of various *Paramecium* ciliates (16,48), and some Holosporales have demonstrated the ability to invade novel hosts experimentally (49,50). However, the distribution and genetic differences observed between the Holosporales genomes presented here suggest that these endosymbionts are and have been associated with their respective hosts for a long period.

368           The nature of the symbioses between *Holosporales* bacteria and their hosts is  
369   complex and varied (13). Midha et al. (2021) hypothesized that *B. saltans* is dependent on  
370   its endosymbiont due to the presence of three putative addiction polymorphic toxin/anti-  
371   toxin systems encoded in the “*Ca. B. vickermanii*” genome (18). These systems typically  
372   encode a long-lived toxin molecule alongside an antitoxin with a shorter half-life, so that if  
373   the system is lost, the toxin is activated and becomes lethal. We found evidence for a  
374   putative toxin/anti-toxin system in the two genomes belonging to genus “*Ca.*  
375   *Bodovickermanii*”, but not in five genomes in the novel genus. However, the identity and/or  
376   functionality of the toxin/anti-toxin systems described in this study is speculative, as the  
377   proteins and loci differ from those in “*Ca. B. vickermanii*” by encoding fewer orphan  
378   modules. The lack of a similar system in the genomes of the novel *Holosporales* genus  
379   presented in this study suggests that alternative processes may be maintaining this symbiosis.  
380   All kinetoplasts including *B. saltans* lack the biosynthetic pathway required for heme  
381   biosynthesis (36,51), as do the *Bodo* spp. presented in this study. The Kinetoplastids  
382   *Angomonas* and *Strigomonas* harbour endosymbiotic beta-proteobacteria which provide  
383   their hosts with nutrients including heme, essential amino acids, and vitamins (35,52,53).  
384   We did find partially complete pathways for heme biosynthesis ( $\geq 70\%$  complete) in the  
385   novel genus, and a partially complete pathway for Lysine biosynthesis ( $\geq 66\%$  complete) in  
386   all the *Bodo* *Holosporales* genomes. Therefore, the two *Holosporales* genera presented here  
387   could demonstrate two alternative symbiotic strategies; with genus “*Ca. Bodovickermanii*”  
388   becoming addictive to its host through the toxin/antitoxin system, while the novel genus is  
389   possibly a source of heme and/or other essential nutrients. All seven *Holosporales* genomes  
390   presented in this paper encode a Type VI secretion system, conserved throughout  
391   *Holosporales* and hypothesized to play a role in host-endosymbiont interactions (13).

392

393           Previous studies have shown that *B. saltans* shares the majority of its metabolic  
394 pathways with the parasitic lineages *Trypanosoma* and *leishmania* (11,36). Our results show  
395 that the metabolic capacity of *B. saltans* is shared with the novel *Bodo* spp. presented here.  
396 We also show that the *Bodo* genus contains an abundance of unexplored and uncategorized  
397 protein diversity. The lineage contains far more unique orthogroups and proteins than the  
398 parasitic lineages. However, as most of these protein sequences have no known homology  
399 to functional domains in databases such as UniProt and KEGG; the functional implications of  
400 this unique gene repertoire remains unknown.

401

## 402   **Conclusions**

403           Most of our knowledge of protist genomic diversity and their symbionts comes from  
404 studies of species that are culturable. However, we know that environmental samples  
405 contain a wealth of underexplored diversity. This study uses single-cell sequencing and  
406 comparative genomics to show that seven *Bodo* spp. cells from a single environmental  
407 sample represent three potentially novel species, each harbouring a novel and unique  
408 species of bacterial endosymbiont. Comparing these data, we demonstrate the varied  
409 nature of these symbioses and show that single-cell sequencing is a powerful method for  
410 exploring the diversity of uncultured protists and their cobionts.

411

## Methods

### Sample collection

Surface water (~1 m) was collected from the River Leam (52.287295, -1.547563), Royal Leamington Spa (UK) in August 2022. Initially, the sample was prefiltered to remove larger debris and then concentrated using different polycarbonate filter pore sizes (Millipore) to obtain concentrated subsamples of different protist size ranges (10 to 40 µm and 0.8 to 5 µm). The subsamples were supplemented with 2-3 autoclaved barley grains to support heterotrophic/mixotrophic growth via bacteria increase and incubated for one week prior to cell sorting. Single-cell nuclei were stained with 1xSybrGreen for 10 minutes and sorted into 96-well microplates (pre-filled with 5ul autoclaved/sterile filtered media), using fluorescence-activated cell sorting (FACS; flow rate=1) and selecting against chlorophyll a-negative cells while selecting for SybrGreen-positive cells. After cell sorting, 10 µl of RLT-plus lysis buffer (Qiagen) was added to the wells and the plate was frozen at -80°C until further processing.

### Whole genome amplification, library preparation, and sequencing

A modified G&T-seq protocol (54) was carried out as follows. Using a magnetic separator, Dynabeads MyOne Streptavidin C1 (Invitrogen) beads were washed according to the manufacturer's guidance and then incubated with 2 × Binding & Wash buffer (10 mM Tris-HCl pH 7.5, 1 mM EDTA, 2 M NaCl) and Biotinylated Oligo-dT primer (IDT, 5'-/BiotinTEG/AAG CAG TGG TAT CAA CGC AGA GTA CTT TTT TTT TTT TTT TTT TTT TTT TVN-3') at 100 µM for 30 minutes at room temperature on a rotator. The oligo-treated

beads were washed four times in 1 × Binding & Wash buffer (5 mM Tris-HCl pH 7.5, 0.5 mM EDTA, 1 M NaCl) and then suspended in 1 × SuperScript II First Strand Buffer (Invitrogen) supplemented with SUPERaseIn RNase Inhibitor (Invitrogen) to a final concentration of 1 U/μl. The lysate was thawed on ice. 10 μl of prepared oligo-dT beads was added to each well containing 12 μl cell lysate using a Dragonfly Discovery liquid dispenser (SPT Labtech). The lysate plate was sealed and incubated on a ThermoMixer C (Eppendorf) with a heated lid at 21°C for 20 minutes shaking at 1000 rpm. Using a Fluent 480 liquid handling robot (Tecan) and a Magnum FLX magnetic separator (Alpaqua), the lysate supernatant was transferred to a new plate, and the beads were washed twice in a custom wash buffer (50 mM Tris-HCl pH 8.3, 75 mM KCl, 3 mM MgCl<sub>2</sub>, 10 mM DTT, 0.5% Tween-20). The supernatant from the washes was added to the left-over cell lysate containing the genomic DNA which was stored at -20°C overnight. The mRNA was not used for this study. The remaining cell lysate was thawed and subjected to a 0.6 × vols Ampure XP clean-up with 80% ethanol. The bead-bound gDNA was isothermally amplified for 3 hours at 30°C then 10 minutes at 65°C using a miniaturised (1/5 vols) Repli-g Single-Cell assay (Qiagen). The amplified gDNA was cleaned up with 0.8 × vols Ampure XP and 80% ethanol, then eluted in 10 mM Tris-HCl.

Sequencing libraries for this project were constructed by the Technical Genomics Group at the Earlham Institute, Norwich, UK. Initial libraries were constructed for shallow depth sequencing as follows: gDNA was quantified by fluorescence (Quant-iT HS-DNA, Invitrogen) on an Infinite Pro 200 plate reader (Tecan) then normalised to a final concentration of 0.2 ng/μl in 10 mM Tris-HCl. The Mosquito HV and Dragonfly Discovery liquid handling instruments (SPT Labtech) were used to prepare miniaturised (1/12.5 vols) Nextera XT (Illumina) dual-indexed sequencing libraries as follows: A tagmentation mastermix comprising two parts TD buffer (Illumina) and one part ATM (Illumina) was

prepared. Using the Dragonfly Discovery, 1.2 µl tagmentation mastermix was dispensed per well to a 384-well skirted PCR plate (Eppendorf). Using the Mosquito HV, 0.4 µl of normalised DNA was transferred to each well containing tagmentation mastermix. The plate was sealed and spun down then incubated at 55°C for 10 minutes on a thermal cycler. Using the Dragonfly Discovery, 0.4 µl 0.2% SDS was dispensed to each reaction. The plate was sealed, spun down and then incubated at room temperature for 5 minutes. Using the Dragonfly Discovery 1.2 µl NPM (Illumina) was added to each reaction. Using the Mosquito HV, 0.8 µl index primers pairs (i5 + i7) at a concentration of 0.5 µM was added to each reaction ensuring a unique combination in each well. The libraries were amplified under the following thermal cycler conditions: 72°C for 3 minutes, 95°C for 30 seconds, 12 cycles (95°C for 10 seconds, 55°C for 30 seconds, 72°C for 60 seconds), 72°C for 5 minutes, 4°C hold. The libraries were pooled and cleaned up using 0.8 × vols Ampure XP and 80% ethanol. The library pools were eluted in 20 µl 10mM Tris-HCl and assessed using a Bioanalyzer HS DNA assay (Agilent), HS DNA Qubit assay (Invitrogen) and finally an Illumina Library Quantification Kit assay (KAPA). These reads from these libraries were used for taxonomic assessment of the cells.

After taxonomic assessment deep sequencing libraries for the *Bodo* cells were constructed using the KAPA High Throughput Library Prep Kit (Roche Part No: KK8234/07961901001). Where possible, 1µg of genomic DNA was sheared to 450bp using the Covaris ML230 Sonicator (Covaris) and the ends of the DNA were repaired; 3' to 5' exonuclease activity removed the 3' overhangs and the polymerase activity filled in the 5' overhangs creating blunt ends. A single 'A' nucleotide was added to the 3' ends of the blunt fragments to allow for the ligation of barcoded adapters (6bp - Perkin Elmer NEXTFLEX DNA Barcodes 1-48 (NOVA-514101/2/3/4)) or (12bp - Perkin Elmer NEXTFLEX-HT (NOVA-

51474/5/6/7)) at a concentration of 6µM prior to a 0.8x clean up using Beckman Coulter AMPure XP beads (A63882). The size of the libraries was estimated using an Agilent High Sensitivity DNA chip (5067-4626) and the concentrations were quantified by fluorescence with a High Sensitivity Qubit assay (ThermoFisher Q32854).

Both shallow and deep sequencing libraries were sequenced on either an Illumina NovaSeq 6000 (cells A8, A10, B2, F10, G10, H10) with SP Reagent Kit v1.5 kit (300 cycles), , using Real-time Analysis (RTA) (version 3.4.4) and Control Software (version 1.7.5) or an Illumina NovaSeq X Plus (cell B7) with 10B Reagent Kit (300 Cycle) using RTA (version 4.6.7) and Control Software (version 1.2.2) to produce 150 bp paired-end reads. The resulting BCL files were converted to fastq with bcl2fastq (version 2.0).

### **Genome assembly, curation and classification**

Sequencing reads were trimmed using Trim Galore (version 0.6.6) (<https://github.com/FelixKrueger/TrimGalore>) with Cutadapt (version 3.4) (55). Genome assemblies were generated using SPAdes (version 3.15.5) (56) with single-cell mode enabled (--sc) and k-mer sizes 21, 33, 55, and 77.

Scaffolds less than 1,000 nt long were discarded from the assembly. Each assembly was manually curated and contaminant scaffolds/bins were removed using a combination of metagenomic binning with MetaBAT2 (57) based on tetra-nucleotide frequencies and taxonomic classification with CAT (v5.2) (58), Blobtools (v 1.1.1) (59) and Tiara (v1.0.1) (60) and EukRep (v0.6.6)(61). Bins that were classified as majority eukaryotic by all four classifiers and majority “Euglenozoa” by Blobtools were retained, as well as any unbinned scaffold classified as “Euglenozoa” by Blobtools or CAT.

The Holosporales bins were first identified from the taxonomic classifications described above. However, all raw assembly bins were assessed for completeness using CheckM (v1.1.2) (62), and all bins > 70% complete were further classified GTDB-TK (v2.3.2) on KBase. In all the cases, only one bin in each assembly was classified as Alphaproteobacteria by GTDB-TK, and was the bin retained as the *Holosporlaes* endosymbiont MAG. The retained scaffold sets were manually checked which resulted in the removal of one scaffold from the B2 Holosporales bin that was a contaminate from another well on the 96 well plate.

The general statistics, classification summaries, CheckM results for all genome bins, and the GTDB-TK classifications are tabulated in Additional File 5.

The SSU rDNA BLASTs were done through Geneious Prime. In each case, the subject sequence with the highest bitscore was considered the top hit.

## **Gene prediction annotation**

Eukaryotic gene prediction was done using Companion (v2.2.11) (63) at the WebServer, with default settings and using *Bodo saltans* as the reference. rDNAs were annotated with Barrnap (v0.9). For one sample (B7), the SSU rDNA sequence was annotated as two overlapping fragments, which were manually merged into one sequence by alignment in Geneious Prime. Four of the seven SSU rDNA sequences are truncated (See Table 1 and Additional File 1). Three of these, B2, G10, and H10 were manually extended by alignment of the scaffolds and truncated SSU rDNA sequences annotated by Barrnap in Geneious Prime and extracting the longest region with shared homology from each scaffold.



All three sequences were extended by ~70 bp. For G10, the SSU rDNA annotation was truncated due to end of the scaffold, while B2 and H10 are truncated due to loss of homology, possibly due to miss-assembly of those scaffolds. Holosporales gene predictions were done using PROKKA (v1.14.6) (64).

Completeness of the genomes and protein sets was assessed with BUSCO (v5.3.2) (65).

### **Phylogenetic analyses of SSU rDNA sequences**

SSU rDNA sequences were collected from GeneBank, filtering for sequences that were at least 1,000 nucleotides long (Additional File 9). The sequences were aligned with MAFFT (v7.520) using the FFT-NS-i model (66). The alignment was trimmed manually in Geneious Prime to remove all positions with less than 50% coverage. The untrimmed alignment used to build the tree is in Additional file 1, while the sequence accessions are in Additional file 11. The Maximum-Likelihood (ML) tree was built with IQ-TREE (v2.3.2) (67) using the TIM3e+I+R3 model, which was the best fit model determined by ModelFinder (68), with 1,000 non-parametric bootstrap replicates, and rooting at two outgroup species *Dimastigella trypaniformis* and *Rhynchomonas nasuta*. The tree was visualized and plotted with its associated distance matrix using Interactive Tree Of Life (iTOL) (v7.2)(69).

### **Phylogenomic analyses**

For the Kinetoplastea, protein sets for related species were collected from GeneBank (Additional File 12). Orthologous gene sets were identified using Orthofinder (v2.5.4) (70).

Protein sequences for shared single copy orthogroups (n = 488) were aligned with MAFFT (v7.520) using the FFT-NS-2 model (66), and trimmed with TrimAl (v2.0) using the gappyout option (71). The trimmed alignments were concatenated with AMAS (72) into a matrix with 234,727 amino acid sites. The ML tree was produced from a partitioned analysis undertaken using IQ-TREE (v2.3.2) with a partitioning scheme that merged the 488 proteins into 20 partitions, with rooting at the outgroup *Perkinsella* sp.. Model selection was performed by ModelFinder, and 1,000 non-parametric bootstraps were run.

For the Holosporales, protein sets were collected from GeneBank for related species and several closely related MAGs (Additional File 10). For one MAG, GCA\_002422845.1, a protein set was not available on GeneBank, so an annotation was done using PROKKA (v1.14.6). Orthologous gene sets were identified using Orthofinder (v2.5.4). Protein sequences from shared single copy orthogroups (n = 24) were aligned with MAFFT (v7.520) using the FFT-NS-2 model and trimmed with TrimAl (v2.0) using the gappyout option. The trimmed alignments were concatenated with AMAS into a matrix with 8,004 amino acid sites. The ML tree was produced from a partitioned analysis undertaken using IQ-TREE (v2.3.2) with a partitioning scheme that merged the 24 proteins into 6 partitions, with rooting at the outgroup containing *Temperatibacter marinus* and *Kordiimonas pumila*. Model selection was performed by ModelFinder, and 1000 non-parametric bootstraps were run.

The Kinetoplastea tree (Figure 5) is the Species Tree from All Genes (STAG) inferred species tree produced by Orthofinder. It is inferred from 331 gene trees from the orthogroups where all species are present.

All trees were visualized and plotted either using the Interactive Tree of Life (iTOL) or in R using the package ggtree (v3.14.0).

## Functional analyses

Pfams were assigned to each protein set using InterProScan (v5.52-86.0) using option -aap Pfams and with the Pre-calculated match lookup service disabled (----disable-precac). Pfam counts were tabulated in R. A PCA was run on the count matrix for *Bodo* spp. after removal of Pfam with no variation, which reduced the dataset from 2938 variables to 2545. The PCA was run in R using the function prcomp with scaling and centering.

KEGG identifiers were assigned to each protein set using KofamKOALA or kofamscan (v1.3.0) (73). The completeness of each KEGG module was calculated using kegg-pathway-completeness-tool (v1.3.0) (<https://github.com/EBI-Metagenomics/kegg-pathways-completeness-tool>).

Venn diagrams showing the proportion of shared orthogroups were generated in R using the package ggven (v0.1.10). For an orthogroup to be missing from a species, it must be missing from all genomes in that species. For an orthogroup to be present, it can be present in one or more genomes in that species.

The presence of bacterial secretion systems was assessed manually by visualizing the results of KofamKOALA on the KEGG website using the KEGG mapper tool Reconstruct. The number of unique components present in each protein set that mapped to each secretions system was counted and compared to the total number of unique components listed for each system. The one exception was for the Type VI system, which was annotated using a

combination of KEGG Reconstruct, BLASTp and psiblast, where the Type VI proteins from *Ca. B. vickermanii* were used as queries to search a database of the protein sets from all seven Holosporales MAGs. The query sequence identifiers, and the significant hit subject identifiers are listed in Additional File 13.

The toxin anti/toxin systems were investigated manually using BLASTp and psiblast, where the proteins of *Ca. B. vickermanii* systems were used as queries to search a database of the protein sets from all seven Holosporales MAGs. N terminal signal peptides were identified using SignalP v 6.0 at the webserver.

## AAI, ANI, and AF analyses

Pairwise amino acid identity (AAI) was calculated for the *Bodo* spp. using the aai.ruby script and BLAST+ v(2.16.0) that is part of the Enveomics Collection at <https://github.com/lmrodriguezr/enveomics> (74). For the Holosporales, pairwise AAI was calculated using FastAAI (34). ANI and AF were calculated using the tool Skani (v0.2.2) (21).

## Figures

All plots were generated in R and edited for publication using Illustrator.

## Declarations

## Availability of data and materials

The raw reads and annotated assemblies for Bodo and Bodo Holosporales have been deposited in the European Nucleotide Archive under accession PRJEB97217. The assembly sequence files, protein sequence files and gff files used in this study have been deposited in Zenodo at 10.5281/zenodo.16948154.

## **Competing interests**

The authors declare that there are no conflicts of interest.

## **Funding**

This work was funded by the Wellcome Trust Darwin Tree of Life Awards (218328 and 226458), and by the Biotechnology and Biological Sciences Research Council (BBSRC), part of UK Research and Innovation, through the Earlham Institute's Core Capability Grants (BB/CCG1720/1 and BB/CCG2220/1), its Strategic Programme Grant Decoding Biodiversity (BBX011089/1) and its constituent Work Package 2 BBS/E/ER/230002B, its National Capability BBS/E/T/000PR9816 (NC1 - Supporting EI's ISPs and the UK Community with Genomics and Single Cell Analysis), and Transformative Genomics, National Bioscience Research Infrastructure (BBS/E/ER/23NB0006). TAR is supported by a Royal Society University Research Fellowship (URF/R/191005). Part of this work was delivered with support for the physical HPC infrastructure and data center delivered via the NBI Computing infrastructure for Science (CiS) group.

## **Authors' contributions**

Sally D. Warring, Conceptualization, Data curation, Formal analysis, Investigation,  
Methodology, Resources, Visualization, Writing | Jamie McGowan, Conceptualization, Data  
curation, Investigation, Methodology, Resources, Software, Writing | Estelle S. Kilias,  
Investigation, Methodology, Resources, Writing | James Lipscombe, Investigation,  
Methodology, Resources, Writing | Elisabet Alacid, Investigation, Methodology, Resources,  
Writing | Tom Barker, Investigation, Methodology | Leah Catchpole, Investigation,  
Methodology, Project administration, Writing | Seanna McTaggart, Funding acquisition,  
Project administration, Writing | Karim Gharbi, Funding acquisition, Methodology,  
Resources, Supervision, Writing | Thomas A. Richards, Conceptualization, Funding  
acquisition, Resources, Supervision, Writing | David Swarbreck, Conceptualization, Funding  
acquisition, Supervision | Neil Hall, Conceptualization, Funding acquisition, Resources,  
Supervision, Writing.

## Acknowledgements

We would like to acknowledge the members of the Technical Genomics and Core  
Bioinformatics groups at the Earlham Institute, and note the specific contributions of Chris  
Watkins, Sacha Lucchini, Kendall Baker, and Neil Shearer. We also acknowledge the work  
delivered via the Laboratory Managers and Research Computing Groups at EI who manage  
and deliver High Performance Computing at EI.

## References

1. Adl SM, Simpson AGB, Lane CE, Lukeš J, Bass D, Bowser SS, et al. The Revised Classification of Eukaryotes. *J Eukaryot Microbiol.* 2012 Sept;59(5):429–514.

- 658 2. Moreira D, López-García P, Vickerman K. An updated view of kinetoplastid phylogeny using  
659 environmental sequences and a closer outgroup: proposal for a new classification of the class  
660 Kinetoplastea. *Int J Syst Evol Microbiol.* 2004 Sept 1;54(5):1861–75.
- 661 3. Simpson AGB, Stevens JR, Lukeš J. The evolution and diversity of kinetoplastid flagellates. *Trends*  
662 *Parasitol.* 2006 Apr;22(4):168–74.
- 663 4. Simpson AGB, Lukeš J, Roger AJ. The Evolutionary History of Kinetoplastids and Their  
664 Kinetoplasts. *Mol Biol Evol.* 2002 Dec 1;19(12):2071–83.
- 665 5. Callahan HA, Litaker RW, Noga EJ. Molecular Taxonomy of the Suborder Bodonina (Order  
666 Kinetoplastida), Including the Important Fish Parasite, *Ichthyobodo necator*. *J Eukaryot*  
667 *Microbiol.* 2002 Mar;49(2):119–28.
- 668 6. Deschamps P, Lara E, Marande W, Lopez-Garcia P, Ekelund F, Moreira D. Phylogenomic Analysis  
669 of Kinetoplastids Supports That Trypanosomatids Arose from within Bodonids. *Mol Biol Evol.*  
670 2011 Jan 1;28(1):53–8.
- 671 7. Brooker BE. Fine structure of *Bodo saltans* and *Bodo caudatus* (Zoomastigophora: Protozoa) and  
672 their affinities with the Trypanosomatidae. *Bull Br Mus Nat Hist Zool.* 1971;22(3):89–102.
- 673 8. Vickerman K, Preston T. Comparative cell biology of the kinetoplastid flagellates. *Biol*  
674 *Kinetoplastida.* 1979;2:35–130.
- 675 9. Dolezel D, Jirků M, Maslov DA, Lukes J. Phylogeny of the bodonid flagellates (Kinetoplastida)  
676 based on small-subunit rRNA gene sequences. *Int J Syst Evol Microbiol.* 2000 Sept 1;50(5):1943–  
677 51.
- 678 10. Von Der Heyden S, Chao EE, Vickerman K, Cavalier-Smith T. Ribosomal RNA Phylogeny of  
679 Bodonid and Diplonemid Flagellates and the Evolution of Euglenozoa. *J Eukaryot Microbiol.* 2004  
680 July;51(4):402–16.
- 681 11. Jackson AP, Otto TD, Aslett M, Armstrong SD, Bringaud F, Schlacht A, et al. Kinetoplastid  
682 Phylogenomics Reveals the Evolutionary Innovations Associated with the Origins of Parasitism.  
683 *Curr Biol.* 2016 Jan;26(2):161–72.
- 684 12. Szokoli F, Castelli M, Sabaneyeva E, Schrällhammer M, Krenek S, Doak TG, et al. Disentangling  
685 the Taxonomy of Rickettsiales and Description of Two Novel Symbionts (“*Candidatus Bealeia*  
686 *paramacronuclearis*” and “*Candidatus Fokinia cryptica*”) Sharing the Cytoplasm of the Ciliate  
687 Protist *Paramecium biaurelia*. Drake HL, editor. *Appl Environ Microbiol.* 2016 Dec  
688 15;82(24):7236–47.
- 689 13. Castelli M, Petroni G. An Evolutionary-Focused Review of the Holosporales  
690 (Alphaproteobacteria): Diversity, Host Interactions, and Taxonomic Re-ranking as Holosporineae  
691 Subord. *Nov. Microb Ecol.* 2025 Mar 14;88(1):15.
- 692 14. Schrällhammer M, Potekhin A. Epidemiology of Nucleus-Dwelling Holospora: Infection,  
693 Transmission, Adaptation, and Interaction with *Paramecium*. In: Kloc M, editor. *Symbiosis:*  
694 *Cellular, Molecular, Medical and Evolutionary Aspects* [Internet]. Cham: Springer International  
695 Publishing; 2020 [cited 2025 Sept 24]. p. 105–35. (Results and Problems in Cell Differentiation;  
696 vol. 69). Available from: [https://link.springer.com/10.1007/978-3-030-51849-3\\_4](https://link.springer.com/10.1007/978-3-030-51849-3_4)

- 697 15. Schrällhammer M, Castelli M, Petroni G. Phylogenetic relationships among endosymbiotic R-  
698 body producer: Bacteria providing their host the killer trait. Syst Appl Microbiol. 2018  
699 May;41(3):213–20.
- 700 16. Fujishima M. Infection and Maintenance of Holospora Species in Paramecium caudatum. In:  
701 Fujishima M, editor. Endosymbionts in Paramecium [Internet]. Berlin, Heidelberg: Springer  
702 Berlin Heidelberg; 2009 [cited 2025 June 10]. p. 201–25. (Microbiology Monographs; vol. 12).  
703 Available from: [https://link.springer.com/10.1007/978-3-540-92677-1\\_8](https://link.springer.com/10.1007/978-3-540-92677-1_8)
- 704 17. Fujishima M, Kawai M, Yamamoto R. *Paramecium caudatum* acquires heat-shock resistance in  
705 ciliary movement by infection with the endonuclear symbiotic bacterium *Holospora obtusa*.  
706 FEMS Microbiol Lett. 2005 Feb;243(1):101–5.
- 707 18. Midha S, Rigden DJ, Siozios S, Hurst GDD, Jackson AP. Bodo saltans (Kinetoplastida) is dependent  
708 on a novel Paracaedibacter-like endosymbiont that possesses multiple putative toxin-antitoxin  
709 systems. ISME J. 2021 June;15(6):1680–94.
- 710 19. Jamet A, Nassif X. New Players in the Toxin Field: Polymorphic Toxin Systems in Bacteria. mBio.  
711 2015 July;6(3):e00285-15.
- 712 20. Scheckenbach F, Wylezich C, Mylnikov AP, Weitere M, Arndt H. Molecular Comparisons of  
713 Freshwater and Marine Isolates of the Same Morphospecies of Heterotrophic Flagellates. Appl  
714 Environ Microbiol. 2006 Oct;72(10):6638–43.
- 715 21. Shaw J, Yu YW. Fast and robust metagenomic sequence comparison through sparse chaining  
716 with skani. Nat Methods. 2023 Nov;20(11):1661–5.
- 717 22. Schneider D, Zühlke D, Poehlein A, Riedel K, Daniel R. Metagenome-Assembled Genome  
718 Sequences from Different Wastewater Treatment Stages in Germany. Thrash JC, editor.  
719 Microbiol Resour Announc. 2021 July 8;10(27):e00504-21.
- 720 23. Muñoz-Gómez SA, Hess S, Burger G, Lang BF, Susko E, Slamovits CH, et al. An updated phylogeny  
721 of the Alphaproteobacteria reveals that the parasitic Rickettsiales and Holosporales have  
722 independent origins. eLife. 2019 Feb 25;8:e42535.
- 723 24. Parks DH, Rinke C, Chuvochina M, Chaumeil PA, Woodcroft BJ, Evans PN, et al. Recovery of  
724 nearly 8,000 metagenome-assembled genomes substantially expands the tree of life. Nat  
725 Microbiol. 2017 Sept 11;2(11):1533–42.
- 726 25. Liu L, Wang Y, Yang Y, Wang D, Cheng SH, Zheng C, et al. Charting the complexity of the  
727 activated sludge microbiome through a hybrid sequencing strategy. Microbiome. 2021  
728 Dec;9(1):205.
- 729 26. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarszewski A, Chaumeil PA, et al. A standardized  
730 bacterial taxonomy based on genome phylogeny substantially revises the tree of life. Nat  
731 Biotechnol. 2018 Nov;36(10):996–1004.
- 732 27. Probst AJ, Ladd B, Jarett JK, Geller-McGrath DE, Sieber CMK, Emerson JB, et al. Differential depth  
733 distribution of microbial function and putative symbionts through sediment-hosted aquifers in  
734 the deep terrestrial subsurface. Nat Microbiol. 2018 Jan 29;3(3):328–36.



- 735 28. Hess S, Suthaus A, Melkonian M. “Candidatus Finniella” (Rickettsiales, Alphaproteobacteria),  
736 Novel Endosymbionts of Viridiraptorid Amoeboflagellates (Cercozoa, Rhizaria). Voordouw G,  
737 editor. Appl Environ Microbiol. 2016 Jan 15;82(2):659–70.
- 738 29. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI analysis of  
739 90K prokaryotic genomes reveals clear species boundaries. Nat Commun. 2018 Nov  
740 30;9(1):5114.
- 741 30. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. DNA-DNA  
742 hybridization values and their relationship to whole-genome sequence similarities. Int J Syst Evol  
743 Microbiol. 2007 Jan;57(Pt 1):81–91.
- 744 31. Konstantinidis KT, Tiedje JM. Towards a Genome-Based Taxonomy for Prokaryotes. J Bacteriol.  
745 2005 Sept 15;187(18):6258–64.
- 746 32. Rodriguez-R LM, Konstantinidis KT. Bypassing Cultivation To Identify Bacterial Species: Culture-  
747 independent genomic approaches identify credibly distinct clusters, avoid cultivation bias, and  
748 provide true insights into microbial species. Microbe Mag. 2014 Mar 1;9(3):111–8.
- 749 33. Park MJ, Kim YJ, Park M, Yu J, Namirimu T, Roh YR, et al. Establishment of Genome Based  
750 Criteria for Classification of the Family Desulfovibrionaceae and Proposal of Two Novel Genera,  
751 Alkalidesulfovibrio gen. nov. and Salidesulfovibrio gen. nov. Front Microbiol. 2022 May  
752 25;13:738205.
- 753 34. Gerhardt K, Ruiz-Perez CA, Rodriguez-R LM, Jain C, Tiedje JM, Cole JR, et al. FastAAI: efficient  
754 estimation of genome average amino acid identity and phylum-level relationships using  
755 tetramers of universal proteins. Nucleic Acids Res. 2025 Apr 22;53(8):gkaf348.
- 756 35. Harmer J, Yurchenko V, Nenarokova A, Lukeš J, Ginger ML. Farming, slaving and enslavement:  
757 histories of endosymbioses during kinetoplastid evolution. Parasitology. 2018  
758 Sept;145(10):1311–23.
- 759 36. Opperdoes FR, Butenko A, Flegontov P, Yurchenko V, Lukeš J. Comparative Metabolism of  
760 Free-living *Bodo saltans* and Parasitic Trypanosomatids. J Eukaryot Microbiol. 2016  
761 Sept;63(5):657–78.
- 762 37. Burki F, Sandin MM, Jamy M. Diversity and ecology of protists revealed by metabarcoding. Curr  
763 Biol. 2021 Oct;31(19):R1267–80.
- 764 38. Majda S, Boenigk J, Beisser D. Intraspecific Variation in Protists: Clues for Microevolution from  
765 *Poteroispumella lacustris* (Chrysophyceae). Archibald J, editor. Genome Biol Evol. 2019 Sept  
766 1;11(9):2492–504.
- 767 39. Lachance MA, Lee DK, Hsiang T. Delineating yeast species with genome average nucleotide  
768 identity: a calibration of ANI with haplontic, heterothallic *Metschnikowia* species. Antonie Van  
769 Leeuwenhoek. 2020 Dec;113(12):2097–106.
- 770 40. De Albuquerque NRM, Haag KL. Using average nucleotide identity ( ANI ) to evaluate  
771 microsporidia species boundaries based on their genetic relatedness. J Eukaryot Microbiol. 2023  
772 Mar;70(2):e12944.
- 773 41. Konstantinidis KT, Tiedje JM. Prokaryotic taxonomy and phylogeny in the genomic era:  
774 advancements and challenges ahead. Curr Opin Microbiol. 2007 Oct;10(5):504–9.

- 775 42. Qin QL, Xie BB, Zhang XY, Chen XL, Zhou BC, Zhou J, et al. A Proposed Genus Boundary for the  
776 Prokaryotes Based on Genomic Insights. *J Bacteriol.* 2014 June 15;196(12):2210–5.
- 777 43. Makołowski W, Zhang J, Boguski MS. Comparative analysis of 1196 orthologous mouse and  
778 human full-length mRNA and protein sequences. *Genome Res.* 1996 Sept;6(9):846–57.
- 779 44. Wibberg D, Rupp O, Blom J, Jelonek L, Kröber M, Verwaaijen B, et al. Development of a  
780 *Rhizoctonia solani* AG1-IB Specific Gene Model Enables Comparative Genome Analyses between  
781 Phytopathogenic *R. solani* AG1-IA, AG1-IB, AG3 and AG8 Isolates. Yu JH, editor. *PLOS ONE.* 2015  
782 Dec 21;10(12):e0144769.
- 783 45. Wibberg D, Stadler M, Lambert C, Bunk B, Spröer C, Rückert C, et al. High quality genome  
784 sequences of thirteen Hypoxylaceae (Ascomycota) strengthen the phylogenetic family backbone  
785 and enable the discovery of new taxa. *Fungal Divers.* 2021 Jan;106(1):7–28.
- 786 46. Groussin M, Mazel F, Alm EJ. Co-evolution and Co-speciation of Host-Gut Bacteria Systems. *Cell*  
787 *Host Microbe.* 2020 July;28(1):12–22.
- 788 47. Russo L, Miller AD, Tooker J, Bjornstad ON, Shea K. Quantitative evolutionary patterns in  
789 bipartite networks: Vicariance, phylogenetic tracking or diffuse co-evolution? *Methods Ecol Evol.*  
790 2018 Mar;9(3):761–72.
- 791 48. Schrällhammer M, Potekhin A. Epidemiology of Nucleus-Dwelling *Holospira*: Infection,  
792 Transmission, Adaptation, and Interaction with *Paramecium*. In: Kloc M, editor. *Symbiosis:*  
793 *Cellular, Molecular, Medical and Evolutionary Aspects* [Internet]. Cham: Springer International  
794 Publishing; 2020 [cited 2025 June 10]. p. 105–35. (Results and Problems in Cell Differentiation;  
795 vol. 69). Available from: [https://link.springer.com/10.1007/978-3-030-51849-3\\_4](https://link.springer.com/10.1007/978-3-030-51849-3_4)
- 796 49. Frelief PF, Loy JK, Kruppenbach B. Transmission of Necrotizing Hepatopancreatitis in *Penaeus*  
797 *vannamei*. *J Invertebr Pathol.* 1993 Jan;61(1):44–8.
- 798 50. Schulz F, Lagkouvardos I, Wascher F, Aistleitner K, Kostanjšek R, Horn M. Life in an unusual  
799 intracellular niche: a bacterial symbiont infecting the nucleus of amoebae. *ISME J.* 2014 Aug  
800 1;8(8):1634–44.
- 801 51. Kořený L, Lukeš J, Oborník M. Evolution of the haem synthetic pathway in kinetoplastid  
802 flagellates: An essential pathway that is not essential after all? *Int J Parasitol.* 2010  
803 Feb;40(2):149–56.
- 804 52. Alves JMP, Voegtly L, Matveyev AV, Lara AM, Da Silva FM, Serrano MG, et al. Identification and  
805 Phylogenetic Analysis of Heme Synthesis Genes in Trypanosomatids and Their Bacterial  
806 Endosymbionts. Lopez-Garcia P, editor. *PLoS ONE.* 2011 Aug 10;6(8):e23518.
- 807 53. Klein CC, Alves JMP, Serrano MG, Buck GA, Vasconcelos ATR, Sagot MF, et al. Biosynthesis of  
808 Vitamins and Cofactors in Bacterium-Harboring Trypanosomatids Depends on the Symbiotic  
809 Association as Revealed by Genomic Analyses. Parkinson J, editor. *PLoS ONE.* 2013 Nov  
810 19;8(11):e79786.
- 811 54. Macaulay IC, Haerty W, Kumar P, Li YI, Hu TX, Teng MJ, et al. G&T-seq: parallel sequencing of  
812 single-cell genomes and transcriptomes. *Nat Methods.* 2015 June;12(6):519–22.
- 813 55. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.  
814 *EMBnet.journal.* 2011 May 2;17(1):10–2.

- 815 56. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A New  
816 Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. *J Comput Biol.* 2012  
817 May;19(5):455–77.
- 818 57. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive binning algorithm  
819 for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ.*  
820 2019;7:e7359.
- 821 58. von Meijenfeldt FAB, Arkhipova K, Cambuy DD, Coutinho FH, Dutilh BE. Robust taxonomic  
822 classification of uncharted microbial sequences and bins with CAT and BAT. *Genome Biol.* 2019  
823 Oct 22;20(1):217.
- 824 59. Laetsch DR, Blaxter ML. BlobTools: Interrogation of genome assemblies. *F1000Research.* 2017  
825 July 31;6:1287.
- 826 60. Karlicki M, Antonowicz S, Karnkowska A. Tiara: deep learning-based classification system for  
827 eukaryotic sequences. *Bioinforma Oxf Engl.* 2022 Jan 3;38(2):344–50.
- 828 61. West PT, Probst AJ, Grigoriev IV, Thomas BC, Banfield JF. Genome-reconstruction for eukaryotes  
829 from complex natural microbial communities. *Genome Res.* 2018 Apr;28(4):569–80.
- 830 62. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of  
831 microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.* 2015  
832 July;25(7):1043–55.
- 833 63. Haese-Hill W, Crouch K, Otto TD. Annotation and visualization of parasite, fungi and arthropod  
834 genomes with Companion. *Nucleic Acids Res.* 2024 July 5;52(W1):W39–44.
- 835 64. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics.* 2014 July  
836 15;30(14):2068–9.
- 837 65. Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO Update: Novel and  
838 Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of  
839 Eukaryotic, Prokaryotic, and Viral Genomes. Kelley J, editor. *Mol Biol Evol.* 2021 Sept  
840 27;38(10):4647–54.
- 841 66. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements  
842 in performance and usability. *Mol Biol Evol.* 2013 Apr;30(4):772–80.
- 843 67. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-  
844 TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Mol*  
845 *Biol Evol.* 2020 May 1;37(5):1530–4.
- 846 68. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermini LS. ModelFinder: fast model  
847 selection for accurate phylogenetic estimates. *Nat Methods.* 2017 June;14(6):587–9.
- 848 69. Letunic I, Bork P. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree  
849 display and annotation tool. *Nucleic Acids Res.* 2024 July 5;52(W1):W78–82.
- 850 70. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative genomics.  
851 *Genome Biol.* 2019 Dec;20(1):238.

- 852 71. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment  
853 trimming in large-scale phylogenetic analyses. *Bioinformatics*. 2009 Aug 1;25(15):1972–3.
- 854 72. Borowiec ML. AMAS: a fast tool for alignment manipulation and computing of summary  
855 statistics. *PeerJ*. 2016 Jan 28;4:e1660.
- 856 73. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, et al. KofamKOALA: KEGG  
857 Ortholog assignment based on profile HMM and adaptive score threshold. Valencia A, editor.  
858 *Bioinformatics*. 2020 Apr 1;36(7):2251–2.
- 859 74. Rodríguez-R LM, Konstantinidis KT. The enveomics collection: a toolbox for specialized analyses  
860 of microbial genomes and metagenomes [Internet]. *PeerJ Preprints*; 2016 [cited 2025 Apr 1].  
861 Available from: <https://peerj.com/preprints/1900v1>

862

863

## Figure Legends

**Figure 1 – SSU rDNA phylogeny of *Bodo* spp.** **A.** Pairwise ANI of SSU rDNA sequences from the seven single cell genomes **B.** ML phylogeny of SSU rDNA sequences from *Bodo* spp. and related genera. The tree scale is substitutions per site. The seven single cell genomes are shaded yellow, blue and pink to distinguish the three clades they form. *B. saltans* lake Konstanz is shaded purple. The two columns on the right side of the tree show the pairwise ANI between each taxon and the SSU rDNA from *B. saltans* lake Konstanz and *B. edax*.

**Figure 2 – Genomic diversity of *Bodo* spp.** **A.** ML phylogeny of 488 single copy orthologous proteins. Tree scale is substitutions per site **B.** Heatmap showing pairwise average AAI values for all *Bodo* genomes **C.** Heatmap showing pairwise ANI (lower triangle) and AF (upper triangle) **D.** Number and proportion of shared orthogroups for each *Bodo* spp. **E.** Bar plot showing the proportion of protein annotations in each genome that have a Pfam annotation **F.** PCA plot generated from the tally of Pfam domain present in each genome. In all plots the three *Bodo* spp. from single cell genomes are shaded yellow, blue and pink to distinguish the three clades they form while *B. saltans* lake Konstanz is shaded purple.

**Figure 3 – Genomic diversity of Holosporales endosymbionts.** **A.** ML phylogeny of 24 single copy orthologous proteins. Tree scale is substitutions per site **B.** Heatmap showing pairwise ANI (lower triangle) and AF (upper triangle) for the Holosporales associated with *Bodo* single cells **C.** Number and proportion of shared orthogroups for each Holosporales species. In all

plot the Holosporales from the seven single cell genomes are shaded yellow, blue and pink to distinguish the three clades they form while “*Ca. B. vickermanii*” is shaded purple.

**Figure 4 – Functional diversity among Holosporales endosymbionts.** Heatmap in blue showing the completeness values for KEGG modules found in each *Bodo* Holosporales genome and *Ca. B. vickermanii* genome. Only modules that are at least 50% complete in one or more genome are shown. Heatmap in red shows the number of proteins belonging to each of the bacterial secretion systems found in the genomes. The block to the right of the heatmap shows the total number of proteins belonging to each secretion system, as listed on the KEGG Brite database.

**Figure 5 – Genomic uniqueness in Genus *Bodo*.** Species tree inferred by STAG. Tree scale is substitutions per site. The support values are the number of individual gene trees that contain that bipartition. The bar plots show the total number of orthogroups and the number of unique orthogroups in each genome protein set. Pie charts are shaded to show the proportion of orthogroups unique to the three genera *Bodo*, *Trypanosoma* and *Leishmania*.

**Figure 6 – Functional comparison of Kinetoplastida.** Heatmap showing the completeness values for KEGG modules in each genome. Only modules that are at least 60% complete in one or more genomes are shown.

## 907 Additional files

908 **Additional File 1 - *Bodo* and related spp. SSU rDNA alignment.** FASTA file containing the  
909 untrimmed SSU rDNA alignment used to build the tree in Figure 1.

910 **Additional File 2 - AAI summary statistics for *Bodo* spp.** Excel sheet listing the summary  
911 metrics output from the aai.rb script. Column “no. proteins used” shows that number of  
912 pairwise comparisons used to calculate the average AAI.

913 **Additional File 3 – *Bodo skani* AF full matrix.** TSV file showing the AF calculated by skani in  
914 both directions for each pairwise comparison.

915 **Additional File 4 - Pfam counts and PCA loadings.** Excell workbook with two sheets. The  
916 first sheet shows the Pfam frequencies in each *Bodo* genome. The second sheet shows the  
917 loadings given to each Pfam for PC1-8.

918 **Additional File 5 - *Bodo* SC raw genome assemblies statistics per bin.** Excell sheet  
919 tabulating statistics for all the raw single-cell genome assemblies, split by MetaBat2 bins.

920 **Additional File 6 - FastAAI distance matrix.** Excell sheet containing the AAI values for the  
921 Holsoporales heatmap in Figure 3A.

922 **Additional File 7 – *Holosporlaes skani* AF full matrix.** TSV file showing the AF calculated by  
923 skani in both directions for each pairwise comparison.

924 **Additional File 8 - *Holosporales* all KEGG completeness.** Excell sheet showing the  
925 completeness values for KEGG modules identified in each *Holosporales* protein set.

926 **Additional File 9 - Toxin antitoxins in *Bodo Holosporlaes* B7 F10 figure.** PDF document  
927 showing a: A. schematic of the putative toxin/antitoxin loci in B7 and F10 B. alignment  
928 between the large toxin protein and alternative toxin, showing region of homology C.  
929 SingnalP plots of N termini of the putative multi-domain toxins from B7 and F10.

930 **Additional File 10 - Kinetoplastids all KEGG completeness.** Excell sheet showing the  
931 completeness values for KEGG modules identified in each *Bodo* and Kinetoplastea protein  
932 set.

933 **Additional File 11 - SSU rDNA sequences used in phylogeny.** Excell sheet with NCBI  
934 accession identifiers for all SSU rDNA sequences used to construct the phylogeny in Figure 1.

935 **Additional File 12- all NCBI protein sets.** Excell workbook with two sheets listing the NCBI  
936 accession identifiers for all protein sets used in this study. Kinetoplastea sets are on sheet 1  
937 and *Holosporales* sets are on sheet two.

938 **Additional File 13 - TypeVI secretion system protein IDs.** Excell sheet listing the Type VI  
939 secretion system protein identifiers used as queries from *Ca. B. vickermanii*, and the protein  
940 identifiers from the *Holosporales* genomes that were significant hits.

Figure 1

bioRxiv preprint doi: <https://doi.org/10.1101/2025.10.03.678719>; this version posted October 3, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.







Figure 2



**Figure 3**




Figure 4

bioRxiv preprint doi: <https://doi.org/10.1101/2025.10.03.678719>; this version posted October 3, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.




Figure 5




Figure 6

bioRxiv preprint doi: <https://doi.org/10.1101/2025.10.03.678719>; this version posted October 3, 2025. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

