
GREATER TROCHANTERIC PAIN SYNDROME

Background

- Greater Trochanteric Pain Syndrome (GTPS) is a common condition that causes pain on the outside of the hip, often called "trochanteric bursitis." This pain mainly happens because of deterioration of or injury (tendinopathy) to the gluteus medius and gluteus minimus tendons. This injury or deterioration can irritate the surrounding tissues.
- GTPS is more commonly seen in women, adults over the age of 40, individuals with other hip or back pain, and those with poor running technique.
- Diagnosis is made through review of your medical history, physical exam, and, in some cases, imaging such as x-ray, MRI, or ultrasound.
- Treatment includes rest, avoiding painful activities, physical therapy, medication, injections, and, in some rare cases, surgery.



Rehab Instructions

Sit to Stand (Single leg squat with a chair)

1. Put a chair against a wall for stability.
2. Sit on the edge of the chair with one leg back and the leg other forward.
3. As you stand up, the leg that's staggered back will straighten first.
4. Once you're standing up, slowly lower yourself back onto the chair.
5. Keep your chest up and don't let your back arch or bend.
6. Use a chair or stool with a high seat height to make the exercise easier. Start with what's tolerable!

Complete 2 sets of 20 reps 1 x a day. Progress to 3 sets of 20 reps 1 x a day.





Bridges




1. Lie on your back with your knees bent and feet flat on the floor, keeping your hips shoulder-width apart.
2. Contract your glute muscles and lift your hips towards the ceiling.
3. Keep your back flat; do not overextend (arch) your back.
4. Hold for 15–20 seconds, then slowly return to the start position.

Complete 3 sets of 10 reps 1 x a day.

Wall Sits with Resistance Band (Abduction)




1. Step into a resistance band and move it up to around your thighs, just above your knees.
2. Stand with your feet shoulder-width apart and your back against a wall.
3. Bend your knees and hips to about 90 degrees to get into a wall-sit position. If needed, a shallower squat is acceptable.
4. Pull your knees apart against the resistance band.
5. Hold this position for 3–5 seconds.
6. Then slowly return your knees to the starting position, controlling the movement.

Complete 3 sets of 10 reps 1 x a day.

Single Knee to Chest Stretch



1. Lie on your back with your feet resting on the floor.
2. Pull one knee to your chest and hold for 10–15 seconds.
3. While keeping that same knee bent, pull it towards the opposite shoulder and hold for another 10–15 seconds.
4. You should feel this stretch in your lower back, glutes, and upper hamstring muscles.
5. Repeat the stretch on the other side.

Complete 3 sets for 20–30 seconds total on each side 1 x a day.

Practical Tips For Success

- Complete these exercises in a slow, controlled manner. Be consistent and perform them 6 days a week.
- A running gait analysis may be helpful for you to optimize your biomechanics and prevent injury.
- Excess body weight loads the gluteal tendons. Maintaining a healthy body weight may be helpful in reducing symptoms.

References

Kane, S. F., Olewinski, L. H., & Tamminga, K. S. (2019). Management of chronic tendon injuries. *American Family Physician, 100*(3), 147–157.

Ladurner, A., Fitzpatrick, J., & O'Donnell, J. M. (2021). Treatment of gluteal tendinopathy: A systematic review and stage-adjusted treatment recommendation. *Orthopaedic Journal of Sports Medicine, 9*(7). doi:10.1177/23259671211016850

Moore, D., Semciw, A. I., & Pizzari, T. (2020). A systematic review and meta-analysis of common therapeutic exercises that generate highest muscle activity in the gluteus medius and gluteus minimus segments. *International Journal of Sports Physical Therapy, 15*(6), 856–881. doi:10.26603/ijspt20200856

Photos used with permission in collaboration with the Musculoskeletal Injury Rehabilitation Research for Operational Readiness (MIRROR) program and the Geneva Foundation.