

Jan Engelmann

Tropical Cyclone Tracking with varying parameter thresholds

Semester Thesis

Institute for Atmospheric and Climate Science - Atmospheric Physics
Swiss Federal Institute of Technology (ETH) Zurich

Supervision

Bernhard Enz
Prof. Dr. Ulrike Lohmann

November 2020

Contents

Abstract	1
1 Introduction	2
1.1 Impact on society	2
1.2 Underlying Physics	2
1.3 Previous work on TC Tracking	4
2 Data and Methods	5
2.1 Simulation Data	5
2.2 Algorithm	5
2.2.1 TC candidate search	6
2.2.2 Creating TC tracks from previously found TC candidates	6
2.2.3 Varying parameter thresholds	7
2.3 Tracking Data Analysis	7
2.3.1 Algorithm output data format	7
3 Results	9
3.1 Filtering out noise	9
3.2 Validating Results	10
3.3 Variation of the Warm Core Criterion strength	10
3.4 Comparison of the Warm Core Criterion and the Vorticity Threshold	10
3.5 TC Genesis regions	11
4 Conclusion	15
A Additional TC Genesis Plots	17
Nomenclature	19

Abstract

An existing algorithm for tracking tropical cyclones in ICON output was sped up. The increased speed was used in order to evaluate assumptions on characterizing parameters for TCs. These parameters were varied and the results compared.

Chapter 1

Introduction

1. DONE Tcs are relevant for society due to their changing intensity and more people living on the coast
2. They exhibit interesting physics (tiefdrucksysteme mit Warmcore)
3. They are studied analytically, with field experiments and with simulations
4. Here their climatology is analyzed by analyzing simulation data

Tropical cyclones also known as Hurricanes or Typhoons are storms of extreme nature in many regards. Not only are they the most deadly and expensive natural catastrophes in the United States but also their physics is quite challenging with many open questions remaining.[1] Finally they are very likely to play a crucial role in the global heat balance and moisture circulation.[2][3]

1.1 Impact on society

While tropical cyclones form and intensify above the ocean, they have the largest impact on society during landfall. The damage happens due to a combination of strong winds and catastrophic storm surges. On average hurricanes inflict normalized damages of about \$10-billion/year in the United States.[4] A single strong storm can cause thousands of deaths. Hurricane Katrina e.g. in 2005 took the lives of over 1200 people.[5] Due to these enormous implications for society and opportunities to save lives and money, active research is happening on tropical cyclone impact reduction. With an unsure impact of climate change on tropical cyclone frequency and intensity and a trend of urbanization on the American east-coast, improving the understanding of tropical cyclones is of great importance.

1.2 Underlying Physics

Tropical Cyclones can be classified using the Saffir-Simpson wind scale. It is defined by the maximum wind observed in the cyclone. Due to their characteristic structure this wind usually occurs at the eyewall. This choice is motivated by the strong correlation between wind speeds and the inflicted damages.[6] The exact categorization can be seen in table 1.1.

Their occurrence frequency is displayed in Fig. 1.1

Tropical Cyclones are rotating low pressure systems that exhibit a warm core. They rotate around a pressure minimum with low wind speeds which is enclosed by the eyewall which in turn is the radius of maximum wind. The rotation around the center happens in different bands of updraft that are alternated with rain bands. Finally the the warm updraft in the center cools down on its way to the top. Resulting in a cold outflow at the top of the storm. This structure is depicted in Fig. 1.2. In order for a storm to develop a number of TC genesis criteria have to be met. Not all of

Tropical Cyclones		Other tropical low pressure systems	
category	wind speed [m/s]	name, category	wind speed [m/s]
1	33–42	tropical depression, -1	≤ 17
2	43–49	tropical storm, 0	18–32
3	50–58		
4	58–70		
5	≥ 70		

Table 1.1: Simpson scale defined by 1-minute maximum sustained winds [6]. The category number of the other tropical low pressure systems was assigned by the author and will be used in the results section.




Figure 1.1: Average cumulative number of storms in the Atlantic. Named systems are mostly tropical storms but can also be tropical depressions.[7]

them have to be satisfied but they do offer a good indicator for the probability of storm formation. The criteria are as follows:

- sea surface temperature above 26.5°C to at least a depth of 50m
- sufficiently moist mid-troposphere for deep convection
- Appreciable moisture flux at the ocean-air interface to sustain a conditionally unstable thermodynamic environment (Montgomery & Smith, 2017)
- A distance of at least 5° from the equator, so that the Coriolis effect is strong enough to initiate the cyclone's rotation (Palmén, 1948).
- A pre-existing weather disturbance with sufficient vorticity and convergence, e.g. a tropical easterly wave.
- Low vertical wind shear between the surface and the upper troposphere

[9]

1. DONE Classification between tc, tornado, tropical depression...
2. DONE Mention climatology and categorisation




Figure 1.2: Structure of a Tropical Cyclone in the Northern hemisphere [8]

3. DONE Storm characterisitics
4. Genesis criteria
5. Carnot Cycle
6. Shortly mentioning WISHE Paradigm
7. Mention 1 or 2 of Heini Wernlis Publications
8. use emanuel review paper and cloud dynamics course material

1.3 Previous work on TC Tracking

1. Mention work of Christoph Raible/Raibler from Bern regarding TC Tracking
2. See what Heini Wernli has done

Chapter 2

Data and Methods

The purpose of this project was to evaluate the parameter thresholds for tracking TCs in ICON simulation data. In the following section the settings used for generating the simulation data and the algorithm itself will be explained.

2.1 Simulation Data

ICON Model

The Icosahedral Nonhydrostatic Model (ICON) is developed by the German Weather Service (DWD), the Max Planck Institute for Meteorology and several partner institutes. As the name implies the grid is in a first step generated by mapping the earth's surface to an Icosahedron (platonic solid with 20 equilateral triangles as faces). The faces are then split up into smaller triangles in order to achieve the desired resolution. The model then delivers all typical meteorological quantities on this grid. The model uses the fully compressible and non-hydrostatic version of the Euler equations for the fundamental transport processes. Physical processes which cannot be resolved with the grid size are then parametrised by using complex functions that take into account the grid box mean values of the model variables. Physical processes which need to be treated in this way include solar and thermal radiation, cloud microphysics and turbulent transfer above the earth's surface.[10]

Analysed simulation output

The North Atlantic basin with its surrounding area (TODO xx1 - xx2 lon and yy1 - yy2 lat) was simulated from the beginning of May until the beginning of December. The output data is saved for every six hours. Two different kinds of time lagged ensembles were used. Both used ERA5 reanalysis data for the initial weather state and the monthly boundary conditions. The first however, labeled "ref", used the first day of each month at 6:00 a.m. (TODO !confirm with Bernhard) as the monthly boundary conditions. The second ensemble, labeled "rm", used monthly rolling mean boundary conditions based on the same reanalysis data. The members within each of these two ensembles were created by varying the initial weather state of the simulation. Namely each of the first 10 days of May were used as initial conditions for a separate simulation run. The output data from each run is mapped from the icosahedral to a Latitude-Longitude grid in order to facilitate distance calculations. Finally this results in 20 separate simulation runs that can be searched for TCs.

2.2 Algorithm

An existing algorithm developed by Bernhard Enz and inspired by [11] was improved in regards to runtime, robustness and readability. The gained speed was used in order to run the algorithm on

the same simulation data but with different threshold values that decide whether a TC was found or not. By comparing the different results reasonable thresholds and the importance of the different criteria were determined.

The algorithm consists of two steps. In the first step the TC candidates are found for each time step without taking into account if a storm already existed in the previous time-step. Afterwards all entries are analysed and connected to TC tracks if they lie close to each other temporally and spatially. Both steps will be explained in the following sections.

2.2.1 TC candidate search

TC candidates are found by searching the simulated domain using several criteria. They are summarised in Tab. 2.1 and will be explained in detail in the following sections.

criterion	parameters
sea level pressure minimum	slpdis
minimal vorticity threshold	vormin
warm core criterion	temdif, temdis

Table 2.1: Each criterion can be adjusted by changing its characterising parameter

Sea Level Pressure Minimum

As outlined in Sec. 1.2 TCs are low pressure systems. In fact some of the lowest pressures on earth were measured inside the eyes of TCs. Therefore the first step to finding TC candidates for a specific time-step is locating the sea level pressure minima. Here enormous speed up was achieved by replacing the previous manual minimum finding algorithm with a vectorised version from the image processing library scikit-image [12]. This function finds the local minima and requires them to be a certain distance apart. The stronger minimum is kept if two candidates are within the specified distance **slpdis**. The resulting minima are then further analysed.

Minimal Vorticity Threshold

If the vorticity at a pressure minimum is below the minimum threshold **vormin** it is discarded as a potential TC candidate.

Warm Core Criterion

The last qualifying characteristic is the warm core structure. The temperature at the height of 300 hPa in the pressure minimum is compared to the average temperature of the surrounding area at the same height. The form and strength of this warm core requirement can be adjusted by using the parameters **temdif** and **temdis**. The former corresponds to the required temperature difference between the center of the storm and its environment. Only storms that have a warm core that is at least **temdif** degrees warmer than the environment, are kept as valid TC candidates. The latter specifies the side-length of the square with the storm in the middle which is defined as environment.

Saving of TC information

At the end of each time-step the remaining candidates are saved with their corresponding date, time, position, maximum windspeed and sea level pressure in the center.

2.2.2 Creating TC tracks from previously found TC candidates

The TC tracks are created by checking the candidates for each time-step and comparing them to currently active TCs. At the beginning of each new time-step previously active TCs that were

interrupted are archived if they lived for at least 18 hours and deleted otherwise. Then for each entry it is checked which previous TC is within its maximum travel distance. This distance is set to 72 kilometres since a TC with a top speed of 20 m/s can travel this far within 6 hours. If several entries could have been reached by the same active TC, the entry with the lower sea level pressure is added to the track and the other entries are discarded. If no active TC within range of the entry is found a new active TC is created. Each TC is finally assigned a unique integer for identification.

2.2.3 Varying parameter thresholds

Naturally the results of the algorithm depend on the choice of the five parameters from Tab. 2.1. In order to answer the research question of this project the algorithm was run for different combinations of these parameters. For instance all combinations of the values in Tab. 2.2 were used.

parameter	unit	values
slpdis	m	25000, 50000, 100000
vormin	1/s	1e-6, 1e-5, 1e-4
temdif	K	0.5, 0.75, 1, 1.25
temdis	m	100000, 200000, 400000

Table 2.2: An exemplary list of parameters which results in 108 ($=3^3 \cdot 3^4 \cdot 3$) combinations

In summary, the algorithm is run with every parameter set on all of the 20 simulation runs.

2.3 Tracking Data Analysis

2.3.1 Algorithm output data format

The output of the algorithm is a pandas dataframe with one row per unique TC per timestep. Its columns are described in Tab. 2.3.

information	columns	description
timestamp	date	time and date of the TC entry
position	lon_idx, lat_idx, lon, lat	position of the found TC in longitude/latitude grid indices and coordinates
intensity	maxwind, curr_cat, cat	maximum wind within a 100km distance, snapshot category at the current time, maximum category of the corresponding TC
parameter combination	param_id	id of the parameter combination used for tracking
unique TC identifier	tc_id	unique TC identifier across all parameter combinations and simulation runs
simulation run	mem, exp	variables specifying the member and boundary conditions of the analysed simulation run
genesis sea level temperature	genesis_sst	genesis sea level temperature of the particular TC, identical across all time entries of a TC.

Table 2.3: description of the columns of the output pandas dataframe [13]

Chapter 3

Results

3.1 Filtering out noise

After the tracking and stitching steps are completed, the algorithm results can still be improved by filtering the results. A large improvement was achieved by requiring a certain sea surface temperature at TC genesis.

Sea Surface Temperature Criterion

As outlined in Sec. 1.2 the TCs need warm ocean water as an energy source, especially when they form. It has been shown that the large majority has a SST over 25.5°C [14]. Therefore it is expected that no reasonable TCs are filtered out when requiring a genesis SST of at least 24°C . However as can be seen in Fig. 3.1 a large part of the unwanted tracks in the North of the domain are removed.




Figure 3.1: Comparison of all tracks without and with the SST criterion on the left and right

Analysing geographically unreasonable tracks

Even with the application of the SST-criterion, unreasonable TC tracks remain. For instance the tracks over Wyoming shown in Fig. 3.2 should not be so frequent. To determine the parameters responsible for this, the 20 parameter combinations that account for the large majority of these, share the common feature that they all corresponded to the same weak warm core criterion. They had a **temdif** of 0.5°C and a **temdis** of 400 km. Therefore if only a very low temperature difference is required for an area that can be larger than smaller sized TCs, low pressure systems that do not

correspond to tropical cyclones are tracked. While this may not be surprising, it does emphasise the importance of a well trimmed warm core criterion.




Figure 3.2: Set of unreasonable tracks over Wyoming and the surrounding states

3.2 Validating Results

In Sec. 3.1 it was found that with a sufficiently strong warm core condition the tracks appear in reasonable areas. Before comparing the algorithm output for different parameter combinations, it still remains to be shown that the produced tracks actually follow TCs and not other low pressure systems. For that purpose ten different storms were randomly chosen. For these storms the radial, tangential and vertical wind and the sea level pressure were visualised in the azimuthal mean. It was found that all storms qualitatively exhibit the physically expected structure that was described in Sec. 1.2. The resulting plots for a representative storm can be seen in Fig. ??.

3.3 Variation of the Warm Core Criterion strength

With the aim of understanding the impact of different warm core criteria strengths, the resulting cyclone distributions for a range of different **temdifs** were compared. As can be seen in Fig. 3.4, a weaker warm core criterion leads to a distribution with more lower intensity storms. When comparing with the absolute counts it can be seen that this is the result from weaker storms being tracked for lower **temdifs**. Logically for TCs of category 2 and upwards, no difference in the counts is observed. Furthermore for **temdifs** larger than 1 K, no tropical depressions which correspond to category -1 are tracked. It can therefore be concluded that the warm core criterion can be used very efficiently to filter out noise if only strong TCs are of interest.

3.4 Comparison of the Warm Core Criterion and the Vorticity Threshold

In order to compare the importance of the warm core criterion with the vorticity threshold, TCs were tracked with parameter combinations of varying relative strength of the two criteria. To be concrete, all nine combinations of the parameters in Tab. 3.1 were used.

It was expected that the vorticity threshold should influence the distribution of tracked TCs only if a weak warm core criterion is applied. However, as can be seen in Fig. 3.5, even with a very weak

parameter	unit	values
slpdis	m	100000
vormin	1/s	1e-6, 1e-5, 1e-4
temdif	K	0.5, 1 ,1.5
temdis	m	200000

Table 3.1: Parameter combinations used for the comparison

warm core criterion does the vorticity threshold not influence the storm intensity distributions. A comparison for different values of **temdif** and an analysis of the impact of the vorticity criterion on the storm lifetime, can be found in the Appendix.

3.5 TC Genesis regions

A direct consequence of the observations from Sec. 3.4 is that with a weaker warm core criterion the TCs can be found already when they have not had much time to intensify. This is relevant since from best track data it is expected that most TCs form in the main development region (MDR) which lies roughly between 10 – 20 degree North and 20 – 80 degree West. However, when checking the genesis spots and density in Fig. 3.6, it can be seen that the MDR is not quite satisfied. When using a weaker warm core criterion




Figure 3.3: Azimuthal mean plots. From left to right and top to bottom: radial wind, tangential wind, vertical wind and sea level pressure.




Figure 4.4: Maximum TC category histograms for different temdif parameters. Each histogram on the left has unit norm. The right plot shows the absolute counts of TCs. The X-axis describes the TC categories as defined in Tab 1.1.




Figure 3.5: Snapshot category distribution for a weak warm core criterion and several vorticity thresholds




Figure 3.6: Genesis spots and density for a specific parameter combination.

Chapter 4

Conclusion

1. repeat most important findings
2. mention further algorithm ideas
3. suggest more elaborate parallelization with eg dask for improved simulation data reading

Appendix A

Additional TC Genesis Plots




Figure A.1: Genesis spots and density for a specific parameter combination.




Figure A.2: Genesis spots and density for a specific parameter combination.




Figure A.3: Genesis spots and density for a specific parameter combination.




Figure A.4: Genesis spots and density for a specific parameter combination.




Figure A.5: Genesis spots and density for a specific parameter combination.

Nomenclature

Symbols

EHC	Conditional equation	[−]
e	Willans coefficient	[−]
F, G	Parts of the system equation	[K/s]

Indicies

a	Ambient
air	Air

Acronyms and Abbreviations

ETH	Eidgenössische Technische Hochschule
TC	Tropical Cyclone
DWD	German Weather Service
MPIM	Max Planck Institute for Meteorology
ICON	Icosahedral Nonhydrostatic Model developed by the DWD and the MPIM
SST	sea surface temperature
SLP	sea level pressure
MDR	main development region

Bibliography

- [1] K. Emanuel, “Tropical cyclones,” *Annual Review of Earth and Planetary Sciences*, vol. 31, no. 1, pp. 75–104, 2003.
- [2] A. Jones, J. Haywood, N. Dunstone, K. Emanuel, M. Hawcroft, K. Hodges, and A. Jones, “Impacts of hemispheric solar geoengineering on tropical cyclone frequency,” *Nature Communications*, vol. 8, 12 2017.
- [3] P. Webster and G. Holland, “Role of hurricanes in the global heat balance,” *AGU Fall Meeting Abstracts*, 12 2006.
- [4] R. Pielke, J. Gratz, C. Landsea, D. Collins, M. Saunders, and R. Musulin, “Normalized hurricane damage in the united states: 1900–2005,” *Natural Hazards Review*, vol. 9, 02 2008.
- [5] I. Beven, John L., L. A. Avila, E. S. Blake, D. P. Brown, J. L. Franklin, R. D. Knabb, R. J. Pasch, J. R. Rhome, and S. R. Stewart, “Atlantic Hurricane Season of 2005,” *Monthly Weather Review*, vol. 136, pp. 1109–1173, 03 2008.
- [6] “Saffir-simpson hurricane wind scale.” <https://www.nhc.noaa.gov/aboutsshws.php>. (Accessed on 11/08/2020).
- [7] “Tropical cyclone climatology.” <https://www.nhc.noaa.gov/climo/>. (Accessed on 11/10/2020).
- [8] Kelvinsong, “File:hurricane-en.svg - wikimedia commons.” <https://commons.wikimedia.org/wiki/File:Hurricane-en.svg>. (Accessed on 11/10/2020).
- [9] U. Lohmann, F. Lüönd, and F. Mahrt, *Storms and cloud dynamics*, pp. 285–322. Cambridge University Press, 2016.
- [10] “Wetter und klima - deutscher wetterdienst - numerical weather prediction models.” https://www.dwd.de/EN/research/weatherforecasting/num_modelling/01_num_weather_prediction_modells/num_weather_prediction_models_node.html. (Accessed on 11/11/2020).
- [11] S. Kleppuk, V. Muccione, C. Raible, D. Bresch, P. Köllner-Heck, and T. Stocker, “Tropical cyclones in era40: A detection and tracking method,” *Geophysical Research Letters - GEOPHYS RES LETT*, vol. 35, 05 2008.
- [12] S. van der Walt, J. L. Schönberger, J. Nunez-Iglesias, F. Boulogne, J. D. Warner, N. Yager, E. Gouillart, T. Yu, and the scikit-image contributors, “scikit-image: image processing in Python,” *PeerJ*, vol. 2, p. e453, 6 2014.
- [13] T. pandas development team, “pandas-dev/pandas: Pandas,” Feb. 2020.
- [14] R. Dare and J. McBride, “The threshold sea surface temperature condition for tropical cyclogenesis,” *Journal of Climate - J CLIMATE*, vol. 24, pp. 4570–4576, 09 2011.



Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Institute for Atmospheric and Climate Science - Atmospheric Physics
Prof. Dr. U. Lohmann

Title of work:

Tropical Cyclone Tracking with varying parameter thresholds

Thesis type and date:

Semester Thesis, November 2020

Supervision:

Bernhard Enz
Prof. Dr. Ulrike Lohmann

Student:

Name:	Jan Engelmann
E-mail:	enjan@ethz.ch
Legi-Nr.:	15-936-784
Semester:	3

Statement regarding plagiarism:

By signing this statement, I affirm that I have read and signed the Declaration of Originality, independently produced this paper, and adhered to the general practice of source citation in this subject-area.

Declaration of Originality:

http://www.ethz.ch/faculty/exams/plagiarism/confirmation_en.pdf

Zurich, 16. 11. 2020: _____



Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Declaration of originality

The signed declaration of originality is a component of every semester paper, Bachelor's thesis, Master's thesis and any other degree paper undertaken during the course of studies, including the respective electronic versions.

Lecturers may also require a declaration of originality for other written papers compiled for their courses.

I hereby confirm that I am the sole author of the written work here enclosed and that I have compiled it in my own words. Parts excepted are corrections of form and content by the supervisor.

Title of work (in block letters):

Authored by (in block letters):

For papers written by groups the names of all authors are required.

Name(s):

First name(s):

With my signature I confirm that

- I have committed none of the forms of plagiarism described in the '[Citation etiquette](#)' information sheet.
- I have documented all methods, data and processes truthfully.
- I have not manipulated any data.
- I have mentioned all persons who were significant facilitators of the work.

I am aware that the work may be screened electronically for plagiarism.

Place, date

Signature(s)

For papers written by groups the names of all authors are required. Their signatures collectively guarantee the entire content of the written paper.