Naranjo Causality Assessment & Justification Report

Workora Pharmacovigilance Internship — Causality Assessment

Scoring is based only on information supplied in the fictional case pack. For Q1, a 'Known' expectedness reference was treated as evidence of previous reports; an 'Unexpected' reference was treated as unknown. Questions for which the case supplied no specific evidence were scored 0 rather than inferred.

Case	Drug	Reaction	Score	Category
SYN-PV-011	Metformin	Nausea	5	Probable
SYN-PV-012	Amlodipine	Peripheral oedema	5	Probable
SYN-PV-013	Ciprofloxacin	Tendon pain	5	Probable
SYN-PV-014	Sertraline	Dizziness	2	Possible
SYN-PV-015	Omeprazole	Headache	5	Probable
SYN-PV-016	Losartan	Fatigue	2	Possible
SYN-PV-017	Amoxicillin	Rash	5	Probable
SYN-PV-018	Atorvastatin	Myalgia	5	Probable
SYN-PV-019	Metformin	Nausea	3	Possible
SYN-PV-020	Amlodipine	Peripheral oedema	3	Possible
SYN-PV-021	Ciprofloxacin	Tendon pain	4	Possible
SYN-PV-022	Sertraline	Dizziness	5	Probable
SYN-PV-023	Omeprazole	Headache	2	Possible
SYN-PV-024	Losartan	Fatigue	5	Probable
SYN-PV-025	Amoxicillin	Rash	5	Probable
SYN-PV-026	Atorvastatin	Myalgia	5	Probable
SYN-PV-027	Metformin	Nausea	5	Probable
SYN-PV-028	Amlodipine	Peripheral oedema	2	Possible
SYN-PV-029	Ciprofloxacin	Tendon pain	5	Probable
SYN-PV-030	Sertraline	Dizziness	6	Probable

SYN-PV-011 — Metformin / Nausea

Age/Sex	23 / Unknown
Dose/Route	5 mg once daily / Oral
Indication	Hypertension
Onset	19 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Unknown / Positive
Concomitant medicines	None reported

Q	Score
1	1
2	2
3	0
4	2
5	0
6	0
7	0
8	0
9	0
10	0
Total	**5**
Category	**Probable**

Clinical rationale: The reaction (Nausea) occurred 19 day(s) after Metformin was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. Dechallenge information was unknown, so Q3 scored 0. Recurrence on re-administration strongly supports causality (+2). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 5, classified as Probable.

SYN-PV-012 — Amlodipine / Peripheral oedema

Age/Sex	24 / Female
Dose/Route	5 mg once daily / Oral
Indication	Hypertension
Onset	4 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Unknown / Positive
Concomitant medicines	None reported

Q	Score
1	1
2	2
3	0
4	2
5	0
6	0
7	0
8	0
9	0
10	0
Total	**5**
Category	**Probable**

Clinical rationale: The reaction (Peripheral oedema) occurred 4 day(s) after Amlodipine was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. Dechallenge information was unknown, so Q3 scored 0. Recurrence on re-administration strongly supports causality (+2). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 5, classified as Probable.

SYN-PV-013 — Ciprofloxacin / Tendon pain

Age/Sex	75 / Female
Dose/Route	500 mg twice daily / Oral
Indication	Diabetes
Onset	5 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Negative / Positive
Concomitant medicines	Vitamin D

Q	Score
1	1
2	2
3	0
4	2
5	0
6	0
7	0
8	0
9	0
10	0
Total	**5**
Category	**Probable**

Clinical rationale: The reaction (Tendon pain) occurred 5 day(s) after Ciprofloxacin was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. No improvement after withdrawal was reported, so Q3 scored 0. Recurrence on re-administration strongly supports causality (+2). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 5, classified as Probable.

SYN-PV-014 — Sertraline / Dizziness

Age/Sex	75 / Female
Dose/Route	5 mg once daily / Oral
Indication	Diabetes
Onset	9 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Negative / Negative
Concomitant medicines	Paracetamol

Q	Score
1	1
2	2
3	0
4	-1
5	0
6	0
7	0
8	0
9	0
10	0
Total	**2**
Category	**Possible**

Clinical rationale: The reaction (Dizziness) occurred 9 day(s) after Sertraline was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. No improvement after withdrawal was reported, so Q3 scored 0. No recurrence on re-administration reduces the probability (-1). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 2, classified as Possible.

SYN-PV-015 — Omeprazole / Headache

Age/Sex	33 / Male
Dose/Route	20 mg once daily / Oral
Indication	Hypertension
Onset	16 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Unknown / Positive
Concomitant medicines	None reported

Q	Score
1	1
2	2
3	0
4	2
5	0
6	0
7	0
8	0
9	0
10	0
Total	**5**
Category	**Probable**

Clinical rationale: The reaction (Headache) occurred 16 day(s) after Omeprazole was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. Dechallenge information was unknown, so Q3 scored 0. Recurrence on re-administration strongly supports causality (+2). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 5, classified as Probable.

SYN-PV-016 — Losartan / Fatigue

Age/Sex	66 / Unknown
Dose/Route	20 mg once daily / Oral
Indication	Diabetes
Onset	10 day(s) after treatment began
Expectedness	Unexpected
Dechallenge / Rechallenge	Positive / Negative
Concomitant medicines	None reported

Q	Score
1	0
2	2
3	1
4	-1
5	0
6	0
7	0
8	0
9	0
10	0
Total	**2**
Category	**Possible**

Clinical rationale: The reaction (Fatigue) occurred 10 day(s) after Losartan was started, supporting temporal association. The case pack identifies the reaction as unexpected; no prior-report evidence was supplied, so Q1 was scored 0. Improvement after withdrawal supports causality (+1). No recurrence on re-administration reduces the probability (-1). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 2, classified as Possible.

SYN-PV-017 — Amoxicillin / Rash

Age/Sex	57 / Unknown
Dose/Route	5 mg once daily / Oral
Indication	Hypertension
Onset	21 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Unknown / Positive
Concomitant medicines	Vitamin D

Q	Score
1	1
2	2
3	0
4	2
5	0
6	0
7	0
8	0
9	0
10	0
Total	**5**
Category	**Probable**

Clinical rationale: The reaction (Rash) occurred 21 day(s) after Amoxicillin was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. Dechallenge information was unknown, so Q3 scored 0. Recurrence on re-administration strongly supports causality (+2). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 5, classified as Probable.

SYN-PV-018 — Atorvastatin / Myalgia

Age/Sex	56 / Female
Dose/Route	500 mg twice daily / Oral
Indication	Anxiety
Onset	17 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Unknown / Positive
Concomitant medicines	None reported

Q	Score
1	1
2	2
3	0
4	2
5	0
6	0
7	0
8	0
9	0
10	0
Total	**5**
Category	**Probable**

Clinical rationale: The reaction (Myalgia) occurred 17 day(s) after Atorvastatin was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. Dechallenge information was unknown, so Q3 scored 0. Recurrence on re-administration strongly supports causality (+2). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 5, classified as Probable.

SYN-PV-019 — Metformin / Nausea

Age/Sex	29 / Unknown
Dose/Route	5 mg once daily / Oral
Indication	Anxiety
Onset	9 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Positive / Negative
Concomitant medicines	None reported

Q	Score
1	1
2	2
3	1
4	-1
5	0
6	0
7	0
8	0
9	0
10	0
Total	**3**
Category	**Possible**

Clinical rationale: The reaction (Nausea) occurred 9 day(s) after Metformin was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. Improvement after withdrawal supports causality (+1). No recurrence on re-administration reduces the probability (-1). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 3, classified as Possible.

SYN-PV-020 — Amlodipine / Peripheral oedema

Age/Sex	56 / Unknown
Dose/Route	5 mg once daily / Oral
Indication	Diabetes
Onset	7 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Unknown / Not done
Concomitant medicines	Paracetamol

Q	Score
1	1
2	2
3	0
4	0
5	0
6	0
7	0
8	0
9	0
10	0
Total	**3**
Category	**Possible**

Clinical rationale: The reaction (Peripheral oedema) occurred 7 day(s) after Amlodipine was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. Dechallenge information was unknown, so Q3 scored 0. Rechallenge was not performed/was not known, so Q4 scored 0. Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 3, classified as Possible.

SYN-PV-021 — Ciprofloxacin / Tendon pain

Age/Sex	33 / Unknown
Dose/Route	5 mg once daily / Oral
Indication	Infection
Onset	9 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Positive / Not done
Concomitant medicines	Paracetamol

Q	Score
1	1
2	2
3	1
4	0
5	0
6	0
7	0
8	0
9	0
10	0
Total	**4**
Category	**Possible**

Clinical rationale: The reaction (Tendon pain) occurred 9 day(s) after Ciprofloxacin was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. Improvement after withdrawal supports causality (+1). Rechallenge was not performed/was not known, so Q4 scored 0. Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 4, classified as Possible.

SYN-PV-022 — Sertraline / Dizziness

Age/Sex	43 / Female
Dose/Route	5 mg once daily / Oral
Indication	Infection
Onset	21 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Negative / Positive
Concomitant medicines	Paracetamol

Q	Score
1	1
2	2
3	0
4	2
5	0
6	0
7	0
8	0
9	0
10	0
Total	**5**
Category	**Probable**

Clinical rationale: The reaction (Dizziness) occurred 21 day(s) after Sertraline was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. No improvement after withdrawal was reported, so Q3 scored 0. Recurrence on re-administration strongly supports causality (+2). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 5, classified as Probable.

SYN-PV-023 — Omeprazole / Headache

Age/Sex	26 / Male
Dose/Route	500 mg twice daily / Oral
Indication	Anxiety
Onset	4 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Negative / Negative
Concomitant medicines	None reported

Q	Score
1	1
2	2
3	0
4	-1
5	0
6	0
7	0
8	0
9	0
10	0
Total	**2**
Category	**Possible**

Clinical rationale: The reaction (Headache) occurred 4 day(s) after Omeprazole was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. No improvement after withdrawal was reported, so Q3 scored 0. No recurrence on re-administration reduces the probability (-1). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 2, classified as Possible.

SYN-PV-024 — Losartan / Fatigue

Age/Sex	79 / Unknown
Dose/Route	500 mg twice daily / Oral
Indication	Hypertension
Onset	3 day(s) after treatment began
Expectedness	Unexpected
Dechallenge / Rechallenge	Positive / Positive
Concomitant medicines	Paracetamol

Q	Score
1	0
2	2
3	1
4	2
5	0
6	0
7	0
8	0
9	0
10	0
Total	**5**
Category	**Probable**

Clinical rationale: The reaction (Fatigue) occurred 3 day(s) after Losartan was started, supporting temporal association. The case pack identifies the reaction as unexpected; no prior-report evidence was supplied, so Q1 was scored 0. Improvement after withdrawal supports causality (+1). Recurrence on re-administration strongly supports causality (+2). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 5, classified as Probable.

SYN-PV-025 — Amoxicillin / Rash

Age/Sex	61 / Unknown
Dose/Route	500 mg twice daily / Oral
Indication	Infection
Onset	21 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Negative / Positive
Concomitant medicines	Paracetamol

Q	Score
1	1
2	2
3	0
4	2
5	0
6	0
7	0
8	0
9	0
10	0
Total	**5**
Category	**Probable**

Clinical rationale: The reaction (Rash) occurred 21 day(s) after Amoxicillin was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. No improvement after withdrawal was reported, so Q3 scored 0. Recurrence on re-administration strongly supports causality (+2). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 5, classified as Probable.

SYN-PV-026 — Atorvastatin / Myalgia

Age/Sex	48 / Unknown
Dose/Route	20 mg once daily / Oral
Indication	Hypertension
Onset	12 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Unknown / Positive
Concomitant medicines	Vitamin D

Q	Score
1	1
2	2
3	0
4	2
5	0
6	0
7	0
8	0
9	0
10	0
Total	**5**
Category	**Probable**

Clinical rationale: The reaction (Myalgia) occurred 12 day(s) after Atorvastatin was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. Dechallenge information was unknown, so Q3 scored 0. Recurrence on re-administration strongly supports causality (+2). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 5, classified as Probable.

SYN-PV-027 — Metformin / Nausea

Age/Sex	24 / Unknown
Dose/Route	20 mg once daily / Oral
Indication	Hypertension
Onset	14 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Unknown / Positive
Concomitant medicines	Paracetamol

Q	Score
1	1
2	2
3	0
4	2
5	0
6	0
7	0
8	0
9	0
10	0
Total	**5**
Category	**Probable**

Clinical rationale: The reaction (Nausea) occurred 14 day(s) after Metformin was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. Dechallenge information was unknown, so Q3 scored 0. Recurrence on re-administration strongly supports causality (+2). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 5, classified as Probable.

SYN-PV-028 — Amlodipine / Peripheral oedema

Age/Sex	77 / Unknown
Dose/Route	5 mg once daily / Oral
Indication	Hypertension
Onset	18 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Unknown / Negative
Concomitant medicines	Vitamin D

Q	Score
1	1
2	2
3	0
4	-1
5	0
6	0
7	0
8	0
9	0
10	0
Total	**2**
Category	**Possible**

Clinical rationale: The reaction (Peripheral oedema) occurred 18 day(s) after Amlodipine was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. Dechallenge information was unknown, so Q3 scored 0. No recurrence on re-administration reduces the probability (-1). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 2, classified as Possible.

SYN-PV-029 — Ciprofloxacin / Tendon pain

Age/Sex	35 / Unknown
Dose/Route	5 mg once daily / Oral
Indication	Diabetes
Onset	13 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Unknown / Positive
Concomitant medicines	Paracetamol

Q	Score
1	1
2	2
3	0
4	2
5	0
6	0
7	0
8	0
9	0
10	0
Total	**5**
Category	**Probable**

Clinical rationale: The reaction (Tendon pain) occurred 13 day(s) after Ciprofloxacin was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. Dechallenge information was unknown, so Q3 scored 0. Recurrence on re-administration strongly supports causality (+2). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 5, classified as Probable.

SYN-PV-030 — Sertraline / Dizziness

Age/Sex	82 / Male
Dose/Route	20 mg once daily / Oral
Indication	Anxiety
Onset	17 day(s) after treatment began
Expectedness	Known
Dechallenge / Rechallenge	Positive / Positive
Concomitant medicines	Paracetamol

Q	Score
1	1
2	2
3	1
4	2
5	0
6	0
7	0
8	0
9	0
10	0
Total	**6**
Category	**Probable**

Clinical rationale: The reaction (Dizziness) occurred 17 day(s) after Sertraline was started, supporting temporal association. The case pack identifies the reaction as known, so Q1 was scored +1. Improvement after withdrawal supports causality (+1). Recurrence on re-administration strongly supports causality (+2). Questions 5–10 were scored 0 because the supplied case does not provide the specific evidence required for those questions. Final Naranjo score = 6, classified as Probable.