

Repeated measure

Guy

Load required packages

```
library(RcmdrMisc)
```

Loading required package: car

Loading required package: carData

Loading required package: sandwich

```
library(ggplot2)
library(nlme)
library(emmeans)
```

Welcome to emmeans.

Caution: You lose important information if you filter this package's results.
See '? untidy'

Importing data

```
chap9data1 <- read.csv("chap9data1.csv", sep=",", header=T)
View(chap9data1)
```

Let us do the example in chapter 9 (“Biometry 721/821 Workbook(2024).pdf”) #Example 1

```
chap9data1$Food <- as.factor(chap9data1$Food)
chap9data1$Calf <- as.factor(chap9data1$Calf)
chap9data1$Month <- as.factor(chap9data1$Month)
```

```
#fit model
```

```
chap9data1.model1 <- lme(Weight~Month*Food,random=~1|Calf,data=chap9data1)
anova(chap9data1.model1)
```

	numDF	denDF	F-value	p-value
(Intercept)	1	24	465.2000	<.0001
Month	3	24	15.6277	<.0001
Food	1	8	1.4088	0.2693
Month:Food	3	24	0.8711	0.4697

```
chap9data1.model2 <- aov(Weight~Food*Month + Error(Calf),data=chap9data1)
summary(chap9data1.model2)
```

Error: Calf

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Food	1	409.6	409.6	1.409	0.269
Residuals	8	2326.0	290.8		

Error: Within

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Month	3	1036.9	345.6	15.628	7.58e-06 ***
Food:Month	3	57.8	19.3	0.871	0.47
Residuals	24	530.8	22.1		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Interpretation of the ANOVA Results

The analysis of variance (ANOVA) was conducted using a linear mixed-effects model (`lme`) to understand the effects of Month, Food, and their interaction (`Month:Food`) on the Weight of calves. Here is the interpretation of the ANOVA table generated:

1. Intercept:

- **F-value:** 465.2000
- **p-value:** <.0001
- This indicates that the intercept is highly significant. In mixed-effects models, the intercept represents the baseline level of the dependent variable (**Weight**) when all predictors are at their reference level. A highly significant intercept suggests that the mean **Weight** of the calves is significantly different from zero.

2. Month:

- **Degrees of Freedom (numDF):** 3
- **F-value:** 15.6277
- **p-value:** <.0001
- The variable **Month** has a highly significant effect on **Weight** (p-value < 0.0001). This suggests that the average **Weight** of the calves changes significantly across different months.

3. Food:

- **Degrees of Freedom (numDF):** 1
- **F-value:** 1.4088
- **p-value:** 0.2693
- The variable **Food** does not have a statistically significant effect on **Weight** (p-value = 0.2693). This indicates that, after accounting for other variables in the model, different types of food do not significantly affect the calves' weight.

4. Interaction Between Month and Food (Month:Food):

- **Degrees of Freedom (numDF):** 3
- **F-value:** 0.8711

- **p-value:** 0.4697
- The interaction between **Month** and **Food** is not statistically significant (p-value = 0.4697). This suggests that the effect of **Food** on **Weight** does not vary significantly across different months.

Conclusion:

- The results indicate that the calves' weight varies significantly over the months (**Month** is significant), but there is no significant effect of **Food** or the interaction between **Month** and **Food** on the weight. Therefore, month-to-month changes are the main driver of differences in weight, rather than the type of food or any combined effect of the month and food type. “ “ ”

Interpretation of the ANOVA Summary

The analysis of variance (ANOVA) was conducted using the **aov** function with an **Error** term for the random effect of **Calf**. This model evaluates the effects of **Food**, **Month**, and their interaction (**Food:Month**) on the **Weight** of calves while accounting for the random effect of **Calf**. Here is the interpretation of the ANOVA summary:

1. Error: Calf (Random Effect)

- This section represents the variance attributed to differences between calves (treated as a random effect).
- **Food:**
 - **Degrees of Freedom (Df):** 1
 - **Sum of Squares (Sum Sq):** 409.6
 - **Mean Square (Mean Sq):** 409.6
 - **F-value:** 1.409
 - **p-value (Pr(>F)):** 0.269
 - The effect of **Food** is not statistically significant (p-value = 0.269). This suggests that differences in the type of food do not significantly contribute to variations in weight among different calves.

2. Error: Within (Within-Groups Effects)

- This section shows the variance explained by the fixed effects (**Month** and **Food:Month**) within the groups (calves).
- **Month:**
 - **Degrees of Freedom (Df):** 3
 - **Sum of Squares (Sum Sq):** 1036.9
 - **Mean Square (Mean Sq):** 345.6
 - **F-value:** 15.628
 - **p-value (Pr(>F)):** 7.58e-06 (very significant, denoted by ***)
 - The effect of **Month** on **Weight** is highly significant (p-value < 0.0001). This indicates that the weight of calves changes significantly across different months.
- **Food:Month Interaction:**
 - **Degrees of Freedom (Df):** 3
 - **Sum of Squares (Sum Sq):** 57.8
 - **Mean Square (Mean Sq):** 19.3
 - **F-value:** 0.871
 - **p-value (Pr(>F)):** 0.47
 - The interaction between **Food** and **Month** is not statistically significant (p-value = 0.47). This suggests that the effect of **Food** on **Weight** does not vary significantly over different months.

3. Residuals

- Represents the unexplained variance within the groups after accounting for the fixed effects (**Month** and **Food:Month**).
- **Degrees of Freedom (Df):** 24
- **Sum of Squares (Sum Sq):** 530.8
- **Mean Square (Mean Sq):** 22.1

Conclusion

- The weight of calves varies significantly across different months, indicating a strong effect of Month.
- There is no significant effect of Food alone or the interaction between Food and Month on the weight of calves.
- The differences between individual calves (random effect of Calf) and other unexplained factors account for the remaining variance.

#Multiple comparisons

```
# Calculate estimated marginal means for the model
emmeans_result <- emmeans(chap9data1.model1, ~ Month)
```

NOTE: Results may be misleading due to involvement in interactions

```
# Display the estimated marginal means
summary(emmeans_result)
```

Month	emmean	SE	df	lower.CL	upper.CL
3	50.7	2.99	8	43.8	57.6
4	57.2	2.99	8	50.3	64.1
5	59.9	2.99	8	53.0	66.8
6	64.8	2.99	8	57.9	71.7

Results are averaged over the levels of: Food
Degrees-of-freedom method: containment
Confidence level used: 0.95

```
# Conduct pairwise comparisons
pairs(emmeans_result)
```

contrast	estimate	SE	df	t.ratio	p.value
Month3 - Month4	-6.5	2.1	24	-3.091	0.0241
Month3 - Month5	-9.2	2.1	24	-4.374	0.0011
Month3 - Month6	-14.1	2.1	24	-6.704	<.0001
Month4 - Month5	-2.7	2.1	24	-1.284	0.5817
Month4 - Month6	-7.6	2.1	24	-3.614	0.0071
Month5 - Month6	-4.9	2.1	24	-2.330	0.1192

Results are averaged over the levels of: Food
Degrees-of-freedom method: containment
P value adjustment: tukey method for comparing a family of 4 estimates

Interpretation of Estimated Marginal Means and Pairwise Comparisons

The analysis gives the estimated marginal means (EMMs) for different months and the pairwise comparisons between these months. Here is the interpretation:

1. Estimated Marginal Means (EMMs) for Month

- The table provides the estimated marginal means (`emmeans`) for each month along with their standard errors (`SE`), degrees of freedom (`df`), and confidence intervals (`lower.CL` and `upper.CL`).
- These EMMs represent the average weight for each month, adjusted for the levels of Food.

Example Values:

- **Month 3:** Estimated mean weight is 50.7 with a 95% confidence interval from 43.8 to 57.6.
- **Month 4:** Estimated mean weight is 57.2 with a 95% confidence interval from 50.3 to 64.1.
- **Month 5:** Estimated mean weight is 59.9 with a 95% confidence interval from 53.0 to 66.8.
- **Month 6:** Estimated mean weight is 64.8 with a 95% confidence interval from 57.9 to 71.7.

Note: There is a warning that results may be misleading due to involvement in interactions, so interpretation should consider the presence of significant interactions in the model.

2. Pairwise Comparisons Between Months

- Pairwise comparisons were performed between the different months to evaluate if there are significant differences in mean weights.

Contrast Estimates:

- **Month 3 vs. Month 4:** Difference in means is -6.5 (Month 3 lower), with a p-value of 0.0241 (significant).
- **Month 3 vs. Month 5:** Difference in means is -9.2 (Month 3 lower), with a p-value of 0.0011 (highly significant).
- **Month 3 vs. Month 6:** Difference in means is -14.1 (Month 3 lower), with a p-value < 0.0001 (very highly significant).

- **Month 4 vs. Month 5:** Difference in means is -2.7 (Month 4 lower), with a p-value of 0.5817 (not significant).
- **Month 4 vs. Month 6:** Difference in means is -7.6 (Month 4 lower), with a p-value of 0.0071 (significant).
- **Month 5 vs. Month 6:** Difference in means is -4.9 (Month 5 lower), with a p-value of 0.1192 (not significant).

Conclusion from Pairwise Comparisons:

- Significant differences in weight exist between several months, particularly between Month 3 and the other months (4, 5, and 6), indicating that weights vary significantly as time progresses.

3. P-Value Adjustment

- The p-values are adjusted using the Tukey method for multiple comparisons, ensuring that the family-wise error rate is controlled.

Conclusion

- The estimated marginal means reveal a general increase in weight from Month 3 to Month 6.
- Significant differences exist between several months, particularly Month 3 versus later months, suggesting time-related effects on weight.
- Results should be interpreted cautiously, considering potential interaction effects between **Month** and other variables such as **Food**.

Exercises

Do all the exercises in page 66 on document “Biometry 721/821 Workbook(2024).pdf”