

Table S2: Relative risks for the effects of alcohol use on disease outcomes from meta-analyses.

Disease outcome	Sex	Age (years)	I ^a	II	III	IV
Ischemic heart disease [1-3]	Both	30–44	1.00	0.60	0.62	1.00 ^b
		45–50	1.00	0.63	0.65	1.00 ^b
		60–69	1.00	0.82	0.83	1.00 ^b
		70–79	1.00	0.92	0.93	1.00 ^b
		≥80	1.00	0.97	0.98	1.00 ^b
Ischemic stroke [2-4]	Men	30–44	1.00	0.83 ^b	1.07 ^b	3.84
		45–50	1.00	0.88 ^b	1.05 ^b	2.52
		60–69	1.00	0.94 ^b	1.02 ^b	1.69
		70–79	1.00	0.97 ^b	1.01 ^b	1.32
		≥80	1.00	1.00 ^b	1.00 ^b	1.00 ^b
	Women	30–44	1.00	0.88 ^b	0.85 ^b	1.33 ^b
		45–50	1.00	0.91 ^b	0.89 ^b	1.22 ^b
		60–69	1.00	0.96 ^b	0.95 ^b	1.10 ^b
		70–79	1.00	0.98 ^b	0.98 ^b	1.05 ^b
		≥80	1.00	1.00 ^b	1.00 ^b	1.00 ^b
Hemorrhagic stroke [2-4]	Men	30–44	1.00	1.65 ^b	3.16 ^b	6.65
		45–50	1.00	1.42 ^b	2.22 ^b	3.60
		60–69	1.00	1.19 ^b	1.55 ^b	2.18
		70–79	1.00	1.09 ^b	1.26 ^b	1.55
		≥80	1.00	1.00 ^b	1.00 ^b	1.00 ^b
	Women	30–44	1.00	1.30 ^b	2.07 ^b	3.89 ^b
		45–50	1.00	1.20 ^b	1.67 ^b	2.54 ^b
		60–69	1.00	1.09 ^b	1.30 ^b	1.70 ^b
		70–79	1.00	1.04 ^b	1.14 ^b	1.32 ^b
		≥80	1.00	1.00 ^b	1.00 ^b	1.00 ^b
Hypertensive disease [2,5]	Men	30≥	1.00	1.40	2.00	4.10
	Women	30≥	1.00	1.40	2.00	2.00
Cardiac arrhythmias [2,6]	Both	30≥	1.00	1.51	2.23	2.23
Larynx cancer [2,6]	Both	30≥	1.00	1.83	3.90	4.93
Mouth and pharynx cancer [2]	Both	30≥	1.00	1.45	1.85	5.39
Selected other cancers [3]	Both	30≥	1.00	1.10	1.30	1.70
Diabetes mellitus [2]	Men	30≥	1.00	0.99	0.57	0.73
	Women	30≥	1.00	0.92	0.87	1.13

Disease outcome	Sex	Age (years)	I ^a	II	III	IV
Liver cirrhosis [3,6]	Both	30≥	1.00	1.30	9.50	13.00
Pancreatitis [2,5]	Men	30≥	1.00	1.30	1.80	3.20
	Women	30≥	1.00	1.30	1.80	1.80

I, Abstainers; II, <40 g/day of ethanol intake; III, 40–59.9 g/day; IV, ≥60 g/day.

^a Reference category

^b We replaced these statistically insignificant relative risks with 1 in our analysis.

References

1. Corrao G, Rubbiati L, Bagnardi V, Zambon A, Poikolainen K (2000) Alcohol and coronary heart disease: a meta-analysis. *Addiction* 95: 1505-1523.
2. Danaei G, Ding EL, Mozaffarian D, Taylor B, Rehm J, et al. (2009) The preventable causes of death in the United States: comparative risk assessment of dietary, lifestyle, and metabolic risk factors. *PLoS Med* 6: e1000058.
3. Rehm J, Room R, Monteiro M, Gmel G, Kathryn G, et al. (2004) Alcohol Use. In: Ezzati M, Lopez AD, Rodgers A, Murray CJ, editors. *Comparative Quantification of health risks: Global and regional burden of disease attributable to selected major risk factors*. Geneva: WHO.
4. Reynolds K, Lewis B, Nolen JD, Kinney GL, Sathya B, et al. (2003) Alcohol consumption and risk of stroke: a meta-analysis. *JAMA* 289: 579-588.
5. Corrao G, Bagnardi V, Zambon A, Arico S (1999) Exploring the dose-response relationship between alcohol consumption and the risk of several alcohol-related conditions: a meta-analysis. *Addiction* 94: 1551-1573.
6. Gutjahr E, Gmel G, Rehm J (2001) Relation between average alcohol consumption and disease: an overview. *Eur Addict Res* 7: 117-127.