

S1 Table

Variable	Tumor sample	Plasma/ serum	RNA
Ang-2	Zhang L., 2003 [1]: 52 specimen Result: Expression in endothelium and 12% of tumor cells	Sallinen H., 2010 [5]: 150 patients Result: Higher pre-operative Ang-2 levels in cancer. Poor PFS and OS.	Hata K., 2004 [9]: 85 specimens Result: Low Ang-1/ Ang-2 predicted poor OS (univariate analysis)
	Lin Z., 2011 [2]: 21 patients Result: No correlation with clinicopathological parameters, no difference between cancer vs benign	Sallinen H., 2014 [6]: 75 patients Result: Higher Ang-2 and Ang-2/ VEGFR2 in cancer. Poor PFS and OS.	
Tie-2/ Ang-2	Li S., 2010 [3]: 12 specimens Result: Cytoplasmic expression, higher expression in cancer, no correlation with clinical parameters	Backen A., Clin Cancer Res 2014 [7]: 91 patients Result: High Ang-1/Tie-2 improved PFS in bevacizumab treated patients	Hata K., Oncology 2002 [10]: 19 patients Result: No difference with Ang-2 and Tie-2 expression in cancer vs benign tumors
	Brunckhorst M.K., 2014 [4]: 12 cancer and 12 omental metastatic samples Result: Ang-2 expression in tumor cells, Tie-2 in endothelium and fibroblasts. Higher Ang-2 expression in cancer.	Zhou C., 2016 [8]: 92 patients Result: 50% increase of Tie-2 from the nadir predicted tumor progression after bevacizumab	

S2 Table

Variable	Correlation coefficient (r)	<i>p</i>
Ang-2 IRS	-0.64	0.002
Ang-2 IRS*	-0.79	0.004
Ang-2 IRS (HGSC)	-0.87	<0.001
Ang-2 met PP	0.75	0.005
Tie-1 IRS	0.008	0.972
Tie-1 IRS (* /HGSC)	-0.087	0.799
Tie-1 met IRS	0.215	0.502
Tie-2 IRS	0.163	0.479
Tie-2 IRS (* /HGSC)	-0.319	0.339
Tie-2 met IRS	0.211	0.510

p-values determined by Spearman's test, IRS immunoreactive score, HGSC high grade serous carcinoma, Ang-2 met angiopoietin-2 metastasis, PP percentage of positively stained epithelial cells, * serous histologic type

S3 Table

	Ang2	<i>p</i>	Tie1	<i>p</i>	Tie2	<i>p</i>
Grade						
<i>Low</i>						
IRS	8		3		2	
PP	80		65		40	
<i>High</i>						
IRS	6	0.075	2	0.018	2	0.108
PP	65	0.008	25	0.007	30	0.016
Stage						
<i>I-II</i>						
IRS	9		2		2	
PP	75		20		40	
<i>III-IV</i>						
IRS	6	0.185 (0.047*)	2	0.781	2	0.141
PP	65	0.370	35	0.657	35	0.375
Histology						
<i>Serous</i>						
IRS	6	0.436	2	0.002	2	0.002
PP	65	0.110	40	0.002	30	<0.001
<i>Mucinous</i>						
IRS	6	0.867	3	0.029	6	0.001
PP	75	0.255	60	0.024	75	<0.001
<i>Endometrioid</i>						
IRS	6	0.175	2	0.411	2	0.672
PP	65	0.878	20	0.136	33	0.890
<i>Clear cell</i>						
IRS	4	0.477	1	0.001	1	0.095
PP	30	0.020	5	0.003	10	0.024
Residual tumor						
<i>None</i>						
IRS	6		2.5		3	
PP	70		45		40	
<i>≤1cm</i>						
IRS	6	0.109	2.5	0.583	2	0.923
PP	65	0.746	45	0.496	30	0.854
<i>>1cm</i>						
IRS	6	0.012	2	0.465 (0.008*)	2	0.688
PP	60	0.092	25	0.284 (0.020*)	30	0.526
Recurrence						
<i>No</i>						
IRS	6		3		3	

PP	73		45		42.5
<i>Yes</i>					
IRS	6	0.018	2	0.157 (0.025*)	2 0.202
PP	63	0.006	25	0.091 (0.009*)	30 0.137

p values determined by Kruskal-Wallis test followed by Mann-Witney tests when appropriate; *p value reported for serous tumors in brackets when only comparison for this subgroup was statistically significant; IRS, PP are median values; HGSC = high grade serous cancer, IRS immunoreactive score, PP percentage of positively stained cells, Recurrence = recurrence of cancer

References

1. Zhang L, Yang N, Park J, Katsaros D, Fracchioli S, Cao G, et al. Tumor-derived Vascular Endothelial Growth Factor Up-Regulates Angiopoietin-2 in Host Endothelium and Destabilizes Host Vasculature, Supporting Angiogenesis in Ovarian Cancer. *Cancer Research*. 2003;63: 3403-3412.
2. Lin Z, Liu Y, Sun Y, He X. Expression of Ets-1, Ang-2 and maspin in ovarian cancer and their role in tumor angiogenesis. *Journal of experimental & clinical cancer research : CR*. 2011;30: 31. doi: 10.1186/1756-9966-30-31.
3. Li S, Meng L, Zhu C, Wu L, Bai X, Wei J, et al. The universal overexpression of a cancer testis antigen hiwi is associated with cancer angiogenesis. *Oncology reports*. 2010;23: 1063. doi: 10.3892/or_00000733.
4. Brunkhorst MK, Xu Y, Lu R, Yu Q. Angiopoietins Promote Ovarian Cancer Progression by Establishing a Procarcinoma Microenvironment. *American Journal of Pathology, The*. 2014;184: 2285-2296. doi: 10.1016/j.ajpath.2014.05.006.
5. Sallinen H, Heikura T, Laidinen S, Kosma VM, Heinonen S, Ylä-Herttuala S, et al. Preoperative Angiopoietin-2 Serum Levels: A Marker of Malignant Potential in Ovarian Neoplasms and Poor Prognosis in Epithelial Ovarian Cancer. *International Journal of Gynecological Cancer*. 2010;20: 1498-1505. doi: 10.1111/IGC.0b013e3181f936e3.
6. Sallinen H, Heikura T, Koponen J, Kosma VM, Heinonen S, Ylä-Herttuala S, et al. Serum angiopoietin-2 and soluble VEGFR-2 levels predict malignancy of ovarian neoplasm and poor prognosis in epithelial ovarian cancer. *BMC Cancer*. 2014;14: 696. doi: 10.1186/1471-2407-14-696.
7. Backen A, Renehan AG, Clamp AR, Berzuini C, Zhou C, Oza A, et al. The combination of circulating Ang-1 and Tie-2 levels predicts progression-free survival advantage in bevacizumab treated patients with ovarian cancer. *Clin Cancer Res*. 2014.
8. Zhou C, Clamp A, Backen A, Berzuini C, Renehan A, Banks RE, et al. Systematic analysis of circulating soluble angiogenesis-associated proteins in ICON7 identifies Tie2 as a biomarker of vascular progression on bevacizumab. *The British Journal of Cancer*. 2016;115: 228-235. doi: 10.1038/bjc.2016.194.
9. Hata K, Nakayama K, Fujiwaki R, Katabuchi H, Okamura H, Miyazaki K. Expression of the angiopoietin-1, angiopoietin-2, Tie2, and vascular endothelial growth factor gene in epithelial ovarian cancer. *Gynecologic Oncology*. 2004;93: 215-222. doi: 10.1016/j.ygyno.2003.12.031.
10. Hata K, Udagawa J, Fujiwaki R, Nakayama K, Otani H, Miyazaki K. Expression of Angiopoietin-1, Angiopoietin-2, and Tie2 Genes in Normal Ovary with Corpus luteum and in Ovarian Cancer. *Oncology*. 2002;62: 340-348. doi: 10.1159/000065066.