

Supplementary Table 1. Difference in baseline characteristics of patients accepting participation in the follow up study and those rejecting participation

	Accepted Participation	Rejected Participation	p
n (%)	46 (63.9%)	26 (36.1%)	-
Age (year)	63.8 (±6.5)	61.5 (±8.9)	0.20
Sex (male)	37 (80.4%)	19 (73.1%)	0.52
AHI (events/hour)	37.8 (±16.0)	30.9 (±13.0)	0.07
Diabetes duration (year)	16.1 (±9.3)	14.6 (±7.7)	0.50
Systolic BP (mmHg)	143.9 (±15.3)	138.6 (±17.5)	0.19
Diastolic BP (mmHg)	83.3 (±7.8)	80.4 (±9.8)	0.17
cfPWV (m/s)	10.7 (±2.0)	11.1 (±2.2)	0.50
BMI (kg/m ²)	34.7 (±3.9)	34.3 (±5.1)	0.69
HbA1c (mmol/mol)	66.7 (±8.9)	66.3 (±3.9)	0.84
Fasting blood glucose (mmol/l)	9.4 (±2.6)	9.8 (±1.9)	0.43
Smoking			
- Currently (n (%))	6 (13%)	4 (15.4%)	0.92
- Former (n (%))	18 (39.1%)	9 (34.6%)	
- Never (n (%))	22 (47.8%)	13 (50%)	
Antidiabetic medication			
- No medication (n (%))	3 (6.5%)	0 (0%)	0.18
- Metformin (n (%))	26 (78.3%)	22 (84.6%)	0.51
- Insulin (n (%))	29 (63.0%)	18 (69.2%)	0.60
- GLP1 analog (n (%))	27 (58.7%)	12 (46.2%)	0.31
- DPP4 inhibitor (n (%))	3 (6.5%)	4 (15.4%)	0.22
- SGLT2 inhibitor (n (%))	5 (10.9%)	8 (30.8%)	0.04
- Sulphonyl Urea (n (%))	1 (2.2%)	1 (3.8%)	0.68
Blood pressure medication			
- No medication (n (%))	4 (8.7%)	2 (4.3%)	0.88
- Beta blockers (n (%))	18 (39.1%)	10 (38.5%)	0.96
- Thiazide (n (%))	19 (41.3%)	9 (34.6%)	0.58
- ACE-inhibitor (n (%))	24 (52.2%)	11 (42.3%)	0.42
- AT2 antagonist (n (%))	14 (30.4%)	10 (38.5%)	0.49
- Calcium blockers (n (%))	18 (39.1%)	11 (42.3%)	0.79
- Loop diuretics (n (%))	8 (17.4%)	2 (4.3%)	0.25
- Other (n (%))	7 (15.2%)	2 (4.3%)	0.35
Statin use (n(%))	36 (78.3%)	21 (80.8%)	0.56
Asprin use (n(%))	25 (54.3%)	12 (46.2%)	0.61

Abbreviations: BP, blood pressure; cfPWV, carotid-femoral pulse wave velocity; GLP1, glucagon like peptide; DPP4, di-peptidyl peptidase 4; SGLT2, sodium-glucose co-transporter 2; ACE, angiotensin converting enzyme; AT2, angiotensin-

