

Supplemental Table 1. Pneumococcal serotype prevalence by vaccine valency and collection year

Serotypes		Total (n=264)	2016 (n=51)	2017 (n=92)	2018 (n=93)	2019 (n=28)
PCV-13		72 (27%)	17 (33%)	24 (26%)	24 (26%)	7 (25%)
	1	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
	3	5 (2%)	1 (2%)	1 (1%)	3 (3%)	0 (0%)
	4	1 (1%)	0 (0%)	1 (1%)	0 (0%)	0 (0%)
	5	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
	6A	10 (4%)	5 (10%)	1 (1%)	3 (3%)	1 (4%)
	6B	2 (1%)	0 (0%)	1 (1%)	1 (1%)	0 (0%)
	6C*	3 (1%)	0 (0%)	1 (1%)	2 (2%)	0 (0%)
	7F	1 (1%)	1 (1%)	0 (0%)	0 (0%)	0 (0%)
	9V	2 (1%)	2 (4%)	0 (0%)	0 (0%)	0 (0%)
	14	2 (1%)	0 (0%)	2 (2%)	0 (0%)	0 (0%)
	18C	3 (1%)	1 (2%)	1 (1%)	0 (0%)	1 (4%)
	19A	16 (6%)	4 (8%)	8 (9%)	2 (2%)	2 (7%)
	19F	20 (8%)	3 (6%)	5 (5%)	9 (10%)	3 (11%)
	23F	7 (3%)	0 (0%)	3 (3%)	4 (4%)	0 (0%)
Additional PCV-15		3 (1%)	0 (0%)	0 (0%)	3 (3%)	0 (0%)
	22F	1 (1%)	0 (0%)	0 (0%)	1 (1%)	0 (0%)
	33F	2 (1%)	0 (0%)	0 (0%)	2 (2%)	0 (0%)
Additional PCV-20		37 (14%)	10 (20%)	14 (15%)	9 (10%)	4 (14%)
	11A	14 (5%)	4 (8%)	6 (7%)	2 (2%)	2 (7%)
	15B	15 (6%)	4 (8%)	4 (4%)	6 (6%)	1 (4%)
	Other	8 (3%)	2 (4%)	4 (4%)	1 (1%)	1 (4%)
Non-vaccine		152 (58%)	24 (47%)	57 (60%)	61 (62%)	17 (61%)
	7C	10 (4%)	2 (4%)	4 (4%)	2 (2%)	2 (7%)
	9N	5 (2%)	1 (2%)	1 (1%)	2 (2%)	1 (4%)
	15A	9 (3%)	1 (2%)	5 (5%)	2 (2%)	1 (4%)
	15C	6 (2%)	0 (0%)	4 (4%)	1 (1%)	1 (4%)
	16F	11 (4%)	1 (2%)	5 (5%)	4 (4%)	1 (4%)
	17F	7 (3%)	2 (4%)	1 (1%)	2 (2%)	2 (7%)
	21	12 (5%)	0 (0%)	4 (4%)	6 (6%)	2 (7%)
	23A	7 (3%)	2 (4%)	1 (1%)	2 (2%)	2 (7%)
	23B	29 (11%)	3 (6%)	12 (13%)	14 (15%)	0 (0%)
	34	8 (3%)	2 (4%)	2 (2%)	2 (2%)	2 (7%)
	35B	10 (4%)	2 (4%)	2 (2%)	4 (4%)	2 (7%)
	35F	9 (3%)	3 (6%)	2 (2%)	3 (3%)	1 (4%)
	Other	29 (11%)	5 (10%)	11 (12%)	13 (14%)	0 (0%)

*Serotype 6C is included as a PCV-13 serotype due to cross-protection provided by serotype 6A