

Supplementary File 1:

Table of clinical details of spontaneous preterm births within 7 days of TPTL presentation which were not appropriately managed during EQUIPTT trial

Case	Site	QUIPP risk	Delivery gestation (weeks ^{days})	Clinical Situation/impact	Neonatal
1	Intervention	0.2%	30 ⁺²	No steroids or admission- but re-attended and received timely ACS	NNU admission
2	Intervention	0.5%	33 ⁺⁰	Not admitted but re-attended and received timely ACS	NNU admission
3	Intervention	0.6%	33 ⁺³	Not admitted but re-attended and received timely ACS	NNU admission
4	Intervention	16.4%	31 ⁺⁴	Admitted but ACS previously (not within 7 days)	NNU admission
5	Control		34 ⁺⁴	Not admitted, ACS previously (not within 7 days),	NNU + sepsis
6	Control		25 ⁺²	No ACS, twins-	Ex-utero transfer
7	Control		29 ⁺³	No ACS, reattended and received magnesium sulphate	Neonatal death
8	Control		32 ⁺³	No ACS or admission	Nil
9	Control		33 ⁺⁰	No ACS, Delivered same day	NNU admission
10	Control		35 ⁺³	No admission or ACS	Nil
11	Control		34 ⁺⁶	No admission but received ACS	NNU admission
12	Control		34 ⁺²	No steroids, Delivered same day	NNU admission
13	Control		33 ⁺¹	Twins, Admitted 2/3 episodes, ACS not within 7 days	NNU- both twins RDS
14	Control		33 ⁺⁵	Admitted but no ACS	NNU admission, left hypoplastic heart syndrome
15	Control		28 ⁺¹	Admitted but ACS previously (not within 7 days)	NNU admission, RDS

16

Control

27⁺⁴

Admitted but ACS previously (not NNU admission, RDS within 7 days)