

Table 1:

Change in mean heart rate (beats per minute) over time between two groups:

GROUP	N	MEAN	S.D	S.E.M	P value
Group A	30	77.600	9.5217	1.7384166	0.003
Group B	30	68.200	9.3307	1.7035450	

There was statistically significant decrease in mean heart rate in Group B compared to Group A ('p' value = 0.0003 [<0.05]). (Table 1)

Table 2:

Change in mean of mean blood pressure (mm of Hg) over time between two groups:

GROUP	N	MEAN	S.D	S.E.M	P value
Group A	30	84.700	10.6387	1.9423520	0.0078
Group B	30	78.300	6.9736	1.2730898	

There was statistically significant decrease in mean of mean blood pressure in Group B compared to Group A ('p' value = 0.0078 [<0.05]). (Table 2)

Table 3:

Change in Spo₂ (%) over time between two groups:

GROUP	N	MEAN	S.D	S.E.M	P value
Group A	30	99.300	1.0222	0.1866273	1.000
Group B	30	99.300	1.2077	0.2204948	

There was no statistically significant changes in mean Spo₂ in Group B compared to Group A ('p' value = 1.000 [>0.05]). (Table 3)

Table 4:

Comparison of duration of sensory blockade (minutes) between two groups:

GROUP	N	MEAN	S.D	S.E.M	P value
Group A	30	351.500	63.4381	11.5821595	0.6936
Group B	30	358.500	73.1690	13.3587706	

There was no statistically significant difference in duration of sensory blockade between two groups. ('p' value = 0.6936 [>0.05]). (Table 4)

Table 5:

Comparison of duration of motor blockade (minutes) between two groups:

GROUP	N	MEAN	S.D	S.E.M	P value
Group A	30	305.500	62.0532	11.3293125	0.1044
Group B	30	336.500	82.1285	14.9944527	

There was no statistically significant difference in duration of motor blockade between two groups. ('p' value = 0.1044 [>0.05]). (Table 5)

Table 6:

Comparison of duration of post-operative analgesia (minutes) between two groups:

GROUP	N	MEAN	S.D	S.E.M	P value
Group A	30	459.000	68.7299	12.5481412	0.0038
Group B	30	515.500	75.9917	13.8741228	

There was statistically significant prolongation of post-operative analgesia in group B compared to group A. ('p' value = 0.0038 [<0.05]). (Table 6)

Table 7:

Comparison of total ephedrine consumption (milligrams) between two groups:

GROUP	N	MEAN	S.D	S.E.M	P value
-------	---	------	-----	-------	---------

Group A	30	4.800	5.3201	0.9713129	0.0069
Group B	30	9.000	6.2533	1.1416912	

There was statistically significant increase in total ephedrine consumption in group B compared to group A. ('p' value = 0.0069 [<0.05]). (Table 7)

Table 8:

Comparison of total blood loss (milliliter) between two groups:

GROUP	N	MEAN	S.D	S.E.M	P value
Group A	30	354.000	144.6625	26.29154165	0.002
Group B	30	180.500	72.3229	13.15429458	

There was statistically significant decrease in blood loss in group B compared to group A. ('p' value = 0.002 [<0.05]). (Table 8)