

~~Bread components derived from yeast fermented wheat and rye breads increase colitis and mortality in a mouse model of colitis~~

Julia Zimmermann, Luigia De Fazio, Valentina Kaden-Volynets, Bernd Hitzmann, Stephan C. Bischoff*

*Corresponding author: bischoff.stephan@uni-hohenheim.de; Department of Nutritional Medicine/Prevention, University of Hohenheim, Fruwirthstrasse 12, Stuttgart, 70599 Germany

State Institution Alternative (SIA) Commissioner's Report

06/13/2025

State Hospital Wait List

Total SIA Persons served

- | | |
|-------------------------------------|-----------------|
| • OSH – 22 as of 8:42am on 06/13/25 | • 2855 adults |
| • LSH – 5 as of 8:24am on 06/13/25 | • 1572 children |

SIA Adult Census

Cottonwood Springs

Via Christi (adult)

***16 open SIA bed as of 12:43pm 06/13/25

***1 open SIA bed as of 8:27am 06/13/2025

New Admits 06/06/25 to 06/12/25

New Admits 06/06/25 to 06/12/25

- | | |
|-----------------------------|-----------------------------|
| • 12 new SIA admits | • 3 new SIA admits |
| ○ 8 paid by HIS | • 1 paid by HIS |
| ○ 4 Private Insurance | • 2 Private Insurance |
| • 0 Admitted and discharged | • 0 Admitted and discharged |
| • 11 OSH Catchment Area | • 3 OSH Catchment Area |
| • 1 LSH Catchment Area | • 0 LSH Catchment Area |

Current Clients Receiving Services

Current Clients Receiving Services

06/06/25 to 06/12/25

06/06/25 to 06/12/25

- | | |
|-------------------------|------------------------|
| • 28 SIA admits | • 6 SIA admits |
| ○ 15 paid by HIS | • 1 paid by HIS |
| ○ 13 Private Insurance | • 5 Private Insurance |
| • 26 OSH Catchment Area | • 6 OSH Catchment Area |
| • 2 LSH Catchment Area | • 0 LSH Catchment Area |

Cumulative Totals 08/30/21 – 06/12/25
06/12/25

Cumulative Totals 09/27/21 –

- | | |
|---------------------------------|--------------------------------|
| • 1448 Total discharges | • 539 total discharges |
| • 1476 Total SIA clients served | • 545 Total SIA clients served |
| ○ 1268 OSH catchment area | • 516 OSH catchment |
| ○ 208 LSH catchment area | • 29 LSH catchment |

~~Bread components derived from yeast fermented wheat and rye breads increase colitis and mortality in a mouse model of colitis~~

Julia Zimmermann, Luigia De Fazio, Valentina Kaden-Volynets, Bernd Hitzmann, Stephan C. Bischoff*

*Corresponding author: bischoff.stephan@uni-hohenheim.de; Department of Nutritional Medicine/Prevention, University of Hohenheim, Fruwirthstrasse 12, Stuttgart, 70599 Germany

<u>Newton Medical Center</u>	<u>ANEW Health</u>
<u>***0 open SIA bed as of 3:50pm 06/13/25</u>	<u>***2 open SIA bed as of 7:30am 06/12/25</u>
<u>New Admits 06/06/25 to 06/12/25</u>	<u>New Admits 06/06/25 to 06/12/25</u>
• 0 new SIA admits ○ 0 paid by HIS ○ 0 Private Insurance • 0 Admitted and discharged • 0 OSH Catchment Area • 0 LSH Catchment Area	●0 new SIA admits ●0 paid by HIS ●0 Private Insurance ●0 Admitted and discharged ●0 OSH Catchment Area ●0 LSH Catchment Area
<u>Current Clients Receiving Services</u>	<u>Current Clients Receiving Services</u>
<u>06/06/25 to 06/12/25</u>	<u>06/06/25 to 06/12/25</u>
• 0 SIA admits ○ 0 paid by HIS ○ 0 Private Insurance • 0 OSH Catchment Area • 0 LSH Catchment Area	●2 SIA admits ●2 paid by HIS ●0 Private Insurance ●2 OSH Catchment Area ●0 LSH Catchment Area
<u>Cumulative Totals 09/27/21 – 06/12/25</u>	<u>Cumulative Totals 09/27/21 – 06/12/25</u>
• 32 Total discharges • 32 Total SIA client served ○ 8 OSH catchment area ○ 24 LSH catchment area	●273 total discharges ●275 Total SIA clients served ●241 OSH catchment ●34 LSH catchment
<u>Hutchinson Regional</u>	<u>Children's Mercy Camber</u>
<u>***4 open SIA bed as of 4:08pm 06/13/25</u>	<u>***6 open SIA bed as of 9:02am 06/13/25</u>
<u>New Admits 06/06/25 to 06/12/25</u>	<u>New Admits 06/06/25 to 06/12/25</u>
• 0 new SIA admits ○ 0 paid by HIS ○ 0 Private Insurance • 0 Admitted and discharged • 0 OSH Catchment Area • 0 LSH Catchment Area	●1 SIA admits ●0 paid by HIS ●1 Private Insurance ●0 Admitted and discharged ●0 OSH Catchment Area ●1 LSH Catchment Area

Bread components derived from yeast fermented wheat and rye breads increase colitis and mortality in a mouse model of colitis

Julia Zimmermann, Luigia De Fazio, Valentina Kaden Volynets, Bernd Hitzmann, Stephan C. Bischoff*

*Corresponding author: bischoff.stephan@uni-hohenheim.de; Department of Nutritional Medicine/Prevention, University of Hohenheim, Fruwirthstrasse 12, Stuttgart, 70599 Germany

Current Clients Receiving Services

06/06/25 to 06/12/25

- 2 SIA admits
 - 0 paid by HIS
 - 2 Private Insurance
- 1 OSH Catchment Area
- 1 LSH Catchment Area

Current Clients Receiving Services

06/06/25 to 06/12/25

- 1 SIA admits
 - 0 paid by HIS
 - 1 Private Insurance
- 0 OSH Catchment Area
- 1 LSH Catchment Area

Cumulative Totals 04/30/24 – 06/12/25

06/12/25

- 110 Total discharges
- 112 Total SIA clients served

- 18 OSH catchment area
- 94 LSH catchment area

South Central MC

***Facility closed on 07/22/2022

Cumulative Totals 11/27/24 –

06/12/25

- 82 Total discharges
- 83 Total SIA clients

- 56 OSH catchment area
- 27 LSH catchment Area

SCK-Health

***Facility closed on 01/31/2023

Cumulative Totals 09/27/21 – 07/28/22

- 37 total discharges
- 37 Total SIA clients served
 - 21 OSH Catchment area
 - 16 LSH Catchment area

Cumulative Totals 08/17/22 – 01/31/23

- 6 total discharges
- 6 Total SIA clients served
 - 4 OSH Catchment area
 - 2 LSH Catchment area

Prairie View

***Facility closed on 02/22/2024

Cumulative Totals 08/30/2021 – 02/22/24

- 289 total discharges
- 289 Total SIA clients served
 - 105 OSH Catchment area
 - 184 LSH Catchment area

SIA Youth Census

Camber Kansas City

***5 SIA bed as of 9:02am 06/13/25

Camber Wichita

***10 SIA bed as of 9:03am 06/13/25

Bread components derived from yeast fermented wheat and rye breads increase colitis and mortality in a mouse model of colitis

Julia Zimmermann, Luigia De Fazio, Valentina Kaden-Volynets, Bernd Hitzmann, Stephan C. Bischoff*

*Corresponding author: bischoff.stephan@uni-hohenheim.de; Department of Nutritional Medicine/Prevention, University of Hohenheim, Fruwirthstrasse 12, Stuttgart, 70599 Germany

New Admits 06/06/25 to 06/12/25

- 0 new SIA admits
 - 0 paid by HIS
 - 0 Private Insurance
- 0 Admitted and discharged

New Admits 06/06/25 to 06/12/25

- 1 new SIA admits
 - 0 paid by HIS
 - 1 Private Insurance
- 0 Admitted and discharged

Current Clients Receiving Services 06/06/25 to 06/12/25

- 0 SIA admits
 - 0 paid by HIS
 - 0 Private Insurance

Current Clients Receiving Services 06/06/25 to 06/12/25

- 1 SIA admits
 - 0 paid by HIS
 - 1 Private Insurance

Cumulative Totals 8/30/21 – 06/12/25

- 470 Total discharges
- **470 Total SIA clients served**
served

Cumulative Totals 8/30/21 – 06/12/25

- 881 total discharges
- **882 Total SIA clients**

Camber Hays

***10 SIA bed as of 9:03am 06/13/25

New Admits 06/06/25 to 06/12/25

- 0 new SIA admits
 - 0 paid by HIS
 - 0 Private Insurance
- 0 Admitted and discharged

Via Christi (youth)

***1 open SIA bed as of 8:27am 06/13/2025

New Admits 06/06/25 to 06/12/25

- 0 new SIA admits
 - 0 paid by HIS
 - 0 Private insurance
- 0 Admit and discharged

Current Clients Receiving Services 06/06/25 to 06/12/25

- 0 SIA admits
 - 0 paid by HIS
 - 0 Private Insurance

Current Clients Receiving Services 06/06/25 to 06/12/25

- 0 SIA admits
 - 0 paid by HIS
 - 0 Private Insurance

Cumulative Totals 02/06/23 – 06/12/25

- 173 Total discharges
- **173 Total SIA clients served**
served

Cumulative Totals 09/27/21 – 06/12/25

- 24 Total discharged
- **24 Total SIA clients**

~~Bread components derived from yeast fermented wheat and rye breads increase colitis and mortality in a mouse model of colitis~~

Julia Zimmermann, Luigia De Fazio, Valentina Kaden Volynets, Bernd Hitzmann, Stephan C. Bischoff*

*Corresponding author: bischoff.stephan@uni-hohenheim.de; Department of Nutritional Medicine/Prevention, University of Hohenheim, Fruwirthstrasse 12, Stuttgart, 70599 Germany

Children's Mercy Camber

*****6 SIA bed as of 9:01am 06/13/25**

New Admits 06/06/25 to 06/12/25

- 0 new SIA admits
 - 0 paid by HIS
 - 0 Private Insurance
- 0 Admitted and discharged

Current Clients Receiving Services

06/06/25 to 06/12/25

- 0 SIA admits
 - 0 paid by HIS
 - 0 Private Insurance

Cumulative Totals 11/27/24 to 06/12/25

- 23 Total discharges
- **23 Total SIA clients served**

Administrative Transfers:

Administrative Transfers OSH

New Transfers 06/06/25 to 06/12/25

- 0 CWS
- 0 Hutchinson Regional
- 0 Via Christi
- 0 ANEW Health
- 0 SCK – Health

Administrative Transfers LSH

New Transfers 06/06/25 to 06/12/25

- 0 CWS
- 0 Hutchinson Regional
- 0 Via Christi
- 0 ANEW Health
- 0 SCK – Health

Bread components derived from yeast fermented wheat and rye breads increase colitis and mortality in a mouse model of colitis

Julia Zimmermann, Luigia De Fazio, Valentina Kaden Volynets, Bernd Hitzmann, Stephan C. Bischoff*

*Corresponding author: bischoff.stephan@uni-hohenheim.de; Department of Nutritional Medicine/Prevention, University of Hohenheim, Fruwirthstrasse 12, Stuttgart, 70599 Germany

• 0 Newton MC

●0 Newton MC

Cumulative Totals – 08/31/21 to 06/12/25

• 20 CWS

• 2 Prairie View

• 9 Via Christi

• 3 ANEW Health

• 0 SCK – Health

• 0 Newton MC

• 0 South Central MC

• 0 Hutchinson Regional

• 34 Total Admin Transfers

Cumulative Totals – 08/31/21 to 06/12/25

●5 CWS

●8 Prairie View

●0 Via Christi

● 0 ANEW Health

●0 SCK Health

●0 Newton MC

●0 South Central

●4 Hutchinson Regional

●17 Total Admin Transfer

Fiscal Year Recaps:

FY 2022 (07/01/2021 to 06/30/2022)

409 Total Adult Persons Served

655 Total Youth Persons Served

~~Bread components derived from yeast fermented wheat and rye breads increase colitis and mortality in a mouse model of colitis~~

Julia Zimmermann, Luigia De Fazio, Valentina Kaden-Volynets, Bernd Hitzmann, Stephan C. Bischoff*

*Corresponding author: bischoff.stephan@uni-hohenheim.de; Department of Nutritional Medicine/Prevention, University of Hohenheim, Fruwirthstrasse 12, Stuttgart, 70599 Germany

Cottonwood Springs

215 Total Clients served

- 178 OSH Catchment
- 37 LSH Catchment

- 145 HIS Reimbursed
- 70 Insured

Prairie View

93 Total Clients served

- 28 OSH Catchment
- 65 LSH Catchment

- 70 HIS Reimbursed
- 23 Insured

Via Christi (Adults)

64 Total Clients served

- 58 OSH Catchment
- 6 LSH Catchment

- 22 HIS Reimbursed
- 42 Insured

Newton MHC

1 Total Client served

- 0 OSH Catchment
- 1 LSH Catchment

- 0 HIS Reimbursed
- 1 Insured

South Central MHC

36 Total Clients served

- 20 OSH Catchment
- 16 LSH Catchment

- 9 HIS Reimbursed
- 27 Insured

Via Christi (Youth)

10 Total Clients served

- 0 HIS Reimbursed
- 10 Insured

Camber Wichita

425 Total Clients served

- 48 HIS Reimbursed
- 377 Insured

Camber Kansas City

220 Total Clients served

- 34 HIS Reimbursed
- 186 Insured

FY 2023 (07/01/2022 to 06/30/2023)

~~Bread components derived from yeast fermented wheat and rye breads increase colitis and mortality in a mouse model of colitis~~

Julia Zimmermann, Luigia De Fazio, Valentina Kaden-Volynets, Bernd Hitzmann, Stephan C. Bischoff*

*Corresponding author: bischoff.stephan@uni-hohenheim.de; Department of Nutritional Medicine/Prevention, University of Hohenheim, Fruwirthstrasse 12, Stuttgart, 70599 Germany

736 Total Adult Persons Served

Cottonwood Springs

366 Total Clients served

- 310 OSH Catchment
- 56 LSH Catchment
- 255 HIS Reimbursed
- 111 Insured

433 Total Youth Persons Served

Prairie View

122 Total Clients served

- 54 OSH Catchment
- 68 LSH Catchment
- 89 HIS Reimbursed
- 33 Insured

Via Christi (Adults)

152 Total Clients served

- 142 OSH Catchment
- 10 LSH Catchment
- 51 HIS Reimbursed
- 101 Insured

Newton MC

1 Total Client served

- 0 OSH Catchment
- 1 LSH Catchment
- 0 HIS Reimbursed
- 1 Insured

South Central MC

1 Total Client served

- 1 OSH Catchment
- 0 LSH Catchment
- 1 HIS Reimbursed
- 0 Insured

SCK-Health

6 Total Clients served

- 4 OSH Catchment
- 2 LSH Catchment
- 0 HIS Reimbursed
- 6 Insured

ANEW Health

88 Total Clients served

- 78 OSH Catchment
- 10 LSH Catchment
- 37 HIS Reimbursed
- 51 Insured

Via Christi (Youth)

6 Total Clients served

- 0 HIS Reimbursed
- 6 Insured

Camber Wichita

253 Total Clients served

- 30 HIS Reimbursed
- 223 Insured

~~Bread components derived from yeast fermented wheat and rye breads increase colitis and mortality in a mouse model of colitis~~

Julia Zimmermann, Luigia De Fazio, Valentina Kaden Volynets, Bernd Hitzmann, Stephan C. Bischoff*

*Corresponding author: bischoff.stephan@uni-hohenheim.de; Department of Nutritional Medicine/Prevention, University of Hohenheim, Fruwirthstrasse 12, Stuttgart, 70599 Germany

Camber Kansas City

121 Total Clients served

- 20 HIS Reimbursed
- 101 Insured

Camber Hays

53 Total Clients served

- 4 HIS Reimbursed
- 49 Insured

FY 2024 (07/01/2023 to 06/30/2024)

777 Total Adult Persons Served

290 Total Youth Persons Served

Cottonwood Springs

414 Total Clients served

- 355 OSH Catchment
- 59 LSH Catchment

- 232 HIS Reimbursed
- 182 Insured

Prairie View

74 Total Clients served

- 23 OSH Catchment
- 51 LSH Catchment

- 58 HIS Reimbursed
- 16 Insured

Via Christi (Adults)

159 Total Clients served

- 152 OSH Catchment
- 7 LSH Catchment

- 52 HIS Reimbursed
- 107 Insured

Newton MC

11 Total Clients served

- 2 OSH Catchment
- 9 LSH Catchment

- 7 HIS Reimbursed
- 4 Insured

ANEW Health

105 Total Clients served

- 94 OSH Catchment
- 11 LSH Catchment

- 50 HIS Reimbursed
- 55 Insured

Hutchison Regional

14 Total Clients served

- 3 OSH Catchment
- 11 LSH Catchment

- 7 HIS Reimbursed
- 7 Insured

Via Christi (Youth)

6 Total Clients served

- 1 HIS Reimbursed
- 5 Insured

Camber Wichita

132 Total Clients served

- 43 HIS Reimbursed
- 89 Insured

~~Bread components derived from yeast fermented wheat and rye breads increase colitis and mortality in a mouse model of colitis~~

Julia Zimmermann, Luigia De Fazio, Valentina Kaden Volynets, Bernd Hitzmann, Stephan C. Bischoff*

*Corresponding author: bischoff.stephan@uni-hohenheim.de; Department of Nutritional Medicine/Prevention, University of Hohenheim, Fruwirthstrasse 12, Stuttgart, 70599 Germany

Camber Kansas City

78 Total Clients served

- 28 HIS Reimbursed
- 50 Insured

Camber Hays

74 Total Clients served

- 11 HIS Reimbursed
- 63 Insured

FY 2025 (07/01/2024 to 06/30/2025)

933 Total Adult Persons Served

194 Total Youth Persons Served

Cottonwood Springs

481 Total Clients served

- 425 OSH Catchment
- 56 LSH Catchment

- 275 HIS Reimbursed
- 206 Insured

Newton MC

19 Total Clients served

- 6 OSH Catchment
- 13 LSH Catchment

- 4 HIS Reimbursed
- 15 Insured

Via Christi (Adults)

170 Total Clients served

- 164 OSH Catchment
- 6 LSH Catchment

- 40 HIS Reimbursed
- 130 Insured

Hutchinson Regional

98 Total Clients served

- 15 OSH Catchment
- 83 LSH Catchment

- 42 HIS Reimbursed
- 56 Insured

ANEW Health

82 Total Clients served

- 69 OSH Catchment
- 13 LSH Catchment

- 27 HIS Reimbursed
- 55 Insured

Children's Mercy Camber (Adult)

83 Total Clients served

- 56 OSH Catchment
- 27 LSH Catchment

- 66 HIS Reimbursed
- 17 Insured

Bread components derived from yeast fermented wheat and rye breads increase colitis and mortality in a mouse model of colitis

Julia Zimmermann, Luigia De Fazio, Valentina Kaden-Volynets, Bernd Hitzmann, Stephan C. Bischoff*

*Corresponding author: bischoff.stephan@uni-hohenheim.de; Department of Nutritional Medicine/Prevention, University of Hohenheim, Fruwirthstrasse 12, Stuttgart, 70599 Germany

Via Christi (Youth)

2 Total Clients served

- 0 HIS Reimbursed
- 2 Insured

Camber Wichita

72 Total Clients served

- 21 HIS Reimbursed
- 51 Insured

Camber Kansas City

51 Total Clients served

- 17 HIS Reimbursed
- 34 Insured

Camber Hays

46 Total Clients served

- 13 HIS Reimbursed
- 33 Insured

Children's Mercy Camber (Youth)

23 Total Clients served

- 9 HIS Reimbursed
- 14 Insured

Supplementary material

Table S1: Amount of Oligosaccharides in the different breads.

Bread	Oligosaccharides DP≤8 [g/100g]	Oligosaccharides DP>8 [g/100g]	Total Oligosaccharides [g/100g]
W-SF-Y	3.6	1.6	5.3
W-SF-S	3.3	1.4	4.6
W-SF-SB	4.4	2.0	6.4
W-WG-Y	4.2	1.6	5.8
Sp-SF-Y	1.8	1.4	3.2
R-SF-Y	5.8	2.2	8.0

Abbreviations: W, Wheat; Sp, Spelt; R, Rye; SF, Superfine flour; WG, Wholegrain flour; Y, Yeast fermented; S, Sourdough fermented; DP, Degree of Polymerization.

Bread components derived from yeast-fermented wheat and rye breads increase colitis and mortality in a mouse model of colitis

Julia Zimmermann, Luigia De Fazio, Valentina Kaden-Volynets, Bernd Hitzmann, Stephan C. Bischoff*

*Corresponding author: bischoff.stephan@uni-hohenheim.de; Department of Nutritional Medicine/Prevention, University of Hohenheim, Fruwirthstrasse 12, Stuttgart, 70599 Germany

Table S2. Average food intake and liver/body weight ratio, body weight changes in Casp8^{ΔIEC} and Casp8^{fl} mice.

	Genotype	GF	5G	WSFY	WSFS	WSFSB	WWGY	SpSFY	RSFY
Food intake/ mouse [g/day]	Casp8 ^{ΔIEC}	2.6 ±0.26	2.5 ±0.25	2.5 ±0.23	2.4 ±0.05	2.7 ±0.12	2.8 ±0.14	2.5 ±0.21	2.6 ±0.15
	Casp8 ^{fl}	2.5 ±0.00	2.8 ±0.14	2.4 ±0.05	2.5 ±0.15	2.7 ±0.10	2.9 ±0.15	2.7 ±0.17	2.6 ±0.12
Liver/Body weight ratio [g]	Casp8 ^{ΔIEC}	0.05 ±0.00	0.04 ±0.00	0.04 ±0.4	0.05 ±0.01	0.05 ±0.00	0.04 ±0.00	0.05 ±0.01	0.05 ±0.01
	Casp8 ^{fl}	0.04 ±0.00	0.03 ±0.01	0.04 ±0.00	0.04 ±0.01	0.04 ±0.00	0.04 ±0.00	0.04 ±0.00	0.04 ±0.01
Body weight changes [%]	Casp8 ^{ΔIEC}	20.3 ±6.45	8.6 ±3.65	2.9 ±5.80*	2.4 ±3.99*	7.5 ±2.50	2.4 ±5.49*	2.4 ±4.18*	3.7 ±5.23***
	Casp8 ^{fl}	13.8 ±2.99	12.1 ±1.68	10.5 ±1.75	12.0 ±1.84	16.6 ±2.48	14.4 ±2.93	11.0 ±2.54	10.2 ±1.56

Mice were fed a gluten-free diet (GF), GF with 5% gluten-added (5G) or a diet with 50% bread. Values are means ± SEM, statistics by two-factor ANOVA. Group differences were assessed by the Tukey's post hoc test. Only differences compared to the GF diet were found and indicated by asterisks (*P<0.05; ***P<0.001). No genotype-specific differences were found. Abbreviations: W, Wheat; Sp, Spelt; R, Rye; SF, Superfine flour; WG, Wholegrain flour; Y, Yeast fermented; S, Sourdough fermented.

Bread components derived from yeast-fermented wheat and rye breads increase colitis and mortality in a mouse model of colitis

Julia Zimmermann, Luigia De Fazio, Valentina Kaden-Volynets, Bernd Hitzmann, Stephan C. Bischoff*

*Corresponding author: bischoff.stephan@uni-hohenheim.de; Department of Nutritional Medicine/Prevention, University of Hohenheim, Fruwirthstrasse 12, Stuttgart, 70599 Germany




Figure S1: Small intestinal barrier analyzed by FITC-Dextran 4000 in plasma. Abbreviations see Table S1.