

Supplementary Tables 1, 2 and 3

Supplementary Table 1: Day-to-day context for engaging in podiatry for both Pre-FC and Post-FC participants

SUPPLEMENTARY TABLE 1. Median number of acquired points for the initial rotor map with each mapping system.

<u>Motivation</u>	<u>Group</u>	<u>Rhythm</u>	<u>Median (IQR)</u>	<u>Car</u>	<u>Mean rank</u>	<u>Ensi</u>	<u>te</u>	<u>Precision</u>	<u>U</u>	<u>Z</u>	<u>p</u>	<u>Effect size (r)</u>
<u>Contribution to society</u>	<u>Pre-FC</u>		<u>5 (4-5)</u>		<u>55.09</u>				<u>1479.5</u>	<u>-1.052</u>	<u>0.290</u>	<u>0.10</u>
	<u>Post-FC</u>				<u>60.24</u>							
<u>Skills to improve quality</u>	<u>Pre-</u>		<u>5 (5-</u>									
<u>Left atrium</u>	<u>FC</u>	<u>22135</u>	<u>5)2576</u>						<u>1476.5</u>	<u>-1.226</u>	<u>0.220</u>	<u>0.11</u>
<u>(number of lifemaps = 17)</u>		<u>[12603 – 30902]</u>	<u>[2368 – 6019]</u>		<u>52 – 6426</u>							
	<u>Post-FC</u>				<u>60.28</u>							
<u>Build human relationships</u>	<u>Pre-FC</u>		<u>5 (4-5)</u>		<u>58.03</u>				<u>1623.5</u>	<u>-0.010</u>	<u>0.992</u>	<u>0.00</u>
	<u>Post-FC</u>				<u>57.98</u>							
<u>Work with a range of</u>	<u>Pre-</u>		<u>5 (4-</u>									
<u>service users</u>	<u>FC</u>	<u>11546</u>	<u>5)2372 – 2427</u>						<u>1613</u>	<u>-0.082</u>	<u>0.934</u>	<u>0.01</u>
<u>Right atrium</u>					<u>57.7623</u>							
<u>(number of maps = 9)</u>		<u>[10380 – 16678]</u>			<u>17 – 3292</u>							
	<u>Post-FC</u>				<u>58.18</u>							
<u>Work with specific groups</u>	<u>Pre-FC</u>		<u>4 (4-5)</u>		<u>58.85</u>				<u>1582.5</u>	<u>-0.256</u>	<u>0.798</u>	<u>0.02</u>
	<u>Post-FC</u>				<u>57.35</u>							
<u>Fulfilling</u>	<u>Pre-FC</u>		<u>5 (4-5)</u>		<u>54.22</u>				<u>1436</u>	<u>-1.393</u>	<u>0.164</u>	<u>0.13</u>
	<u>Post-FC</u>				<u>60.91</u>							
<u>Working under pressure</u>	<u>Pre-FC</u>		<u>4 (4-5)</u>		<u>59.78</u>				<u>1536</u>	<u>-0.544</u>	<u>0.587</u>	<u>0.05</u>
	<u>Post-FC</u>				<u>56.63</u>							
<u>Non-sedentary profession</u>	<u>Pre-FC</u>		<u>4 (3-5)</u>		<u>56.98</u>				<u>1574</u>	<u>-0.303</u>	<u>0.762</u>	<u>0.03</u>
	<u>Post-FC</u>				<u>58.78</u>							
<u>Work in a team</u>	<u>Pre-FC</u>		<u>4 (4-5)</u>		<u>58.31</u>				<u>1609.5</u>	<u>-0.096</u>	<u>0.924</u>	<u>0.01</u>

	<u>Post-FC</u>		<u>57.76</u>				
<u>Offers variety</u>	<u>Pre-FC</u>	<u>5 (4-5)</u>	<u>56.36</u>	<u>1543</u>	<u>-0.54</u> <u>2</u>	<u>0.588</u>	<u>0.05</u>
	<u>Post-FC</u>		<u>59.26</u>				

Note: IQR, * Points in use. Total number of points was 37183 - 92480 for the left atrium, and 14296 - 19785 for the right atrium.

Data are shown as median [interquartile range (25th - 75th percentiles); Pre-FC, before first career; Post-FC, after first career] or as range, if only two maps were created.

Supplementary Table 2: How information on podiatry is accessed for both Pre-FC and Post-FC participants

SUPPLEMENTARY TABLE 2. Detailed procedural results in each patient.

<u>Resource</u>	<u>Patient number</u>	<u>Gro</u> <u>up</u> <u>Nu</u> <u>mb</u>	<u>Med</u> <u>ian</u> <u>(IQR</u> <u>)Effe</u>	<u>Mean</u> <u>rank</u> <u>Num</u> <u>ber-of</u>	<u>U</u>	<u>z</u>	<u>p</u>	<u>Eff</u> <u>ect</u> <u>siz</u> <u>e</u>	<u>ATs ablated and</u> <u>other performed</u>
-----------------	-----------------------	---	---	---	----------	----------	----------	--	--

			er of rotors	ct of rotor ablation sites with STD*		(r)of ablations (apart from focal ablation of rotors and sites with STD).
<u>Health Website</u>	<u>Pre -FC</u>	<u>1 (0-1)</u>	<u>60.542</u>	<u>1448 AT</u>	<u>-1.063</u>	<u>0.288</u> <u>0.10</u> -Small reentry in the posterior LA wall.
				<u>Post-FC</u>		<u>55.13</u>
<u>Job Website2</u>	<u>Pre -FC</u>	<u>0 (0-0)†</u>	<u>56.54 SR</u>	<u>1552</u>	<u>-0.372</u>	<u>0.710</u> -Small reentry in the RSPV + right superior PVI. -Typical atrial flutter.
	<u>3</u>	<u>Pos t-FC†</u>		<u>58.25</u>		<u>-Right superior PVI.</u>
4	1	AT				-Typical atrial flutter. -Focal AT in the posterior LA wall.
5	3	AT				-Perimitral flutter: ASL. -Typical atrial flutter. -SVC isolation + line between SVC and IVC to connect rotor ablation sites.
6	2	AT				-Small reentry in the LA roof. -Perimitral flutter: MAL. -Typical atrial flutter.