

Midtherm Report  
Jan Grzegorzewski

# **Reaction-diffusion dynamics with fractional Brownian motion**

Summer term: 2016

# **Contents**

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Theory</b>	<b>3</b>
2.1	Brownian Motion . . . . .	3
2.2	Fractional Brownian Motion . . . . .	6
2.2.1	Algorithm . . . . .	9
<b>3</b>	<b>Reactions-Diffusion-Dynamics</b>	<b>20</b>
<b>4</b>	<b>Status Quo &amp; Outlook</b>	<b>21</b>
<b>5</b>	<b>Appendix</b>	<b>22</b>
5.0.2	Code:Gebrochen-rationale Brownische Bewegung . . . . .	22
	<b>Bibliography</b>	<b>25</b>

# List of Figures

2.1	Mean Square Displacement of different $\alpha$ . . . . .	12
2.2	Mean Square Displacement (Ensemble average of 4000 Trajectories)	12
2.3	Mean Square Displacement (Ensemble average fpr different $\Delta t$ ) . . .	13
2.4	Mean Square Displacement (time and ensemble avarge for different $\Delta t$ and with standard deviation of the Mean) . . . . .	14
2.5	Non-gaussian parameter of $\alpha = 0.5$ , $D = 2$ and 5000 trajectories ensemble. . . . .	15
2.6	Non-gaussian parameter of $\alpha = 0.5$ , $D = 2$ and 5000 trajectories ensemble. . . . .	16
2.7	rescaled function for different times . . . . .	17
2.8	variere dt bei gleichem Faktor 1 . . . . .	17
2.9	variere dt bei gleichem Faktor 1, logarithmisch . . . . .	18
2.10	variere Faktor bei gleichem dt . . . . .	18
2.11	variere Faktor bei gleichem dt, logarithmisch . . . . .	19

## **List of Tables**

# 1 Introduction

Anomalous diffusion can be observed in many different areas of nature, in particular related to heterogeneous media (Crowded biological media, porous materials, ...). The most popular phenomenon of anomalous diffusion is a power-law behaviour of the Mean-Square-Displacement ( $MSD \propto t^\alpha$ ), which is violating the Einstein formula  $MSD = 2dDt$  and thereby the central limit theorem. Various theoretical models try to encounter the power-law behaviour of the MSD showing different origins for anomalous diffusion. These models have the main phenomenon, thus the power-law behaviour of the mean square displacement, in common. But differ in some other observables, due to different origins of the anomalous power-law behaviour. By measuring theses different quantities one might reveal the origin of the anomalous behaviour in an experiment.

A basic model which accounts this phenomenon so far is Fractional Brownian Motion. This thesis is going to focus on Fractional Brownian Motion especially in respect to Reaction and Diffusion Dynamics. (Normal Diffusion Limited Smoluchowski equation, chemical master equation, mass action law michaelis menden reaction ????? check it ) is failing miserably in describing Reaction and Diffusion Dynamics while anomalous motion is present.(explain ...because of fluctuations). The overall work aims not only to bring Reaction Diffusion and Fractional Brownian Motion together but also providing a Fractional Brownian Motion Integrator implemented into a **Reaction Diffusion Dynamics** software (ReaDDy). ReaDDy being a particle-based **Reaction Diffusion Dynamics** software is acting on a macromolecular level. It is coarse-graining molecules into spherical particles. Usually using Brownian or Langevin motion as the integrator. A Fractional Brownian Motion Integrator may be an elegant way of adding an approximation due to a crowded environment. The phenomenon of anomalous motion can also be studied with conventional integrators by actually building a crowded environment (for example: by adding obstacles). Nevertheless, the Fractional Brownian Motion approach could save computational time since it is not necessary to simulated each particle explicitly which builds up the crowded environment. Furthermore different studies show that heterogeneous environments have influences on particle segregation in the presents of reactions.

Having an Introduction set, chapter 2 deals with the theoretical foundation for Frac-

tional Brownian Motion. Subsequently, due to computational reasons these theoretical elaborations need to be transferred into discrete space and finally casted into an algorithm. In the following some properties of the algorithm are studied. In Chapter 3 theoretical foundation for reactions are going to be set. The final chapter 4 will deal with the present state of the thesis and give an outlook on further challenges.

## 2 Theory

### 2.1 Brownian Motion

Fractional Brownian motion is a generalized case of standard Brownian Motion. The following section will explore in more detail Brownian motion and normal diffusion. Than the more generalized case will be considerer.

Standard Brownian Motion is a very important and good studied stochastic process. It describes the erratic motion for mesoscopic particles, which first have been documented by Jan Ingenhousz in 1785, in particular for coal dust on the surface of alcohol. Later on in 1827 Robert Brown observed the erratic motion of pollen grains. Brownian Motion has an Gaussian propagator which has its origin in the Central Limit Theorem (CLT) for a sum of independent random variables. Lets assume a set of  $N$  independent variables  $\{X_i\}$  with a finite variance  $\sigma_i^2 = \langle X_i^2 \rangle$  and the mean  $\langle X_i \rangle = 0$ . The definition of a another random variable  $Y$  is given by:

$$Y = \frac{1}{\sqrt{N}} \sum_{j=1}^N X_j \quad (2.1)$$

This scenario in which a random variable is defined of the sum of another can be observed generically in nature. In the following the distribution of  $Y$ ,  $\rho(y)dy = P(y < Y < y + dy)$  in the limit of large  $N$ , is going to be calculated. The Generating function for a random variable  $Y$  is:

$$G_Y(k) = \langle e^{ikY} \rangle = \int e^{ikY} \rho(y)dy \quad (2.2)$$

eq. (2.1) can be inserted into the generating function, which results in:

$$\begin{aligned} G_Y(k) &= \langle e^{\frac{ik}{\sqrt{N}} \sum_{j=1}^N X_j} \rangle \\ G_Y(k) &= \langle \prod_{j=i}^N e^{\frac{ik}{\sqrt{N}} X_j} \rangle \end{aligned}$$

If all  $X_j$  are independent, then:

$$G_Y(k) = \prod_{j=i}^N \langle e^{\frac{ik}{\sqrt{N}} X_j} \rangle = e^{\sum_{j=1}^N A_j(\frac{k}{\sqrt{N}})}$$

with  $A_j(\frac{k}{\sqrt{N}}) = \ln \langle e^{\frac{ik}{\sqrt{N}} X_j} \rangle$

(2.3)

For large N behaviour, we assume  $\frac{k}{\sqrt{N}} \ll 1$  and expand

$$A_j(\frac{k}{\sqrt{N}}) = \ln(1 + \langle X_j \rangle \frac{ik}{\sqrt{N}} - \langle X_j^2 \rangle \frac{k^2}{2N} + \mathcal{O}(N^{-\frac{3}{2}}))$$
(2.4)

with a finite variance  $\sigma_i^2 = \langle X_i^2 \rangle$  and the mean  $\langle X_i \rangle = 0$

$$A_j(\frac{k}{\sqrt{N}}) = -\sigma_j^2 \frac{k^2}{2N} + \mathcal{O}(N^{-\frac{3}{2}}))$$
(2.5)

Thus, the generating function for large N is:

$$G_Y(k) = e^{-\frac{\sigma^2 k^2}{2}}$$
(2.6)

with  $\sigma = \frac{1}{N} \sum_{j=1}^N \sigma_j^2$

The distribution of Y can be calculated via the inverse Fourier Transform:

$$\rho(y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-\frac{\sigma^2 k^2}{2}} e^{iky} dk$$
(2.7)

$$= \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{y^2}{2\sigma^2}}$$
(2.8)

The seemingly innocent assumption of independence for the random variables  $\{X_i\}$  in the summation result in an Gaussian distribution. Microscopic processes, which result in independent random position changes of an particle and add up over time, thus have a Gaussian distribution function for the overall position change. The propagator for Brownian motion has a Gaussian distribution.

**Definition 1** Standard Brownian motion is an stochastic process  $\{W_t\}_{t \geq 0} : \Omega \rightarrow \mathbb{R}^n$  with  $W_t(\omega)$  being the position of a particle with  $\omega \in \Omega$  at time  $t \in T$  in the

*observation time  $T = [0, \infty)$ . It has a fixed  $x \in \mathbb{R}^n$  as its origin. The transition probabilities are:*

$$\begin{aligned} T_t(y|x) &:= (2\pi t)^{-\frac{n}{2}} e^{-\frac{\|x-y\|^2}{2t}} \text{ for } y \in \mathbb{R}^n, t > 0 \\ T_0(y|x) &= \delta(x - y) \end{aligned} \quad (2.9)$$

In the following some properties of Brownian motion are discussed:

- Due to the Kolmogorov's continuity theorem a continuous path  $t \mapsto W_t(\omega)$  almost surely exists. Therefore it is possible to go to a continuous function  $[0, \infty) \rightarrow \mathbb{R}^n$  and define a probability density.

$$p_{t_1, \dots, t_k}(x_1, \dots, x_k) = T_{t_k-t_{k-1}}(x_k|x_{k-1}) \dots T_{t_2-t_1}(x_2|x_1) T_{t_1}(x_1|x) \quad \text{with } 0 \leq t_1 < t_2 < \dots < t_k, k \in \mathbb{N}$$

- Brownian motion is a Gaussian process with mean  $E^x[W_t] = x$ ,  $W_o = x$  Since  $E^x[(W_{t_i} - W_{t_{i-1}})(W_{t_j} - W_{t_{j-1}})] = 0$  all its increments  $\{W_{t_1}, W_{t_2} - W_{t_1}, \dots, W_{t_k} - W_{t_{k-1}}\}$  are independent.
- Brownian motion has stationary increments since  $W_{t-h} - W_h$  has the same distribution for all  $h > 0$ .
- Brownian scaling  $\{\hat{W}_t := \frac{1}{c}W_{c^2t}\}_{t \geq 0}$  if  $\{W_t\}_{t \geq 0}$  is also a Brownian motion. Therefore Brownian motion has self-similar and fractal paths.

For Brownian motion a linear relation between the MSD and the diffusion constant can be shown. The starting point is the diffusion equation for overdamped motion with a constant diffusion coefficient  $D$ .

$$\frac{\partial}{\partial t} C(x, t) = -\frac{\partial}{\partial x} J(x, t) = D \frac{\partial^2}{\partial x^2} C(x, t) \quad (2.10)$$

Flux of particles  $J(x, t) = \frac{\partial}{\partial x} C(x, t)$  as a result of a linear response relation. Under this approximation diffusion only depends on the position and not on the velocity. Therefore it is again a long-time approximation for normal diffusion. The propagator of Brownian motion solves this equation. The Concentration  $C(x, t)$  can be interpreted as a probability distribution if properly normalized as  $\int dx C(x, t) = 1$ .

The Mean Square Displacement (MSD) can be calculated as:

$$\frac{d}{dt} \langle x^2(t) \rangle = \frac{d}{dt} \int dx x^2 C(x, t) = \int dx x^2 \frac{\partial}{\partial t} C(x, t) \quad (2.11)$$

$$= D \int_{-\infty}^{\infty} dx x^2 \frac{\partial^2}{\partial x^2} C(x, t) = -2D \int_{-\infty}^{\infty} dx x \frac{\partial}{\partial x} C(x, t) \quad (2.12)$$

$$= 2D \int_{-\infty}^{\infty} dx C(x, t) = 2D \quad (2.13)$$

In previous equations partial integration has been used and  $C(\pm\infty, t) = 0$  has been assumed. By integration for the initial condition  $x(0) = 0$  one gets the Einstein Formula:  $\langle (x(t) - x(0))^2 \rangle = 2Dt$ , which can easily be extended for 2 and 3 dimensions. It will not be calculated here explicitly, but one can show that also for Smoluchowski eq. which follows from langevin dynamics for a long time limit the initial velocities decorrelates and thus the MSD is also linear with time.

## 2.2 Fractional Brownian Motion

In this section the theoretical foundation for Fractional Brownian Motion will be set and related to standard Brownian Motion.

In the previous section the the Means Square Displacement (MSD) has been shown to be linear with time as a result of the central limit theorem. In normal liquids this behaviour can be seen already at time scales higher than picoseconds [2]. Nevertheless many experiments show that the MSDs has a power law behaviour ( $\delta r^2(t) \propto t^\alpha$  for  $0 < \alpha < 1$ ). Thus the central limit theorem does not hold, not even for long time scales. It can be shown that persistent correlations of the increments can be observed. In soft matter, like polymers, subdiffusive behaviour is typically present in an time window but finally the linear MSD takes over. In particular Fractional Brownian Motion examines the case that the central limit theorem is violated for all time scales.

fractional Brownian scaling

In the following some statistical tools are defined. They are important as the increments of Fractional Brownian motion are no longer assumed to be independent. The single particle density  $\rho(\mathbf{r}, t) = \delta(\mathbf{r} - \mathbf{R}(t))$  describes the density of a particle which is localized at position  $\mathbf{R}(t)$ . Its correlation function  $P(\mathbf{r} - \mathbf{r}', t - t') = V \langle \rho(\mathbf{r}, t) \rho(\mathbf{r}', t') \rangle$  is also called Van Hove self-correlation function.  $V$  refers to the volume. From now on we will consider an isotropic system  $r = |\mathbf{r}|$ . As for Brownian motion with independent increments the correlated increments  $\Delta \mathbf{R}(t)$  of fractional Brownian motion

are assumed to follow a Gaussian distribution with zero mean. Thus the correlation function of the single particle density results in:

$$P(r, t) = [2\pi\delta r^2(t)/d]^{-\frac{d}{2}} e^{\frac{-r^2 d}{2\delta r^2(t)}} \quad (2.14)$$

The van Hove correlation function can be transformed via the spatial Fourier transform into the wave-number representation, which is called the self-intermediate scattering function. Again for isotropic systems one can write  $|\mathbf{k}| = k$ .

$$F_s(k, t) = \langle \rho(\mathbf{k}, t)\rho(\mathbf{k}', t') \rangle = \int d^d r e^{-ikr} P(r, t) \quad (2.15)$$

$$= \langle e^{-ik\Delta R(t)} \rangle \quad (2.16)$$

The intermediate scattering function for the single particle density turns out to be the characteristic or moment generating function of  $\Delta R(t)$  by expanding it for small wave-numbers  $k \rightarrow 0$  one can get the moments. Its logarithm return the cumulants. For Gaussian propagators all but the second cumulants vanish. For non Gaussian transport also the fourth cumulant is non-zero. Therefore it is used to indicate beyond Gaussian transport. The non-Gaussian parameter is defined as:

$$\alpha_2 = \frac{d\delta r^4(t)}{(d+2)[\delta r^2(t)] - 1} \quad (2.17)$$

An other important quantity is the dynamical structure factor, which is the time-frequency Fourier transform of the intermediate scattering function:

$$F_s(\mathbf{k}, z) = \langle \rho(\mathbf{k}, z)\rho(\mathbf{k}', z') \rangle = \int_0^\infty dt e^{-i\mathbf{k}\mathbf{r}} P(r, t) \text{ for } k \rightarrow 0, \text{Im}(z) > 0 \quad (2.18)$$

$$= \frac{1}{-iz} - \frac{k^2}{2d} \int_0^\infty dt e^{izt} \delta r^2(t) + \mathcal{O}(k^2) \quad (2.19)$$

From now on lets a differential equation  $\partial_t \mathbf{R}(t) = \boldsymbol{\eta}(t)$ .  $\boldsymbol{\eta}(t)$  are velocities, which are no more delta-correlated in time as they would be for standard Brownian motion. Velocities can be used to calculate the Velocity Autocorrelation Function (VACF):

$$Z(|t - t'|) = \frac{1}{d} \langle \boldsymbol{\eta}(t)\boldsymbol{\eta}(t') \rangle = \frac{1}{2d} \frac{d^2}{dt^2} \delta r^2(t - t') \quad (2.20)$$

$$(2.21)$$

Subsequently, the VACF in the frequency domain for Fractional Brownian motion will be calculated. The MSD is  $\delta r^2(t) = \langle \Delta R(t) \rangle = 2dK_\alpha t^\alpha$  with  $K_\alpha$  being the generalized diffusion-coefficient:

$$\begin{aligned}\tilde{Z}(z) &= \int_0^\infty dt e^{izt} Z(t) \\ &= 2d \int_0^\infty dt e^{izt} \left[ \frac{d^2}{dt^2} \delta r^2(t) \right] \\ &\stackrel{\text{par.integ.}}{=} 2d \left( \underbrace{\left[ e^{izt} \overbrace{\frac{d^2}{dt^2} 2dK_\alpha t^\alpha}^{=A(t)} \right]_0^\infty - \int_0^\infty dt e^{izt} \left[ \frac{d}{dt} \delta r^2(t) \right]}_{\alpha \leq 2} \right) \\ A(t) &= \frac{d}{dt} \overbrace{\left[ \frac{2dK_\alpha t^{\alpha-1}}{\alpha} \right]}^{B(t)} = \frac{2dK_\alpha t^{\alpha-2}}{\alpha + (\alpha - 1)} \\ \tilde{Z}(z) &\stackrel{\text{par.integ.}}{=} -2d \left( \underbrace{\left[ e^{izt} \overbrace{\frac{d}{dt} 2dK_\alpha t^\alpha}^{=B(t)} \right]_0^\infty - \int_0^\infty dt e^{izt} \delta r^2(t)}_{\alpha \leq 1} \right) \\ &= \int_0^\infty dt e^{izt} \delta r^2(t) \stackrel{\text{Im}(z) > 0}{=} K_\alpha \Gamma(1 + \alpha) (iz)^{1-\alpha}\end{aligned}$$

Eventually, the VACF in the frequency domain will be used to modify standard Brownian motion velocities, which are easily computable, to generate fractional Brownian motion velocities. The starting point is again the differential equation  $\partial_t \mathbf{R}(t) = \boldsymbol{\eta}(t)$ . In the frequency domain they are represented as:

$$\tilde{\boldsymbol{\eta}}(z) = \int_0^\infty dt e^{izt} \boldsymbol{\eta}(t) \tag{2.22}$$

Fractional correlations can be incorporated via its VACF:

$$\tilde{\xi}(z) = \sqrt{2 \operatorname{Re}(\tilde{Z}(z))} \tilde{\eta}(z) \quad (2.23)$$

With  $\tilde{\xi}(z)$  being fractional Brownian velocities in the frequency domain. Its Fourier-back-transform results in Fractional Brownian velocities in the time domain.

$$\xi(t) = \frac{1}{2\pi} \int dz \tilde{\xi}(z) \quad (2.24)$$

### 2.2.1 Algorithm

In the following an algorithm which generates fractional Brownian noise will be introduced. The algorithm is based on the Davis-Harte algorythm [1]. The idea is to use the calculated VACF and thereby modify conventionally generated Gaussian random variables. All the increments should be generated beforehand. With this concept it is more difficult to include forces. Nevertheless it is computationally favourable then computing each increment recursively considering also its history. This would be certainly necessary since all increments are defined to be more then just delta-correlated in time. For computational reasons the elaborations for fractional Brownian motion in the previous chapter on how to generate fractional Brownian increments have to be transformed into a discrete from thereby the solution is no longer exact, which will be shown in the analyse part of the algorithm.

$$\boldsymbol{\eta}(t) \longrightarrow \boldsymbol{\eta}_j(t) \text{ with } j = 0, 1, 2, \dots, n, n = \text{amount of steps} \quad (2.25)$$

For a n-steps long trajectory one can write:

$$\Delta \mathbf{R}_n(t) = \sum_{j=0}^n \boldsymbol{\eta}_j \Delta t \quad (2.26)$$

The MSD then can be written as:

$$\langle \Delta \mathbf{R}_j(t) \rangle = 2dK_\alpha(\Delta t j)^\alpha \quad (2.27)$$

The algorithm goes as follows:

1.  $2n$  independent normally distributed random increments are generated:

$$\boldsymbol{\eta}_k(t) = \mathcal{N}(0, \sqrt{\Delta t}) \text{ with } k = 0, 1, 2, \dots, 2n \quad (2.28)$$

2 times more increments are generated to counteract the boundary problem in the discrete Fourier transform.

2. Via discrete Fourier transform these increments are transformed into the frequency domain:

$$\tilde{\eta}_k(z) = \sum_{j=0}^{2n-1} \eta_k e^{\frac{-i2\pi j k}{2n}} \Delta t \quad (2.29)$$

$$(2.30)$$

By comparison with the eq. (2.22) one can see that:

$$z \rightarrow g\Delta z, \Delta z = \frac{2\pi}{2n\Delta t}, t \rightarrow j\Delta t \text{ and } \int dt \rightarrow \sum \Delta t \quad (2.31)$$

3. Comparable to eq. (2.23) correlations are incorporated:

$$\xi_k(z) = \eta_k(z) \sqrt{2\text{Re}(Z_k(z))} \quad (2.32)$$

with  $\tilde{Z}(z) \rightarrow \tilde{Z}_k(z)$  as introduced in eq. (2.31):

$$\tilde{Z}_k(z) = K_\alpha \Gamma(1 + \alpha) (i2\pi k \Delta \omega)^{1-\alpha} = K_\alpha \Gamma(1 + \alpha) (ik \frac{\pi}{n \Delta t})^{1-\alpha} \quad (2.33)$$

4. The discrete Fourier transform has a downside compared to the continuous Fourier transform, as already noted in the beginning of this section. The VACF is zero at zero-frequency  $Z_{k=0}(0) = 0$ . From eq. (2.32) also the first increment in the frequency domain is zero  $\xi_{k=0}(z) = 0$ . Due to eq. (2.29) also the following relation holds:

$$\xi_{k=0}(z) = \sum_{j=0}^{2n-1} \eta_j e^0 \Delta t = \Delta R \quad (2.34)$$

$\Delta R$  ist in dem Fall die zurückgelegte Entfernung nach  $2n$  Zeitschritten. Dadurch würde das Teilchen nach  $2n$  Schritten wieder zurück an seinen Ursprung kehren. Stattdessen wird Das Nullte Inkrement im Frequenzraum folgendermaßen berechnet:

$$\eta_{fbR_{g=0}} = \mathcal{N}(0, \sqrt{2K_\alpha(2n\Delta t)^\alpha}) \quad (2.35)$$

gerechnet. Dies wäre korrekt, wenn Gebrochen-rationale Brownische Bewegung ein Markowischer Prozess wäre. Da dies nicht der Fall ist, entsteht an dieser Stelle eine Approximation. Um sie genauer zu verstehen muss ich mich

damit noch weiter beschäftigen. Um den Einfluss der Approximation zu verringern wurde in Schritt 1 die doppelte Menge an Inkrementen erstellt. Die Hoffnung ist, dass der Einfluss der Approximation mit zunehmender Entfernung von  $2\Delta R_{fbr}$  immer geringer wird und bei  $\Delta R_{fbr}$  bereits vernachlässigbar ist.

5. In der Zeitdomäne ergeben sich die Inkremente durch die Rücktransformation als:

$$\eta_{fbr_j} = \frac{1}{2n} \sum_{g=0}^{2n-1} \eta_g e^{\frac{2\pi i j g}{2n}} \quad (2.36)$$

Es wird nur die erste Hälfte der Inkremente,  $\eta_{fbr_j}$  für  $j = (0, 1, \dots, n)$ , weiter verwendet.

Für die drei dimensionale Gebrochen-rationale Brownische Bewegung kann für jede kartesische Komponente der soeben beschriebene Algorithmus verwendet werden, da davon ausgegangen wird, dass die Kartesischen Komponenten der Inkremente untereinander nicht korreliert sind.

### **Analyse of Algorythm**

Hier kommen die Plots aus dem Analyse Skript rein.




Figure 2.1: Mean Square Displacement of different  $\alpha$ 

Figure 2.2: Mean Square Displacement (Ensemble average of 4000 Trajectories)

Figure 2.3: Mean Square Displacement (Ensemble average fpr different  $\Delta t$ )




Figure 2.4: Mean Square Displacement (time and ensemble average for different  $\Delta t$  and with standard deviation of the Mean)




Figure 2.5: Non-gaussian parameter of  $\alpha = 0.5$ ,  $D = 2$  and 5000 trajectories ensemble.




Figure 2.6: Non-gaussian parameter of  $\alpha = 0.5$ ,  $D = 2$  and 5000 trajectories ensemble.





Figure 2.7: rescaled function for different times

Figure 2.8: variere  $dt$  bei gleichem Faktor 1




Figure 2.9: variere  $dt$  bei gleichem Faktor 1, logarithmisch




Figure 2.10: variere Faktor bei gleichem  $dt$




Figure 2.11: variere Faktor bei gleichem dt, logarithmisch

### **3 Reactions-Diffusion-Dynamics**

## **4 Status Quo & Outlook**

# 5 Appendix

## 5.0.2 Code: Gebrochen-rationale Brownische Bewegung

```
1 # -*- coding: utf-8 -*-
2 --author__ = 'janek'
3 import numpy as np # module for scientific computing
4 from scipy import integrate
5 import matplotlib.pyplot as plt # module for plotting "a la" matlab
6 import test_cython.genereatefracincrements as ginc
7
8
9 class Felix_Method():
10
11     def __init__(self, D, particles, length, alpha, dt, x=2, version="python"):
12         self.D=D
13         self.K_alpha=D
14         self.particles=particles
15         self.n=length
16         self.alpha=alpha
17         self.ki=self.n*x
18         self.frq=np.fft.freq(np.pi*2.)/(dt))
19         self.t=np.array(range(length))
20         self.dt=dt
21         self.version=version
22
23     def z(self):
24         """
25             :param D: Diffusionskoeffizient
26             :param alpha: Anomalieparameter
27
28
29             :return: z: Correlationsfunktion im Frequenzraum welche auf die
30             gew hnliche diffusion angewendet wird.
31
32
33
```

```

35     """
36     z=((((+1j* self .frq ))** (1-self .alpha ))* self .K_alpha *np.math .
37     gamma(1+ self .alpha ))
38     return z
39
40     def compute_trajectory( self ):
41         """
42             :param D: Diffusionskoeffizient
43             :param particles: Anzahl an Teilechen welche simuliert werden
44             sollen .
45
46             :param length: L nge der Trajektorien , welche simuliert werden
47             sollen .
48
49             :param alpha: Anomalieparameter
50
51             :return: ( Particles x Length ) Array of Trajectories of all
52             particles. Close to :cite:`Graigmile2003`"
53
54         """
55
56         #plt.plot( self .frq ,”r”)
57         #plt.plot( np.arange(-np.pi/ self .dt , np.pi/ self .dt ,(np.pi/(
58             self .n* self .dt ))))
59         #plt.show()
60         r_t_allparticles=[]
61         if self .version == "cpp":
62             inc = ginc.pyIncrements( self .n, self .particles )
63             inc.generateIncrements( self .D, self .dt , self .alpha )
64             a=inc.returnIncrements()
65             r_t=np.cumsum(a[:,0,:], axis=1)
66             r_t[:,0]=0
67             return r_t
68         if self .version == "python":
69             for particle in range( self .particles):
70                 #r = np.random.RandomState(1234)
71                 v_t=(np.random.normal(0,np.sqrt( self .dt ),size= self .ki))
72                 #todo mit matrix.shape k nnen man eine zuf llige verteilung von
73                 #allen daten generieren
74                 v_t=np.array(v_t)
75                 v_fraq=np.fft.fft(v_t)
76                 v_ano_fraq= np.sqrt( self .z().real*2.)*v_fraq
77                 v_ano_fraq[0]=np.random.normal(0,np.sqrt(2.* self .K_alpha
78                     *( self .ki* self .dt )** self .alpha ))

```

```

73         #v_ano_frq [ self .n]=np .sqrt ( self .z () [ self .n]. real* self .n
    *2)* v_t [ self .ki-1]
    #v_ano_frq [ self .n-1]=np .sqrt ( self .z () [ self .n]. real* self .
    .n*2)* v_t [ self .ki-1]
    v_ano_t=np .fft .ifft ( v_ano_frq)
    #r_t1=np .cumsum( v_ano_t [: self .n]. real) #Ort bei
75 anomaler Diffusion in Abh ngigkeit von der zeit
    r_t=np .zeros( self .n)
    r_t [1:]=np .cumsum( v_ano_t [: self .n]. real) [1:] #Ort bei
77 anomaler Diffusion in Abh ngigkeit von der zeit
    r_t_allparticles.append(r_t) # Trajektorie bei anomaler
    Diffusion f r alle teilchen
    return np .array( r_t_allparticles)
79

```

..../simulation.py

## Bibliography

- [1] Peter F. Craigmile.  
Simulating a class of stationary Gaussian processes using the Davies-Harte algorithm, with application to long memory processes.  
*Journal of Time Series Analysis*, 24(5):505–511, 2003.
- [2] Felix Höfling and Thomas Franosch.  
Anomalous transport in the crowded world of biological cells.  
*Reports on Progress in Physics*, 76(4):046602, apr 2013.

inspired by [http://www.weizmann.ac.il/weizsites/mukamel/files/2013/01/Brownian\\_Motion.pdf](http://www.weizmann.ac.il/weizsites/mukamel/files/2013/01/Brownian_Motion.pdf)