

Linear Regression

Newton's three sisters

February 16, 2025

Department of Statistics
Sungshin Women's University

1 Distribution of the RSS Values

2 Hypothesis Testing for $\hat{\beta}_j \neq 0$

3 Coefficient of Determination and the Detection of Collinearity

4 Confidence and Prediction Intervals

- Hat matrix defined by $\hat{y} = Hy$

$$\hat{y} = X\hat{\beta} = X(X^T X)^{-1} X^T y = Hy$$

$$H \triangleq X(X^T X)^{-1} X^T$$

- Some properties

$$H^2 = X(X^T X)^{-1} X^T \cdot X(X^T X)^{-1} X^T = X(X^T X)^{-1} X^T = H$$

$$(I - H)^2 = I - 2H + H^2 = I - H$$

$$HX = X(X^T X)^{-1} X^T \cdot X = X$$

- RSS defined

$$RSS \triangleq ||y - \hat{y}||^2$$

- Using hat matrix

$$\begin{aligned} y - \hat{y} &= y - Hy = (I - H)y = (I - H)(X\beta + \varepsilon) \\ &= (X - HX)\beta + (I - H)\varepsilon = (I - H)\varepsilon \end{aligned}$$

$$RSS \triangleq ||y - \hat{y}||^2 = \{(I - H)\varepsilon\}^T (I - H)\varepsilon = \varepsilon^T (I - H)^2 \varepsilon = \varepsilon^T (I - H)\varepsilon$$

Eigenvalues of H and Null space of $(I - H)$

- Proof by contrapositive

$$Hx = x \Rightarrow (I - H)x = 0$$

$$(I - H)x = 0 \Rightarrow Hx = x$$

- Dimensions of the eigenspaces of H is $p + 1$

Proof using $\text{rank}(X) = p + 1$

$$\text{rank}(H) \leq \min\{\text{rank}(X(X^T X)^{-1}), \text{rank}(X)\} \leq \text{rank}(X) = p + 1$$

$$\text{rank}(H) \geq \text{rank}(HX) = \text{rank}(X) = p + 1$$

- Dimensions of the null space of $I - H$ is $N - (p + 1)$

$$P(I - H)P^T = \text{diag}(\underbrace{1, \dots, 1}_{N-p-1}, \underbrace{0, \dots, 0}_{p+1})$$

Residual Sum of Squares (RSS) and Eigenvalue

- We define $v = P\varepsilon \in \mathbb{R}^N$, then from $\varepsilon = P^T v$

$$\text{RSS} = \varepsilon^T (I - H) \varepsilon = (P^T v)^T (I - H) P^T v = v^T P (I - H) P^T v$$

$$= [v_1, \dots, v_{N-p-1}, v_{N-p}, \dots, v_N] \begin{bmatrix} 1 & 0 & \dots & \dots & \dots & 0 \\ 0 & \ddots & 0 & \dots & \dots & \vdots \\ \vdots & 0 & 1 & \dots & \dots & 0 \\ 0 & 0 & 0 & \dots & \dots & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \dots & 0 & \dots & \dots & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ \vdots \\ v_{N-p-1} \\ v_{N-p} \\ \vdots \\ v_N \end{bmatrix}$$

$$= \sum_{i=1}^{N-p-1} v_i^2$$

- Let $w \in \mathbb{R}^{N-p-1}$ be

- Average

$$E[v] = E[P\varepsilon] = 0$$

$$E[w] = 0$$

- Covariance

$$E[vv^t] = E[P\varepsilon(P\varepsilon)^T] = PE[\varepsilon\varepsilon^t]P = P\sigma^2IP^T = \sigma^2I$$

$$E[ww^T] = \sigma^2I$$

- We have RSS

$$\frac{RSS}{\sigma^2} \sim \chi_{N-p-1}^2$$

1 Distribution of the RSS Values

2 Hypothesis Testing for $\hat{\beta}_j \neq 0$


3 Coefficient of Determination and the Detection of Collinearity

4 Confidence and Prediction Intervals

Test statistic T


- A t distribution with $N - P - 1$ degrees of freedom
- We decide that hypothesis $\beta_j = 0$ should be rejected.
- $U \sim N(0, 1)$, $V \sim \chi_m^2$,

$$T \triangleq U / \sqrt{V/m}$$



Significance level

- $\alpha = 0.01, 0.05$
- Null hypothesis $\beta_j = 0$



Example 23

- For $p = 1$, since

$$X^T X = \begin{bmatrix} 1 & \cdots & 1 \\ x_1 & \cdots & x_N \end{bmatrix} \begin{bmatrix} 1 & x_1 \\ \vdots & \vdots \\ 1 & x_n \end{bmatrix} = N \begin{bmatrix} 1 & \bar{x} \\ \bar{x} & \frac{1}{N} \sum_{i=1}^N x_i^2 \end{bmatrix}$$

- The inverse is

$$(X^T X)^{-1} = \frac{1}{N} \begin{bmatrix} 1 & \bar{x} \\ \bar{x} & \frac{1}{N} \sum_{i=1}^N x_i^2 \end{bmatrix}^{-1} = \frac{1}{\sum_{i=1}^N (x_i - \bar{x})^2} \begin{bmatrix} \frac{1}{N} \sum_{i=1}^N x_i^2 & -\bar{x} \\ -\bar{x} & 1 \end{bmatrix}$$

- Which means that

$$B_0 = \frac{\frac{1}{N} \sum_{i=1}^N x_i^2}{\sum_{i=1}^N (x_i - \bar{x})^2} \quad \text{and} \quad B_1 = \frac{1}{\sum_{i=1}^N (x_i - \bar{x})^2}$$

Example 23 (contd.)

- For $B = (X^T X)^{-1}$, $B\sigma^2$ is covariance matrix of $\hat{\beta}$
- $B_j\sigma^2$ is the variance of $\hat{\beta}_j$
- Because \bar{x} is positive, the correlation between $\hat{\beta}_0$ and $\hat{\beta}_1$ is negative

$$t = \frac{\hat{\beta}_j - \beta_j}{\text{SE}(\hat{\beta}_j)} \sim t_{N-p-1}$$

- It remains to be shown that U and V are independent

$$U \triangleq \frac{\hat{\beta}_j - \beta_j}{\sqrt{B_j}\sigma} \sim N(0,1) \quad \text{and} \quad V \triangleq \chi_{N-p-1}^2$$

- Sufficient to show that $y - \hat{y}$ and $\hat{\beta} - \beta$ are independent

$$(\hat{\beta} - \beta)(y - \hat{y})^T = (X^T X)^{-1} X^T \varepsilon \varepsilon^T (I - H)$$

- From $E\varepsilon\varepsilon^T = \sigma^2 I$ and $HX = X$,

$$E(\hat{\beta} - \beta)(y - \hat{y})^T = 0$$

- 1 Distribution of the RSS Values
- 2 Hypothesis Testing for $\hat{\beta}_j \neq 0$
- 3 Coefficient of Determination and the Detection of Collinearity**
- 4 Confidence and Prediction Intervals

- We define a matrix $W \in \mathbb{R}^{N \times N}$ such that all the elements are $1/N$
 $Wy \in \mathbb{R}^N$ are $\bar{y} = Wy = \sum_{i=1}^N y_i$ for $y_1, \dots, y_N \in \mathbb{R}$
- Residual sum of squares RSS

$$\text{RSS} = \|\hat{y} - y\|^2 = \|(I - H)\varepsilon\|^2 = \|(I - H)y\|^2$$

- Explained sum of squares ESS

$$\text{ESS} \triangleq \|\hat{y} - \bar{y}\|^2 = \|\hat{y} - Wy\|^2 = \|(H - W)y\|^2$$

- Total sum of squares TSS

$$\text{TSS} \triangleq \|y - \bar{y}\|^2 = \|(I - W)y\|^2$$

- Coefficient of determination

$$R^2 = \frac{\text{ESS}}{\text{TSS}} = 1 - \frac{\text{RSS}}{\text{TSS}}$$

- Correlation between the covariates and response

$$\hat{\rho} \triangleq \frac{\sum_{i=1}^N (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^N (x_i - \bar{x})^2 \sum_{i=1}^N (y_i - \bar{y})^2}}$$

$$\begin{aligned} \frac{\text{ESS}}{\text{TSS}} &= \frac{\hat{\beta}_1^2 \|x - \bar{x}\|^2}{\|y - \bar{y}\|^2} = \left\{ \frac{\sum_{i=1}^N (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^N (x_i - \bar{x})^2} \right\}^2 \frac{\sum_{i=1}^N (x_i - \bar{x})^2}{\sum_{i=1}^N (y_i - \bar{y})^2} \\ &= \frac{\left\{ \sum_{i=1}^N (x_i - \bar{x})(y_i - \bar{y}) \right\}^2}{\sum_{i=1}^N (x_i - \bar{x})^2 \sum_{i=1}^N (y_i - \bar{y})^2} = \hat{\rho}^2 \end{aligned}$$

- Variance inflation factors

$$\text{VIF} \triangleq \frac{1}{1 - R_{X_j|X_{-j}}^2}$$

- The minimum value of VIF is one, and we say that the collinearity of covariate is strong when its VIF value is large

- 1 Distribution of the RSS Values
- 2 Hypothesis Testing for $\hat{\beta}_j \neq 0$
- 3 Coefficient of Determination and the Detection of Collinearity
- 4 Confidence and Prediction Intervals

- We have showed how to obtain the estimate $\hat{\beta}$ of $\beta \in \mathbb{R}^{p+1}$, confidence interval of $\hat{\beta}$ as follows

$$\beta_i = \hat{\beta}_i \pm t_{N-p-1}(\alpha/2)\text{SE}(\hat{\beta}_i), \quad \text{for } i = 0, 1, \dots, p$$

- Confidence interval of $x_*\hat{\beta}$ for another point $x_* \in \mathbb{R}^{p+1}$

- The average

$$E[x_*\hat{\beta}] = x_*E[\hat{\beta}]$$

- The variance

$$V[x_*\hat{\beta}] = x_*V(\hat{\beta})x_*^T = \sigma^2 x_*(X^T X)^{-1}x_*^T$$

- We define

$$\hat{\sigma} \triangleq \sqrt{\text{RSS}/(N-p-1)}, \quad \text{SE}(x_*\hat{\beta}) \triangleq \hat{\sigma}\sqrt{x_*(X^T X)^{-1}x_*^T}$$

Confidence and Prediction Intervals in Regression

- $C \sim t_{N-p-1}$

- variance in the difference between $x_*\hat{\beta}$ and $y_* \triangleq x_*\beta + \varepsilon$

$$V[x_*\hat{\beta} - (x_*\beta + \varepsilon)] = V[x_*(\hat{\beta} - \beta)] + V[\varepsilon] = \sigma^2 x_*(X^T X)^{-1} x_*^T + \sigma^2$$

- Similarly, we can derive the following

$$P \triangleq \frac{x_*\hat{\beta} - y_*}{\text{SE}(x_*\hat{\beta} - y_*)} = \frac{x_*\hat{\beta} - y_*}{\sigma(1 + \sqrt{x_*(X^T X)^{-1} x_*^T})} / \sqrt{\frac{\text{RSS}}{\sigma^2} / (N - p - 1)} \sim t_{N-p-1}$$

- The confidence and prediction intervals

$$\begin{aligned} x_*\beta &= x_*\hat{\beta} \pm t_{N-p-1}(\alpha/2)\hat{\sigma}\sqrt{x_*(X^T X)^{-1} x_*^T} \\ y_* &= x_*\hat{\beta} \pm t_{N-p-1}(\alpha/2)\hat{\sigma}\sqrt{1 + x_*(X^T X)^{-1} x_*^T} \end{aligned}$$

Q & A

Thank you :)