

HOMEWORK

CONTENTS

1. Homework 1 (Due: Apr 5)	2
2. Homework 2 (Due: Apr 19)	4
3. Homework 3 (Due: May 10)	7

1. HOMEWORK 1 (DUE: APR 5)

Problem 1.1. Let A be a nonempty set and let k be a positive integer with $k \leq |A|$. The symmetric group S_A acts on the set B consisting of all subsets of A of cardinality k by

$$\sigma \cdot \{a_1, \dots, a_k\} = \{\sigma(a_1), \dots, \sigma(a_k)\}.$$

- (1) Prove that this is a group action.
- (2) Describe explicitly how the elements $(1\ 2)$ and $(1\ 2\ 3)$ act on the six 2-element subsets of $\{1, 2, 3, 4\}$.

Problem 1.2. Let H be a group acting on a set A . Prove that the relation \sim on A defined by $a \sim b$ if and only if $a = hb$ for some $h \in H$ is an equivalence relation. (For each $x \in A$ the equivalence class of x under \sim is called the orbit of x under the action of H . The orbits under the action of H partition the set A .)

Problem 1.3. In each of parts (1) to (5) give the number of nonisomorphic abelian groups of the specified order - do not list the groups:

- (1) order 100
- (2) order 576
- (3) order 1155
- (4) order 42875
- (5) order 2704

Problem 1.4. In each of parts (1) to (5) give the lists of invariant factors for all abelian groups of the specified order:

- (1) order 270
- (2) order 9801
- (3) order 320
- (4) order 105
- (5) order 44100

Problem 1.5. In each of parts (1) to (5) give the lists of elementary divisors for all abelian groups of the specified order and then match each list with the corresponding list of invariant factors found in the preceding problem:

- (1) order 270
- (2) order 9801
- (3) order 320
- (4) order 105
- (5) order 44100

Problem 1.6. Let R be a ring with identity and let S be a subring of R containing the identity. Prove that if u is a unit in S then u is a unit in R . Show by example that the converse is false.

Problem 1.7. Let R be a ring with $1 \neq 0$.

- (1) Prove that if a is a zero divisor, then it is not a unit.
- (2) Prove that if $ab = ac$ and $a \neq 0$ is not a zero divisor, then $b = c$.

Problem 1.8. Assume R is commutative with $1 \neq 0$. Prove that if P is a prime ideal of R and P contains no zero divisors then R is an integral domain.

Problem 1.9. Let R be a ring with $1 \neq 0$. Let $A = (a_1, a_2, \dots, a_n)$ be a nonzero finitely generated ideal of R . Prove that there is an ideal B which is maximal with respect to the property that it does not contain A . [Use Zorn's Lemma.]

Problem 1.10. Let n_1, n_2, \dots, n_k be integers which are relatively prime in pairs: $(n_i, n_j) = 1$ for all $i \neq j$.

- (1) Show that the Chinese Remainder Theorem implies that for any $a_1, \dots, a_k \in \mathbb{Z}$ there is a solution $x \in \mathbb{Z}$ to the simultaneous congruences

$$x \equiv a_1 \pmod{n_1}, \quad x \equiv a_2 \pmod{n_2}, \quad \dots, \quad x \equiv a_k \pmod{n_k}$$

and that the solution x is unique $\pmod{n = n_1 n_2 \dots n_k}$.

- (2) Let $n'_i = n/n_i$ be the quotient of n by n_i , which is relatively prime to n_i by assumption. Let t_i be the inverse of $n'_i \pmod{n_i}$. Prove that the solution x in (a) is given by

$$x = a_1 t_1 n'_1 + a_2 t_2 n'_2 + \dots + a_k t_k n'_k \pmod{n}.$$

Note that the elements t_i can be quickly found by the Euclidean Algorithm as described in Section 2 of the Preliminaries chapter (writing $an_i + bn'_i = (n_i, n'_i) = 1$ gives $t_i = b$) and that these then quickly give the solutions to the system of congruences above for any choice of a_1, a_2, \dots, a_k .

- (3) Solve the simultaneous system of congruences

$$x \equiv 1 \pmod{8}, \quad x \equiv 2 \pmod{25}, \quad \text{and} \quad x \equiv 3 \pmod{81}$$

and the simultaneous system

$$y \equiv 5 \pmod{8}, \quad y \equiv 12 \pmod{25}, \quad \text{and} \quad y \equiv 47 \pmod{81}$$

2. HOMEWORK 2 (DUE: APR 19)

For all problems, suppose that R is a ring with $1 \neq 0$ and M is a left R -module.

Problem 2.1. An element m of the R -module M is called a *torsion element* if $rm = 0$ for some nonzero element $r \in R$. The set of torsion elements is denoted

$$\text{Tor}(M) = \{m \in M \mid rm = 0 \text{ for some nonzero } r \in R\}.$$

- (1) Prove that if R is an integral domain then $\text{Tor}(M)$ is a submodule of M (called the torsion submodule of M).
- (2) Give an example of a ring R and an R -module M such that $\text{Tor}(M)$ is not a submodule. [Consider the torsion elements in the R -module R .]
- (3) If R has zero divisors show that every nonzero R -module has nonzero torsion elements.

Problem 2.2. (1) If N is a submodule of M , the *annihilator of N in R* is defined to be $\{r \in R \mid rn = 0 \text{ for all } n \in N\}$. Prove that the annihilator of N in R is a 2-sided ideal of R .

- (2) If I is a right ideal of R , the *annihilator of I in M* is defined to be $\{m \in M \mid am = 0 \text{ for all } a \in I\}$. Prove that the annihilator of I in M is a submodule of M .
- (3) Let M be the abelian group (i.e., \mathbb{Z} -module) $\mathbb{Z}/24\mathbb{Z} \times \mathbb{Z}/15\mathbb{Z} \times \mathbb{Z}/50\mathbb{Z}$.
 - (a) Find the annihilator of M in \mathbb{Z} (i.e., a generator for this principal ideal).
 - (b) Let $I = 2\mathbb{Z}$. Describe the annihilator of I in M as a direct product of cyclic groups.

Problem 2.3. (1) Let $F = \mathbb{R}$, let $V = \mathbb{R}^2$ and let T be the linear transformation from V to V which is rotation clockwise about the origin by $\pi/2$ radians. Show that V and 0 are the only $F[x]$ -submodules for this T .

- (2) Let $F = \mathbb{R}$, let $V = \mathbb{R}^2$ and let T be the linear transformation from V to V which is projection onto the y -axis. Show that $V, 0$, the x -axis and the y -axis are the only $F[x]$ -submodules for this T .
- (3) Let $F = \mathbb{R}$, let $V = \mathbb{R}^2$ and let T be the linear transformation from V to V which is rotation clockwise about the origin by π radians. Show that every subspace of V is an $F[x]$ -submodule for this T .

Problem 2.4. (1) For any left ideal I of R define

$$IM = \left\{ \sum_{\text{finite}} a_i m_i \mid a_i \in I, m_i \in M \right\}$$

to be the collection of all finite sums of elements of the form am where $a \in I$ and $m \in M$. Prove that IM is a submodule of M .

- (2) Let A_1, A_2, \dots, A_n be R -modules and let B_i be a submodule of A_i for each $i = 1, 2, \dots, n$. Prove that

$$(A_1 \times \cdots \times A_n) / (B_1 \times \cdots \times B_n) \cong (A_1/B_1) \times \cdots \times (A_n/B_n).$$

- (3) Let I be a left ideal of R and let n be a positive integer. Prove that

$$R^n / IR^n \cong R/IR \times \cdots \times R/IR \quad (n \text{ times}).$$

- (4) Assume R is commutative. Prove that $R^n \cong R^m$ if and only if $n = m$, i.e., two free R -modules of finite rank are isomorphic if and only if they have the same rank. [Apply the previous problem with I a maximal ideal of R . You may use the fact that if F is a field, then $F^n \cong F^m$ if and only if $n = m$.]

Problem 2.5. Let I be a nonempty index set and for each $i \in I$ let M_i be an R -module. The direct product of the modules M_i is defined to be their direct product as abelian groups (cf. Exercise 15 in Section 5.1) with the action of R componentwise multiplication. The direct sum of the modules M_i is defined to be the restricted direct product of the abelian groups M_i (cf. Exercise 17 in Section 5.1) with the action of R componentwise multiplication. In other words, the direct sum of the M_i 's is the subset of the direct product, $\prod_{i \in I} M_i$, which consists of all elements $\prod_{i \in I} m_i$ such

that only finitely many of the components m_i are nonzero; the action of R on the direct product or direct sum is given by $r \prod_{i \in I} m_i = \prod_{i \in I} r m_i$ (cf. Appendix I for the definition of Cartesian products of infinitely many sets). The direct sum will be denoted by $\oplus_{i \in I} M_i$.

- (1) Prove that the direct product of the M_i 's is an R -module and the direct sum of the M_i 's is a submodule of their direct product.
- (2) Show that if $R = \mathbb{Z}$, $I = \mathbb{Z}^+$ and M_i is the cyclic group of order i for each i , then the direct sum of the M_i 's is not isomorphic to their direct product. [Look at torsion.]

Problem 2.6. (1) Show that the element " $2 \otimes 1$ " is 0 in $\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$ but is nonzero in $2\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$.

- (2) Show that $\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C}$ and $\mathbb{C} \otimes_{\mathbb{C}} \mathbb{C}$ are both left \mathbb{R} -modules but are not isomorphic as \mathbb{R} -modules.
- (3) Show that $\mathbb{Q} \otimes_{\mathbb{Z}} \mathbb{Q}$ and $\mathbb{Q} \otimes_{\mathbb{Q}} \mathbb{Q}$ are isomorphic left \mathbb{Q} -modules. [Show they are both 1-dimensional vector spaces over \mathbb{Q} .]
- (4) If R is any integral domain with quotient field Q , prove that $(Q/R) \otimes_R (Q/R) = 0$.
- (5) Let $\{e_1, e_2\}$ be a basis of $V = \mathbb{R}^2$. Show that the element $e_1 \otimes e_2 + e_2 \otimes e_1$ in $V \otimes_{\mathbb{R}} V$ cannot be written as a simple tensor $v \otimes w$ for any $v, w \in \mathbb{R}^2$.

Problem 2.7. Suppose R is commutative and $N \cong R^n$ is a free R -module of rank n with R -module basis e_1, \dots, e_n .

- (1) For any nonzero R -module M show that every element of $M \otimes N$ can be written uniquely in the form $\sum_{i=1}^n m_i \otimes e_i$ where $m_i \in M$. Deduce that if $\sum_{i=1}^n m_i \otimes e_i = 0$ in $M \otimes N$ then $m_i = 0$ for $i = 1, \dots, n$.
- (2) Show that if $\sum m_i \otimes n_i = 0$ in $M \otimes N$ where the n_i are merely assumed to be R linearly independent then it is not necessarily true that all the m_i are 0. [Consider $R = \mathbb{Z}$, $n = 1$, $M = \mathbb{Z}/2\mathbb{Z}$, and the element $1 \otimes 2$.

Problem 2.8. Suppose that

$$\begin{array}{ccccc} A & \xrightarrow{\psi} & B & \xrightarrow{\varphi} & C \\ \alpha \downarrow & & \beta \downarrow & & \gamma \downarrow \\ A' & \xrightarrow{\psi'} & B' & \xrightarrow{\varphi'} & C' \end{array}$$

is a commutative diagram of groups and that the rows are exact. Prove that

- (1) if φ and α are surjective, and β is injective then γ is injective. [If $c \in \ker \gamma$, show there is a $b \in B$ with $\varphi(b) = c$. Show that $\varphi'(\beta(b)) = 0$ and deduce that $\beta(b) = \psi'(a')$ for some $a' \in A'$. Show there is an $a \in A$ with $\alpha(a) = a'$ and that $\beta(\psi(a)) = \beta(b)$. Conclude that $b = \psi(a)$ and hence $c = \varphi(b) = 0$.]
- (2) if ψ' , α , and γ are injective, then β is injective,
- (3) if φ , α , and γ are surjective, then β is surjective,
- (4) if β is injective, α and φ are surjective, then γ is injective,
- (5) if β is surjective, γ and ψ' are injective, then α is surjective.

Problem 2.9. Let P_1 and P_2 be R -modules. Prove that $P_1 \oplus P_2$ is a projective R -module if and only if both P_1 and P_2 are projective.

Problem 2.10. Let $0 \longrightarrow L \xrightarrow{\psi} M \xrightarrow{\varphi} N \longrightarrow 0$ be a sequence of R -modules.

- (1) Prove that the associated sequence

$$0 \longrightarrow \text{Hom}_R(D, L) \xrightarrow{\psi'} \text{Hom}_R(D, M) \xrightarrow{\varphi'} \text{Hom}_R(D, N) \longrightarrow 0$$

is a short exact sequence of abelian groups for all R -modules D if and only if the original sequence is a split short exact sequence. [To show the sequence splits, take $D = N$ and show the lift of the identity map in $\text{Hom}_R(N, N)$ to $\text{Hom}_R(N, M)$ is a splitting homomorphism for φ .]

(2) Prove that the associated sequence

$$0 \longrightarrow \operatorname{Hom}_R(N, D) \xrightarrow{\varphi'} \operatorname{Hom}_R(M, D) \xrightarrow{\psi'} \operatorname{Hom}_R(L, D) \longrightarrow 0$$

is a short exact sequence of abelian groups for all R -modules D if and only if the original sequence is a split short exact sequence.

3. HOMEWORK 3 (DUE: MAY 10)

Problem 3.1 (Section 12.1, Exercise 2). Let M be a module over the integral domain R .

- (1) Suppose that M has rank n and that x_1, x_2, \dots, x_n is any maximal set of linearly independent elements of M . Let $N = Rx_1 + \dots + Rx_n$ be the submodule generated by x_1, x_2, \dots, x_n . Prove that N is isomorphic to R^n and that the quotient M/N is a torsion R -module (equivalently, the elements x_1, \dots, x_n are linearly independent and for any $y \in M$ there is a nonzero element $r \in R$ such that ry can be written as a linear combination $r_1x_1 + \dots + r_nx_n$ of the x_i).
- (2) Prove conversely that if M contains a submodule N that is free of rank n (i.e., $N \cong R^n$) such that the quotient M/N is a torsion R -module then M has rank n . [Let y_1, y_2, \dots, y_{n+1} be any $n+1$ elements of M . Use the fact that M/N is torsion to write ry_i as a linear combination of a basis for N for some nonzero elements r_1, \dots, r_{n+1} of R . Use an argument as in the proof of Proposition 3 to see that the ry_i , and hence also the y_i , are linearly dependent.]

Problem 3.2 (Section 12.1, Exercise 3). Let R be an integral domain and let A and B be R -modules of ranks m and n , respectively. Prove that the rank of $A \oplus B$ is $m+n$. [Use the previous exercise.]

Problem 3.3 (Section 12.1, Exercise 4). Let R be an integral domain, let M be an R -module and let N be a submodule of M . Suppose M has rank n , N has rank r and the quotient M/N has rank s . Prove that $n = r + s$. [Let x_1, x_2, \dots, x_s be elements of M whose images in M/N are a maximal set of independent elements and let $x_{s+1}, x_{s+2}, \dots, x_{s+r}$ be a maximal set of independent elements in N . Prove that x_1, x_2, \dots, x_{s+r} are linearly independent in M and that for any element $y \in M$ there is a nonzero element $r \in R$ such that ry is a linear combination of these elements. Then use Exercise 2.]

Problem 3.4 (Section 12.1, Exercise 5). Let $R = \mathbb{Z}[x]$ and let $M = (2, x)$ be the ideal generated by 2 and x , considered as a submodule of R . Show that $\{2, x\}$ is not a basis of M . [Find a nontrivial R -linear dependence between these two elements.] Show that the rank of M is 1 but that M is not free of rank 1 (cf. Exercise 2).

Problem 3.5 (Section 12.1, Exercise 6). Show that if R is an integral domain and M is any nonprincipal ideal of R then M is torsion free of rank 1 but is not a free R -module.

Problem 3.6 (Section 12.1, Exercise 7). Let R be any ring, let A_1, A_2, \dots, A_m be R -modules and let B_i be a submodule of A_i , $1 \leq i \leq m$. Prove that

$$(A_1 \oplus A_2 \oplus \dots \oplus A_m) / (B_1 \oplus B_2 \oplus \dots \oplus B_m) \cong (A_1/B_1) \oplus (A_2/B_2) \oplus \dots \oplus (A_m/B_m)$$

Problem 3.7 (Section 12.1, Exercise 8). Let R be a P.I.D., let B be a torsion R -module and let p be a prime in R . Prove that if $pb = 0$ for some nonzero $b \in B$, then $\text{Ann}(B) \subseteq (p)$.

Problem 3.8 (Section 12.1, Exercise 9). Give an example of an integral domain R and a nonzero torsion R -module M such that $\text{Ann}(M) = 0$. Prove that if N is a finitely generated torsion R -module then $\text{Ann}(N) \neq 0$.

Problem 3.9 (Section 12.1, Exercise 10). For p a prime in the P.I.D. R and N an R -module prove that the p -primary component of N is a submodule of N and prove that N is the direct sum of its p -primary components (there need not be finitely many of them).

Problem 3.10 (Section 12.1, Exercise 15). Prove that if R is a Noetherian ring then R^n is a Noetherian R -module. [Fix a basis of R^n . If M is a submodule of R^n show that the collection of first coordinates of elements of M is a submodule of R hence is finitely generated. Let m_1, m_2, \dots, m_k be elements of M whose first coordinates generate this submodule of R . Show that any element of M can be written as an R -linear combination of m_1, m_2, \dots, m_k plus an element of M whose first coordinate is 0. Prove that $M \cap R^{n-1}$ is a submodule of R^{n-1} where R^{n-1} is the set of elements of R^n with first coordinate 0 and then use induction on n .