

# Enumeration of multiplex juggling card sequences using generalized $q$ -derivatives

Jang Soo Kim  
Sungkyunkwan University

Joint work with Yumin Cho, Jaehyun Kim, and Nakyung Lee  
(Gyeonggi Science High School for the Gifted)

GIST

May 10, 2024

# What is juggling?

- **Juggling** is an act of manipulating objects in some interesting ways.

# What is juggling?

- **Juggling** is an act of manipulating objects in some interesting ways.
- **Toss juggling** is an act of throwing and catching objects.



# What is juggling?

- **Juggling** is an act of manipulating objects in some interesting ways.
- **Toss juggling** is an act of throwing and catching objects.



- **Contact juggling** is an act of manipulating objects in contact with the body.



# What is juggling?

- **Juggling** is an act of manipulating objects in some interesting ways.
- **Toss juggling** is an act of throwing and catching objects.



- **Contact juggling** is an act of manipulating objects in contact with the body.



# What is juggling?

- **Juggling** is an act of manipulating objects in some interesting ways.
- **Toss juggling** is an act of throwing and catching objects.



- **Contact juggling** is an act of manipulating objects in contact with the body.



## How to model juggling mathematically?

- We assume that every throw and catch is performed at an integer beat.

## How to model juggling mathematically?

- We assume that every throw and catch is performed at an integer beat.
- Every throw can be represented as a nonnegative integer  $k$ .

## How to model juggling mathematically?


- We assume that every throw and catch is performed at an integer beat.
- Every throw can be represented as a nonnegative integer  $k$ .
- A  **$k$ -throw** means throwing a ball at beat  $i$  that will land at beat  $i + k$ .

## How to model juggling mathematically?

- We assume that every throw and catch is performed at an integer beat.
- Every throw can be represented as a nonnegative integer  $k$ .
- A  **$k$ -throw** means throwing a ball at beat  $i$  that will land at beat  $i + k$ .
- By convention a **0-throw** means nothing happens at that beat.

## How to model juggling mathematically?

- We assume that every throw and catch is performed at an integer beat.
- Every throw can be represented as a nonnegative integer  $k$ .
- A  **$k$ -throw** means throwing a ball at beat  $i$  that will land at beat  $i + k$ .
- By convention a **0-throw** means nothing happens at that beat.
- For example,



## Simple juggling

- For **simple juggling**, we assume that at every beat the juggler can throw at most one ball and catch at most one ball.

## Simple juggling

- For **simple juggling**, we assume that at every beat the juggler can throw at most one ball and catch at most one ball.
- A **(simple) juggling sequence** is a sequence  $a_1 a_2 \dots a_n$  such that if we draw the diagram with this sequence then at every beat

## Simple juggling

- For **simple juggling**, we assume that at every beat the juggler can throw at most one ball and catch at most one ball.
- A **(simple) juggling sequence** is a sequence  $a_1 a_2 \dots a_n$  such that if we draw the diagram with this sequence then at every beat
  - there is at most one ball in,

## Simple juggling


- For **simple juggling**, we assume that at every beat the juggler can throw at most one ball and catch at most one ball.
- A **(simple) juggling sequence** is a sequence  $a_1 a_2 \dots a_n$  such that if we draw the diagram with this sequence then at every beat
  - there is at most one ball in,
  - there is at most one ball out,

## Simple juggling

- For **simple juggling**, we assume that at every beat the juggler can throw at most one ball and catch at most one ball.
- A **(simple) juggling sequence** is a sequence  $a_1 a_2 \dots a_n$  such that if we draw the diagram with this sequence then at every beat
  - there is at most one ball in,
  - there is at most one ball out,
  - there is a ball in if and only if there is a ball out.


## Simple juggling

- For **simple juggling**, we assume that at every beat the juggler can throw at most one ball and catch at most one ball.
- A **(simple) juggling sequence** is a sequence  $a_1a_2 \dots a_n$  such that if we draw the diagram with this sequence then at every beat
  - ① there is at most one ball in,
  - ② there is at most one ball out,
  - ③ there is a ball in if and only if there is a ball out.
- For example,



## Simple juggling


- For **simple juggling**, we assume that at every beat the juggler can throw at most one ball and catch at most one ball.
- A **(simple) juggling sequence** is a sequence  $a_1 a_2 \dots a_n$  such that if we draw the diagram with this sequence then at every beat
  - there is at most one ball in,
  - there is at most one ball out,
  - there is a ball in if and only if there is a ball out.
- For example,



- The number of balls for a juggling sequence  $a_1 \dots a_n$  is the average  $\frac{a_1 + \dots + a_n}{n}$ .


## Enumeration of simple juggling patterns

- **Juggling cards** tell us what happens at each beat:




## Enumeration of simple juggling patterns

- **Juggling cards** tell us what happens at each beat:




- The pattern 4053 can be understood as follows.




## Enumeration of simple juggling patterns

- **Juggling cards** tell us what happens at each beat:




- The pattern 4053 can be understood as follows.




- The following theorem was first proved by Buhler, Eisenbud, Graham, Wright in 1994 using permutations and later by Ehrenborg and Readdy in 1996 using juggling cards.

## Enumeration of simple juggling patterns

- **Juggling cards** tell us what happens at each beat:



- The pattern 4053 can be understood as follows.




- The following theorem was first proved by Buhler, Eisenbud, Graham, Wright in 1994 using permutations and later by Ehrenborg and Readdy in 1996 using juggling cards.


### Theorem

## Enumeration of simple juggling patterns

- **Juggling cards** tell us what happens at each beat:



- The pattern 4053 can be understood as follows.




- The following theorem was first proved by Buhler, Eisenbud, Graham, Wright in 1994 using permutations and later by Ehrenborg and Readdy in 1996 using juggling cards.

### Theorem


- *The number of juggling patterns with at most  $b$  balls and length  $n$  is  $(b + 1)^n$ .*

## Enumeration of simple juggling patterns

- **Juggling cards** tell us what happens at each beat:



- The pattern 4053 can be understood as follows.




- The following theorem was first proved by Buhler, Eisenbud, Graham, Wright in 1994 using permutations and later by Ehrenborg and Readdy in 1996 using juggling cards.

### Theorem


- *The number of juggling patterns with at most  $b$  balls and length  $n$  is  $(b + 1)^n$ .*
- *The number of juggling patterns with  $b$  balls and length  $n$  is  $(b + 1)^n - b^n$ .*

## Enumeration of simple juggling patterns

- **Juggling cards** tell us what happens at each beat:



- The pattern 4053 can be understood as follows.



- The following theorem was first proved by Buhler, Eisenbud, Graham, Wright in 1994 using permutations and later by Ehrenborg and Readdy in 1996 using juggling cards.

### Theorem

- *The number of juggling patterns with at most  $b$  balls and length  $n$  is  $(b + 1)^n$ .*
- *The number of juggling patterns with  $b$  balls and length  $n$  is  $(b + 1)^n - b^n$ .*
- *The number of juggling patterns with  $b$  balls and smallest period  $n$  is*

$$\frac{1}{n} \sum_{d|n} \mu\left(\frac{n}{d}\right) ((b+1)^d - b^d).$$

## Multiplex juggling


- Recall: For **simple juggling**, we assume that at every beat the juggler can throw at most one ball and catch at most one ball.

## Multiplex juggling

- Recall: For **simple juggling**, we assume that at every beat the juggler can throw at most one ball and catch at most one ball.
- A **multiplex juggling sequence** is a sequence  $A_1A_2 \cdots A_n$  of sets  $A_i = \{a_{i,1}, a_{i,2}, \dots, a_{i,t_i}\}$  such that if we draw the diagram with this sequence then at every beat there are the same number of balls in and balls out.


## Multiplex juggling

- Recall: For **simple juggling**, we assume that at every beat the juggler can throw at most one ball and catch at most one ball.
- A **multiplex juggling sequence** is a sequence  $A_1A_2 \cdots A_n$  of sets  $A_i = \{a_{i,1}, a_{i,2}, \dots, a_{i,t_i}\}$  such that if we draw the diagram with this sequence then at every beat there are the same number of balls in and balls out.
- For example, the multiplex juggling sequence  $\{1, 4\}, \{1\}$  is



## Multiplex juggling


- Recall: For **simple juggling**, we assume that at every beat the juggler can throw at most one ball and catch at most one ball.
- A **multiplex juggling sequence** is a sequence  $A_1 A_2 \cdots A_n$  of sets  $A_i = \{a_{i,1}, a_{i,2}, \dots, a_{i,t_i}\}$  such that if we draw the diagram with this sequence then at every beat there are the same number of balls in and balls out.
- For example, the multiplex juggling sequence  $\{1, 4\}, \{1\}$  is



- If every  $A_i$  has cardinality at most  $k$ , we say that the juggling sequence has **hand capacity  $k$** .

## Multiplex juggling


- Recall: For **simple juggling**, we assume that at every beat the juggler can throw at most one ball and catch at most one ball.
- A **multiplex juggling sequence** is a sequence  $A_1 A_2 \cdots A_n$  of sets  $A_i = \{a_{i,1}, a_{i,2}, \dots, a_{i,t_i}\}$  such that if we draw the diagram with this sequence then at every beat there are the same number of balls in and balls out.
- For example, the multiplex juggling sequence  $\{1, 4\}, \{1\}$  is



- If every  $A_i$  has cardinality at most  $k$ , we say that the juggling sequence has **hand capacity  $k$** .
- The number of balls for a multiplex juggling sequence  $A_1 \cdots A_n$  is the average  $\frac{\|A_1\| + \cdots + \|A_n\|}{n}$ , where  $\|A_i\|$  is the sum of elements in  $A_i$ .


## Enumeration of multiplex juggling patterns

- We can try to use the juggling card technique.




## Enumeration of multiplex juggling patterns

- We can try to use the juggling card technique.




- For example, the pattern  $\{1, 2\}, \{5\}, \{1\}$  is represented as




## Enumeration of multiplex juggling patterns


- We can try to use the juggling card technique.



- For example, the pattern  $\{1, 2\}, \{5\}, \{1\}$  is represented as




- However, the same pattern has a different representation:




## Enumeration of multiplex juggling patterns


- We can try to use the juggling card technique.



- For example, the pattern  $\{1, 2\}, \{5\}, \{1\}$  is represented as




- However, the same pattern has a different representation:



- All we can say is that the number of multiplex juggling patterns with at most  $b$  balls and length  $n$  is **at most**  $(\binom{b}{0} + \binom{b}{1} + \dots + \binom{b}{b})^n = (2^b)^n = 2^{bn}$ .


## Multiplex juggling cards

- In 2019, Butler, Choi, Kim, and Seo introduced a different type of multiplex juggling cards.




## Multiplex juggling cards

- In 2019, Butler, Choi, Kim, and Seo introduced a different type of multiplex juggling cards.




- This card scheme represents a juggling pattern uniquely.




## Multiplex juggling cards

- In 2019, Butler, Choi, Kim, and Seo introduced a different type of multiplex juggling cards.



- This card scheme represents a juggling pattern uniquely.



- However, the cards cannot be arranged independently.

## Three natural statistics

- **Goal:** study enumerative properties of the number of multiplex juggling card sequences.

## Three natural statistics

- **Goal:** study enumerative properties of the number of multiplex juggling card sequences.
- We consider three natural parameters of multiplex juggling card sequences: the **number  $b$  of balls**, the **capacity  $k$** , and the **length  $\ell$**  of a sequence of cards.

## Three natural statistics

- **Goal:** study enumerative properties of the number of multiplex juggling card sequences.
- We consider three natural parameters of multiplex juggling card sequences: the **number  $b$  of balls**, the **capacity  $k$** , and the **length  $\ell$**  of a sequence of cards.
- Let  $J(b, k, \ell)$  denote the number of multiplex juggling card sequences with given parameters  $k$ ,  $b$ , and  $\ell$ .

## Three natural statistics

- **Goal:** study enumerative properties of the number of multiplex juggling card sequences.
- We consider three natural parameters of multiplex juggling card sequences: the **number  $b$  of balls**, the **capacity  $k$** , and the **length  $\ell$**  of a sequence of cards.
- Let  $J(b, k, \ell)$  denote the number of multiplex juggling card sequences with given parameters  $k$ ,  $b$ , and  $\ell$ .
- Note that  $J(b, k, 1)$  is the number of multiplex juggling cards with  $b$  balls and capacity  $k$ .

## The number of multiplex juggling cards

- Let's consider

$$\sum_{b \geq 0} J(b, k, 1)x^b.$$

## The number of multiplex juggling cards

- Let's consider

$$\sum_{b \geq 0} J(b, k, 1)x^b.$$

- If  $k = 1$ , by definition,

$$\sum_{b \geq 0} J(b, 1, 1)x^b = \sum_{b \geq 0} (b + 1)x^b = \frac{1}{(1 - x)^2}.$$

# The number of multiplex juggling cards

- Let's consider

$$\sum_{b \geq 0} J(b, k, 1)x^b.$$

- If  $k = 1$ , by definition,

$$\sum_{b \geq 0} J(b, 1, 1)x^b = \sum_{b \geq 0} (b + 1)x^b = \frac{1}{(1 - x)^2}.$$

- For the other extreme case  $k = \infty$ , Butler et al. (2019) showed that the sequence  $\{J(b, \infty, 1)\}_{b \geq 0}$  satisfies the following simple linear recurrence:  $J(0, \infty, 1) = 1$ ,  $J(1, \infty, 1) = 2$ ,  $J(2, \infty, 1) = 7$ , and for  $b \geq 3$ ,

$$J(b, \infty, 1) = 4J(b - 1, \infty, 1) - 2J(b - 2, \infty, 1).$$

# The number of multiplex juggling cards

- Let's consider

$$\sum_{b \geq 0} J(b, k, 1)x^b.$$

- If  $k = 1$ , by definition,

$$\sum_{b \geq 0} J(b, 1, 1)x^b = \sum_{b \geq 0} (b + 1)x^b = \frac{1}{(1 - x)^2}.$$

- For the other extreme case  $k = \infty$ , Butler et al. (2019) showed that the sequence  $\{J(b, \infty, 1)\}_{b \geq 0}$  satisfies the following simple linear recurrence:  $J(0, \infty, 1) = 1$ ,  $J(1, \infty, 1) = 2$ ,  $J(2, \infty, 1) = 7$ , and for  $b \geq 3$ ,

$$J(b, \infty, 1) = 4J(b - 1, \infty, 1) - 2J(b - 2, \infty, 1).$$

- By the standard method for linear recurrences, this is equivalent to

$$\sum_{b \geq 0} J(b, \infty, 1)x^b = \frac{1 - 2x + x^2}{1 - 4x + 2x^2}.$$

# The number of multiplex juggling cards

- Let's consider

$$\sum_{b \geq 0} J(b, k, 1)x^b.$$

- If  $k = 1$ , by definition,

$$\sum_{b \geq 0} J(b, 1, 1)x^b = \sum_{b \geq 0} (b + 1)x^b = \frac{1}{(1 - x)^2}.$$

- For the other extreme case  $k = \infty$ , Butler et al. (2019) showed that the sequence  $\{J(b, \infty, 1)\}_{b \geq 0}$  satisfies the following simple linear recurrence:  $J(0, \infty, 1) = 1$ ,  $J(1, \infty, 1) = 2$ ,  $J(2, \infty, 1) = 7$ , and for  $b \geq 3$ ,

$$J(b, \infty, 1) = 4J(b - 1, \infty, 1) - 2J(b - 2, \infty, 1).$$

- By the standard method for linear recurrences, this is equivalent to

$$\sum_{b \geq 0} J(b, \infty, 1)x^b = \frac{1 - 2x + x^2}{1 - 4x + 2x^2}.$$

- For the case  $k = 2$ , Butler et al. (2019) conjectured the following generating function formula:

$$\sum_{b \geq 0} J(b, 2, 1)x^b = \frac{1 - x + x^2 + x^3}{(1 - x - x^2)^3}.$$

## Our results

- We prove Butler et al.'s conjecture:

$$\sum_{b \geq 0} J(b, 2, 1)x^b = \frac{1 - x + x^2 + x^3}{(1 - x - x^2)^3}.$$

## Our results

- We prove Butler et al.'s conjecture:

$$\sum_{b \geq 0} J(b, 2, 1)x^b = \frac{1 - x + x^2 + x^3}{(1 - x - x^2)^3}.$$

- More generally, for any capacity  $k$ , we find an explicit formula for the generating function

$$\sum_{b \geq 0} J(b, k, 1)x^b.$$

## Our results

- We prove Butler et al.'s conjecture:

$$\sum_{b \geq 0} J(b, 2, 1)x^b = \frac{1 - x + x^2 + x^3}{(1 - x - x^2)^3}.$$

- More generally, for any capacity  $k$ , we find an explicit formula for the generating function

$$\sum_{b \geq 0} J(b, k, 1)x^b.$$

- We also find an expression for

$$\sum_{b \geq 0} J(b, k, \ell)x^b \tag{1}$$

for any capacity  $k$  and length  $\ell$  by introducing a generalization of the  $q$ -derivative operator.

## Our results

- We prove Butler et al.'s conjecture:

$$\sum_{b \geq 0} J(b, 2, 1)x^b = \frac{1 - x + x^2 + x^3}{(1 - x - x^2)^3}.$$

- More generally, for any capacity  $k$ , we find an explicit formula for the generating function

$$\sum_{b \geq 0} J(b, k, 1)x^b.$$

- We also find an expression for

$$\sum_{b \geq 0} J(b, k, \ell)x^b \tag{1}$$

for any capacity  $k$  and length  $\ell$  by introducing a generalization of the  $q$ -derivative operator.

- As a consequence, we show that (1) is a rational function in  $x$ .

## Definition

## Definition

- A  **$(b, k)$ -embedding** is a sequence  $\gamma = (\gamma_1, \dots, \gamma_r)$  satisfying the following conditions:

## Definition

- A  **$(b, k)$ -embedding** is a sequence  $\gamma = (\gamma_1, \dots, \gamma_r)$  satisfying the following conditions:
  - Each  $\gamma_i$  is a word of the form  $\gamma_i = 0^u 1^v$  for some integers  $u, v \geq 0$  with  $1 \leq u + v \leq k$ .

## Definition

- A  **$(b, k)$ -embedding** is a sequence  $\gamma = (\gamma_1, \dots, \gamma_r)$  satisfying the following conditions:
  - Each  $\gamma_i$  is a word of the form  $\gamma_i = 0^u 1^v$  for some integers  $u, v \geq 0$  with  $1 \leq u + v \leq k$ .
  - The total number of 1's in  $\gamma_1, \dots, \gamma_r$  is at most  $k$ .

## Definition

- A  **$(b, k)$ -embedding** is a sequence  $\gamma = (\gamma_1, \dots, \gamma_r)$  satisfying the following conditions:
  - Each  $\gamma_i$  is a word of the form  $\gamma_i = 0^u 1^v$  for some integers  $u, v \geq 0$  with  $1 \leq u + v \leq k$ .
  - The total number of 1's in  $\gamma_1, \dots, \gamma_r$  is at most  $k$ .
  - The sum of the length of  $\gamma_i$  for all  $i \in [r]$  is equal to  $b$ .

## Definition



- A  **$(b, k)$ -embedding** is a sequence  $\gamma = (\gamma_1, \dots, \gamma_r)$  satisfying the following conditions:
  - Each  $\gamma_i$  is a word of the form  $\gamma_i = 0^u 1^v$  for some integers  $u, v \geq 0$  with  $1 \leq u + v \leq k$ .
  - The total number of 1's in  $\gamma_1, \dots, \gamma_r$  is at most  $k$ .
  - The sum of the length of  $\gamma_i$  for all  $i \in [r]$  is equal to  $b$ .
- Let  $\mathcal{E}(b, k)$  be the set of  $(b, k)$ -embeddings.

## Definition

- A  **$(b, k)$ -embedding** is a sequence  $\gamma = (\gamma_1, \dots, \gamma_r)$  satisfying the following conditions:
  - Each  $\gamma_i$  is a word of the form  $\gamma_i = 0^u 1^v$  for some integers  $u, v \geq 0$  with  $1 \leq u + v \leq k$ .
  - The total number of 1's in  $\gamma_1, \dots, \gamma_r$  is at most  $k$ .
  - The sum of the length of  $\gamma_i$  for all  $i \in [r]$  is equal to  $b$ .
- Let  $\mathcal{E}(b, k)$  be the set of  $(b, k)$ -embeddings.

## Example

The left card corresponds to the embedding  $(0000, 00, 000)$ . The right card corresponds to the embedding  $(011, 1, 00, 001, 11)$ .



# The generating function for the number of cards

## Proposition

For a positive integer  $k$ , we have

$$\sum_{b \geq 0} J(b, k, 1) x^b = [z^k] \left( \frac{1}{1-z} \cdot \frac{1}{1 - \sum_{i=1}^k x^i \sum_{j=0}^i z^j} \right).$$

Here, for a formal power series  $F(z)$  in  $z$ , the notation  $[z^k]F(z)$  means the coefficient of  $z^k$  in  $F(z)$ .

# The generating function for the number of cards

## Proposition

For a positive integer  $k$ , we have

$$\sum_{b \geq 0} J(b, k, 1)x^b = [z^k] \left( \frac{1}{1-z} \cdot \frac{1}{1 - \sum_{i=1}^k x^i \sum_{j=0}^i z^j} \right).$$

Here, for a formal power series  $F(z)$  in  $z$ , the notation  $[z^k]F(z)$  means the coefficient of  $z^k$  in  $F(z)$ .

Proof.

# The generating function for the number of cards

## Proposition

For a positive integer  $k$ , we have

$$\sum_{b \geq 0} J(b, k, 1) x^b = [z^k] \left( \frac{1}{1-z} \cdot \frac{1}{1 - \sum_{i=1}^k x^i \sum_{j=0}^i z^j} \right).$$

Here, for a formal power series  $F(z)$  in  $z$ , the notation  $[z^k]F(z)$  means the coefficient of  $z^k$  in  $F(z)$ .

## Proof.

- Since  $J(0, k, 1) = 1$  and  $J(b, k, 1) = |\mathcal{E}(b, k)|$  for  $b \geq 1$ , we can instead consider the  $(b, k)$ -embeddings for all  $b \geq 1$ .

# The generating function for the number of cards

## Proposition

For a positive integer  $k$ , we have

$$\sum_{b \geq 0} J(b, k, 1)x^b = [z^k] \left( \frac{1}{1-z} \cdot \frac{1}{1 - \sum_{i=1}^k x^i \sum_{j=0}^i z^j} \right).$$

Here, for a formal power series  $F(z)$  in  $z$ , the notation  $[z^k]F(z)$  means the coefficient of  $z^k$  in  $F(z)$ .

## Proof.

- Since  $J(0, k, 1) = 1$  and  $J(b, k, 1) = |\mathcal{E}(b, k)|$  for  $b \geq 1$ , we can instead consider the  $(b, k)$ -embeddings for all  $b \geq 1$ .
- Recall that if  $\gamma = (\gamma_1, \dots, \gamma_r)$  is a  $(b, k)$ -embedding, then  $\gamma_i = 0^{u_i}1^{v_i}$  for some integers  $u_i, v_i \geq 0$  with  $1 \leq u_i + v_i \leq k$  such that  $0 \leq v_1 + \dots + v_r \leq k$  and  $u_1 + \dots + u_r + v_1 + \dots + v_t = b$ .

# The generating function for the number of cards

## Proposition

For a positive integer  $k$ , we have

$$\sum_{b \geq 0} J(b, k, 1)x^b = [z^k] \left( \frac{1}{1-z} \cdot \frac{1}{1 - \sum_{i=1}^k x^i \sum_{j=0}^i z^j} \right).$$

Here, for a formal power series  $F(z)$  in  $z$ , the notation  $[z^k]F(z)$  means the coefficient of  $z^k$  in  $F(z)$ .

## Proof.

- Since  $J(0, k, 1) = 1$  and  $J(b, k, 1) = |\mathcal{E}(b, k)|$  for  $b \geq 1$ , we can instead consider the  $(b, k)$ -embeddings for all  $b \geq 1$ .
- Recall that if  $\gamma = (\gamma_1, \dots, \gamma_r)$  is a  $(b, k)$ -embedding, then  $\gamma_i = 0^{u_i}1^{v_i}$  for some integers  $u_i, v_i \geq 0$  with  $1 \leq u_i + v_i \leq k$  such that  $0 \leq v_1 + \dots + v_r \leq k$  and  $u_1 + \dots + u_r + v_1 + \dots + v_t = b$ .
- Let  $W$  be the set of words  $0^u1^v$  such that  $u, v \geq 0$  and  $1 \leq u + v \leq k$ .

# The generating function for the number of cards

## Proposition

For a positive integer  $k$ , we have

$$\sum_{b \geq 0} J(b, k, 1) x^b = [z^k] \left( \frac{1}{1-z} \cdot \frac{1}{1 - \sum_{i=1}^k x^i \sum_{j=0}^i z^j} \right).$$

Here, for a formal power series  $F(z)$  in  $z$ , the notation  $[z^k]F(z)$  means the coefficient of  $z^k$  in  $F(z)$ .

## Proof.

- Since  $J(0, k, 1) = 1$  and  $J(b, k, 1) = |\mathcal{E}(b, k)|$  for  $b \geq 1$ , we can instead consider the  $(b, k)$ -embeddings for all  $b \geq 1$ .
- Recall that if  $\gamma = (\gamma_1, \dots, \gamma_r)$  is a  $(b, k)$ -embedding, then  $\gamma_i = 0^{u_i} 1^{v_i}$  for some integers  $u_i, v_i \geq 0$  with  $1 \leq u_i + v_i \leq k$  such that  $0 \leq v_1 + \dots + v_r \leq k$  and  $u_1 + \dots + u_r + v_1 + \dots + v_t = b$ .
- Let  $W$  be the set of words  $0^u 1^v$  such that  $u, v \geq 0$  and  $1 \leq u + v \leq k$ .
- For a word  $w = 0^u 1^v \in W$ , let  $\ell(w) = u + v$  and  $\ell_1(w) = v$ . By definition,

$$\sum_{w \in W} x^{\ell(w)} z^{\ell_1(w)} = \sum_{i=1}^k x^i \sum_{j=0}^i z^j.$$

## Proof continued

- Thus

$$\frac{1}{1 - \sum_{i=1}^k x^i \sum_{j=0}^i z^j} = 1 + \sum_{r \geq 1} \sum_{(w_1, \dots, w_r) \in W^r} x^{\ell(w_1) + \dots + \ell(w_r)} z^{\ell_1(w_1) + \dots + \ell_1(w_r)}. \quad (2)$$

## Proof continued

- Thus

$$\frac{1}{1 - \sum_{i=1}^k x^i \sum_{j=0}^i z^j} = 1 + \sum_{r \geq 1} \sum_{(w_1, \dots, w_r) \in W^r} x^{\ell(w_1) + \dots + \ell(w_r)} z^{\ell_1(w_1) + \dots + \ell_1(w_r)}. \quad (2)$$

- On the other hand, for  $b \geq 1$ ,  $\mathcal{E}(b, k)$  is the set of  $\gamma = (\gamma_1, \dots, \gamma_r) \in W^r$ ,  $r \geq 1$ , such that  $\ell(\gamma_1) + \dots + \ell(\gamma_r) = b$  and  $0 \leq \ell_1(\gamma_1) + \dots + \ell_1(\gamma_r) \leq k$ .

## Proof continued

- Thus

$$\frac{1}{1 - \sum_{i=1}^k x^i \sum_{j=0}^i z^j} = 1 + \sum_{r \geq 1} \sum_{(w_1, \dots, w_r) \in W^r} x^{\ell(w_1) + \dots + \ell(w_r)} z^{\ell_1(w_1) + \dots + \ell_1(w_r)}. \quad (2)$$

- On the other hand, for  $b \geq 1$ ,  $\mathcal{E}(b, k)$  is the set of  $\gamma = (\gamma_1, \dots, \gamma_r) \in W^r$ ,  $r \geq 1$ , such that  $\ell(\gamma_1) + \dots + \ell(\gamma_r) = b$  and  $0 \leq \ell_1(\gamma_1) + \dots + \ell_1(\gamma_r) \leq k$ .
- Thus

$$\begin{aligned} \sum_{b \geq 0} J(b, k, 1) x^b &= 1 + \sum_{b \geq 1} |\mathcal{E}(b, k)| x^b \\ &= 1 + \sum_{r \geq 1} \sum_{\substack{(w_1, \dots, w_r) \in W^r \\ 0 \leq \ell_1(w_1) + \dots + \ell_1(w_r) \leq k}} x^{\ell(w_1) + \dots + \ell(w_r)} \\ &= \sum_{s=0}^k [z^s] \left( 1 + \sum_{r \geq 1} \sum_{(w_1, \dots, w_r) \in W^r} x^{\ell(w_1) + \dots + \ell(w_r)} z^{\ell_1(w_1) + \dots + \ell_1(w_r)} \right). \end{aligned}$$

## Proof continued

- Thus

$$\frac{1}{1 - \sum_{i=1}^k x^i \sum_{j=0}^i z^j} = 1 + \sum_{r \geq 1} \sum_{(w_1, \dots, w_r) \in W^r} x^{\ell(w_1) + \dots + \ell(w_r)} z^{\ell_1(w_1) + \dots + \ell_1(w_r)}. \quad (2)$$

- On the other hand, for  $b \geq 1$ ,  $\mathcal{E}(b, k)$  is the set of  $\gamma = (\gamma_1, \dots, \gamma_r) \in W^r$ ,  $r \geq 1$ , such that  $\ell(\gamma_1) + \dots + \ell(\gamma_r) = b$  and  $0 \leq \ell_1(\gamma_1) + \dots + \ell_1(\gamma_r) \leq k$ .
- Thus

$$\begin{aligned} \sum_{b \geq 0} J(b, k, 1) x^b &= 1 + \sum_{b \geq 1} |\mathcal{E}(b, k)| x^b \\ &= 1 + \sum_{r \geq 1} \sum_{\substack{(w_1, \dots, w_r) \in W^r \\ 0 \leq \ell_1(w_1) + \dots + \ell_1(w_r) \leq k}} x^{\ell(w_1) + \dots + \ell(w_r)} \\ &= \sum_{s=0}^k [z^s] \left( 1 + \sum_{r \geq 1} \sum_{(w_1, \dots, w_r) \in W^r} x^{\ell(w_1) + \dots + \ell(w_r)} z^{\ell_1(w_1) + \dots + \ell_1(w_r)} \right). \end{aligned}$$

- By (2), this is equal to

$$\sum_{s=0}^k [z^s] \left( \frac{1}{1 - \sum_{i=1}^k x^i \sum_{j=0}^i z^j} \right) = [z^k] \left( \frac{1}{1-z} \cdot \frac{1}{1 - \sum_{i=1}^k x^i \sum_{j=0}^i z^j} \right),$$

as desired.

## Another expression

### Definition

## Another expression

### Definition

- A **composition** of  $k$  is a sequence  $\alpha = (\alpha_1, \dots, \alpha_r)$  of positive integers summing to  $k$ .

## Another expression

### Definition

- A **composition** of  $k$  is a sequence  $\alpha = (\alpha_1, \dots, \alpha_r)$  of positive integers summing to  $k$ .
- The **length**  $\ell(\alpha)$  of  $\alpha$  is defined to be the number  $r$  of elements in  $\alpha$ .

## Another expression

### Definition

- A **composition** of  $k$  is a sequence  $\alpha = (\alpha_1, \dots, \alpha_r)$  of positive integers summing to  $k$ .
- The **length**  $\ell(\alpha)$  of  $\alpha$  is defined to be the number  $r$  of elements in  $\alpha$ .
- We denote by  $\text{Comp}(k)$  the set of compositions of  $k$ .

## Another expression

### Definition

- A **composition** of  $k$  is a sequence  $\alpha = (\alpha_1, \dots, \alpha_r)$  of positive integers summing to  $k$ .
- The **length**  $\ell(\alpha)$  of  $\alpha$  is defined to be the number  $r$  of elements in  $\alpha$ .
- We denote by  $\text{Comp}(k)$  the set of compositions of  $k$ .

### Theorem

For a positive integer  $k$ , we have

$$\sum_{b \geq 0} J(b, k, 1) x^b = \sum_{\alpha \in \text{Comp}(k)} \frac{(-1)^{\ell_2(\alpha)} x^{k - \ell(\alpha)}}{(1 - x - \cdots - x^k)^{1 + \ell(\alpha)}},$$

where  $\ell_2(\alpha)$  is the number of parts of  $\alpha$  at least 2.

## Another expression

### Definition

- A **composition** of  $k$  is a sequence  $\alpha = (\alpha_1, \dots, \alpha_r)$  of positive integers summing to  $k$ .
- The **length**  $\ell(\alpha)$  of  $\alpha$  is defined to be the number  $r$  of elements in  $\alpha$ .
- We denote by  $\text{Comp}(k)$  the set of compositions of  $k$ .

### Theorem

For a positive integer  $k$ , we have

$$\sum_{b \geq 0} J(b, k, 1) x^b = \sum_{\alpha \in \text{Comp}(k)} \frac{(-1)^{\ell_2(\alpha)} x^{k - \ell(\alpha)}}{(1 - x - \cdots - x^k)^{1 + \ell(\alpha)}},$$

where  $\ell_2(\alpha)$  is the number of parts of  $\alpha$  at least 2.

### Corollary

For a positive integer  $k$ , we have

$$\sum_{b \geq 0} J(b, k, 1) x^b = \sum_{r=1}^k \sum_{s=0}^r \frac{(-1)^{r-s} \binom{r}{s} \binom{k-r-1}{r-s-1} x^{k-r}}{(1 - x - \cdots - x^k)^{1+r}}.$$

## Example

If  $k = 1$ , then  $\text{Comp}(k) = \{(1)\}$ . Thus

$$\sum_{b \geq 0} J(b, 1, 1)x^b = \frac{1}{(1-x)^2}.$$

## Example

If  $k = 1$ , then  $\text{Comp}(k) = \{(1)\}$ . Thus

$$\sum_{b \geq 0} J(b, 1, 1)x^b = \frac{1}{(1-x)^2}.$$

## Example

If  $k = 2$ , then  $\text{Comp}(k) = \{(2), (1, 1)\}$ . Thus

$$\sum_{b \geq 0} J(b, 2, 1)x^b = \frac{-x}{(1-x-x^2)^2} + \frac{1}{(1-x-x^2)^3} = \frac{1-x+x^2+x^3}{(1-x-x^2)^3}.$$

This proves Butler et al. 's conjecture.

## Example

If  $k = 1$ , then  $\text{Comp}(k) = \{(1)\}$ . Thus

$$\sum_{b \geq 0} J(b, 1, 1)x^b = \frac{1}{(1-x)^2}.$$

## Example

If  $k = 2$ , then  $\text{Comp}(k) = \{(2), (1, 1)\}$ . Thus

$$\sum_{b \geq 0} J(b, 2, 1)x^b = \frac{-x}{(1-x-x^2)^2} + \frac{1}{(1-x-x^2)^3} = \frac{1-x+x^2+x^3}{(1-x-x^2)^3}.$$

This proves Butler et al. 's conjecture.

## Example

If  $k = 3$ , then  $\text{Comp}(k) = \{(3), (2, 1), (1, 2), (1, 1, 1)\}$ . Thus

$$\begin{aligned} \sum_{b \geq 0} J(b, 3, 1)x^b &= \frac{-x^2}{(1-x-x^2-x^3)^2} + \frac{-2x}{(1-x-x^2-x^3)^3} + \frac{1}{(1-x-x^2-x^3)^4} \\ &= \frac{1-2x+x^2+4x^3+3x^4-3x^6-2x^7-x^8}{(1-x-x^2-x^3)^4}. \end{aligned}$$

## Juggling card sequences with fixed capacity

- We found an expression for

$$\sum_{b \geq 0} J(b, k, 1)x^b.$$

# Juggling card sequences with fixed capacity

- We found an expression for


$$\sum_{b \geq 0} J(b, k, 1)x^b.$$

- **Goal:** Find an expression for

$$\sum_{b \geq 0} J(b, k, \ell)x^b.$$


## Embeddings

- We can extend the notion of an embedding of a card to a sequence of cards.



## Embeddings


- We can extend the notion of an embedding of a card to a sequence of cards.



## Observation

## Embeddings

- We can extend the notion of an embedding of a card to a sequence of cards.




### Observation

- In the  $i$ th card, if there are  $r$   $i$ 's then  $r$  balls are thrown at beat  $i$ .

## Embeddings

- We can extend the notion of an embedding of a card to a sequence of cards.



### Observation

- In the  $i$ th card, if there are  $r$   $i$ 's then  $r$  balls are thrown at beat  $i$ .
- The  $r$  thrown balls have an associated word of integers in  $0, 1, \dots, i - 1$ .

## Definition

For a nonnegative integer  $n$ , the **homogeneous symmetric function**  $h_n(x_1, \dots, x_m)$  is defined by

$$h_n(x_1, \dots, x_m) = \sum_{i_1 + \dots + i_m = n} x_1^{i_1} \cdots x_m^{i_m},$$

where the sum is over all  $m$ -tuples  $(i_1, \dots, i_m)$  of nonnegative integers such that  $i_1 + \dots + i_m = n$ .

## Definition

For a nonnegative integer  $n$ , the **homogeneous symmetric function**  $h_n(x_1, \dots, x_m)$  is defined by

$$h_n(x_1, \dots, x_m) = \sum_{i_1 + \dots + i_m = n} x_1^{i_1} \cdots x_m^{i_m},$$

where the sum is over all  $m$ -tuples  $(i_1, \dots, i_m)$  of nonnegative integers such that  $i_1 + \dots + i_m = n$ .

## Definition

For indeterminates  $z_1, \dots, z_k$ , we denote by  $D_{z_1, \dots, z_k}$  the linear operator on the space of formal power series in  $z_k$  defined by

$$D_{z_1, \dots, z_k} z_k^n = h_n(1, z_1, \dots, z_{k-1}) z_k^n.$$

## Definition

For a nonnegative integer  $n$ , the **homogeneous symmetric function**  $h_n(x_1, \dots, x_m)$  is defined by

$$h_n(x_1, \dots, x_m) = \sum_{i_1 + \dots + i_m = n} x_1^{i_1} \cdots x_m^{i_m},$$

where the sum is over all  $m$ -tuples  $(i_1, \dots, i_m)$  of nonnegative integers such that  $i_1 + \dots + i_m = n$ .

## Definition

For indeterminates  $z_1, \dots, z_k$ , we denote by  $D_{z_1, \dots, z_k}$  the linear operator on the space of formal power series in  $z_k$  defined by

$$D_{z_1, \dots, z_k} z_k^n = h_n(1, z_1, \dots, z_{k-1}) z_k^n.$$

## Remark

Note that this can be seen as a generalization of the  **$q$ -derivative operator**  $(\frac{d}{dz})_q$ , which is the linear operator on the space of formal power series in  $z$  defined by

$$\left(\frac{d}{dz}\right)_q z^n = (1 + q + \cdots + q^{n-1}) z^{n-1}.$$

Hence  $D_{q,z}$  is equal to the operator  $(\frac{d}{dz})_{qz}$ , which multiplies  $z$  and then takes the  $q$ -derivative.

# Main result

## Theorem

For fixed positive integers  $k$  and  $\ell$ , we have

$$\begin{aligned} & \sum_{b \geq 0} J(b, k, \ell) x^b \\ &= [z_1^k \cdots z_\ell^k] \left( \frac{1}{1 - z_1} \cdots \frac{1}{1 - z_\ell} D_{z_1, z_2} D_{z_1, z_2, z_3} \cdots D_{z_1, \dots, z_\ell} \frac{1}{2 - h_k(1, x, xz_1, \dots, xz_\ell)} \right). \end{aligned}$$

## Rationality of the generating function

- By the quotient rule in calculus, one can easily deduce that the derivative of a rational function is also a rational function.

## Rationality of the generating function

- By the quotient rule in calculus, one can easily deduce that the derivative of a rational function is also a rational function.
- For the  $q$ -derivative, it is well known that

$$\left( \frac{d}{dx} \right)_q f(x) = \frac{f(x) - f(qx)}{x - qx}.$$

## Rationality of the generating function

- By the quotient rule in calculus, one can easily deduce that the derivative of a rational function is also a rational function.
- For the  $q$ -derivative, it is well known that

$$\left( \frac{d}{dx} \right)_q f(x) = \frac{f(x) - f(qx)}{x - qx}.$$

- This implies that the  $q$ -derivative of a rational function is also a rational function.

## Rationality of the generating function

- By the quotient rule in calculus, one can easily deduce that the derivative of a rational function is also a rational function.
- For the  $q$ -derivative, it is well known that

$$\left( \frac{d}{dx} \right)_q f(x) = \frac{f(x) - f(qx)}{x - qx}.$$

- This implies that the  $q$ -derivative of a rational function is also a rational function.
- Since

$$D_{z_1, z_2} z_2^n = (1 + z_1 + \cdots + z_1^n) z_2^n = \frac{(1 - z_1^{n+1}) z_2^n}{1 - z_1},$$

we have

$$D_{z_1, z_2} f(z_2) = \frac{f(z_2) - z_1 f(z_1 z_2)}{1 - z_1}. \quad (3)$$

## Rationality of the generating function

- By the quotient rule in calculus, one can easily deduce that the derivative of a rational function is also a rational function.
- For the  $q$ -derivative, it is well known that

$$\left( \frac{d}{dx} \right)_q f(x) = \frac{f(x) - f(qx)}{x - qx}.$$

- This implies that the  $q$ -derivative of a rational function is also a rational function.
- Since

$$D_{z_1, z_2} z_2^n = (1 + z_1 + \cdots + z_1^n) z_2^n = \frac{(1 - z_1^{n+1}) z_2^n}{1 - z_1},$$

we have

$$D_{z_1, z_2} f(z_2) = \frac{f(z_2) - z_1 f(z_1 z_2)}{1 - z_1}. \quad (3)$$

- Hence, if  $f(z_2)$  is a rational function in  $z_2$ , then so is  $D_{z_1, z_2} f(z_2)$ .

## Lemma

For integers  $n \geq 0$  and  $m \geq 2$ , we have

$$h_n(1, z_1, \dots, z_m) = \frac{z_{m-1} h_n(1, z_1, \dots, z_{m-1}) - z_m h_n(1, z_1, \dots, z_{m-2}, z_m)}{z_{m-1} - z_m}.$$

## Lemma

For integers  $n \geq 0$  and  $m \geq 2$ , we have

$$h_n(1, z_1, \dots, z_m) = \frac{z_{m-1} h_n(1, z_1, \dots, z_{m-1}) - z_m h_n(1, z_1, \dots, z_{m-2}, z_m)}{z_{m-1} - z_m}.$$

## Lemma

For an integer  $m \geq 2$ , we have

$$D_{z_1, \dots, z_{m+1}} = \frac{z_{m-1} D_{z_1, \dots, z_{m-1}, z_{m+1}} - z_m D_{z_1, \dots, z_{m-2}, z_m, z_{m+1}}}{z_{m-1} - z_m}.$$

## Lemma

For integers  $n \geq 0$  and  $m \geq 2$ , we have

$$h_n(1, z_1, \dots, z_m) = \frac{z_{m-1} h_n(1, z_1, \dots, z_{m-1}) - z_m h_n(1, z_1, \dots, z_{m-2}, z_m)}{z_{m-1} - z_m}.$$

## Lemma

For an integer  $m \geq 2$ , we have

$$D_{z_1, \dots, z_{m+1}} = \frac{z_{m-1} D_{z_1, \dots, z_{m-1}, z_{m+1}} - z_m D_{z_1, \dots, z_{m-2}, z_m, z_{m+1}}}{z_{m-1} - z_m}.$$

## Proposition

Suppose that  $\ell$  and  $m$  are integers with  $2 \leq m \leq \ell$  and let  $z_1, \dots, z_\ell$  be indeterminates.

If  $f(z_1, \dots, z_\ell)$  is a formal power series in  $z_1, \dots, z_\ell$  that is a rational function in  $z_1, \dots, z_\ell$ , then  $D_{z_1, \dots, z_m} f(z_1, \dots, z_\ell)$  is a rational function in  $z_1, \dots, z_\ell$ .

## Corollary

For fixed positive integers  $k$  and  $\ell$ , the generating function

$$\sum_{b \geq 0} J(b, k, \ell) x^b$$

is a rational function in  $x$ .

## Corollary

For fixed positive integers  $k$  and  $\ell$ , the generating function

$$\sum_{b \geq 0} J(b, k, \ell) x^b$$

is a rational function in  $x$ .

Proof.



## Corollary

For fixed positive integers  $k$  and  $\ell$ , the generating function

$$\sum_{b \geq 0} J(b, k, \ell) x^b$$

is a rational function in  $x$ .

## Proof.

- We proved that

$$\sum_{b \geq 0} J(b, k, \ell) x^b = [z_1^k \cdots z_\ell^k] f(z_1, \dots, z_\ell, x) \quad (4)$$

for a formal power series  $f(z_1, \dots, z_\ell, x)$  in  $z_1, \dots, z_\ell, x$  that is a rational function in these indeterminates.



## Corollary

For fixed positive integers  $k$  and  $\ell$ , the generating function

$$\sum_{b \geq 0} J(b, k, \ell) x^b$$

is a rational function in  $x$ .

## Proof.

- We proved that

$$\sum_{b \geq 0} J(b, k, \ell) x^b = [z_1^k \cdots z_\ell^k] f(z_1, \dots, z_\ell, x) \quad (4)$$

for a formal power series  $f(z_1, \dots, z_\ell, x)$  in  $z_1, \dots, z_\ell, x$  that is a rational function in these indeterminates.

- If  $g(z)$  is a formal power series in  $z$  that is a rational function in  $z$  and some other indeterminates, say  $u_1, \dots, u_r$ , then by the quotient rule,  $[z^k] g(z) = k! g^{(k)}(0)$  is a rational function in  $u_1, \dots, u_r$ .



## Corollary

For fixed positive integers  $k$  and  $\ell$ , the generating function

$$\sum_{b \geq 0} J(b, k, \ell) x^b$$

is a rational function in  $x$ .

## Proof.

- We proved that

$$\sum_{b \geq 0} J(b, k, \ell) x^b = [z_1^k \cdots z_\ell^k] f(z_1, \dots, z_\ell, x) \quad (4)$$

for a formal power series  $f(z_1, \dots, z_\ell, x)$  in  $z_1, \dots, z_\ell, x$  that is a rational function in these indeterminates.

- If  $g(z)$  is a formal power series in  $z$  that is a rational function in  $z$  and some other indeterminates, say  $u_1, \dots, u_r$ , then by the quotient rule,  $[z^k] g(z) = k! g^{(k)}(0)$  is a rational function in  $u_1, \dots, u_r$ .
- Therefore the right-hand side of (4) is a rational function of  $x$  as desired.



## Conclusion

- We found an expression for the generating function

$$\sum_{b \geq 0} J(b, k, \ell) x^b$$

and showed that it is a rational function in  $x$ .

## Conclusion

- We found an expression for the generating function

$$\sum_{b \geq 0} J(b, k, \ell) x^b$$

and showed that it is a rational function in  $x$ .

- It can be shown more easily that these are also rational functions:

$$\sum_{k \geq 0} J(b, k, \ell) x^k,$$

$$\sum_{\ell \geq 0} J(b, k, \ell) x^\ell.$$

## Conclusion

- We found an expression for the generating function

$$\sum_{b \geq 0} J(b, k, \ell) x^b$$

and showed that it is a rational function in  $x$ .

- It can be shown more easily that these are also rational functions:

$$\sum_{k \geq 0} J(b, k, \ell) x^k,$$

$$\sum_{\ell \geq 0} J(b, k, \ell) x^\ell.$$

### Problem

Determine whether the following is a rational function in  $x, y$ , and  $z$ :

$$\sum_{b \geq 0} \sum_{k \geq 0} \sum_{\ell \geq 0} J(b, k, \ell) x^b y^k z^\ell.$$

## Real multiplex juggling patterns

- Note that  $J(b, k, \ell)$  can be used to compute the number of ways to juggle  $b$  balls with capacity  $k$  for beats  $1, \dots, \ell$  without any restrictions on the initial and final states of the balls.

## Real multiplex juggling patterns

- Note that  $J(b, k, \ell)$  can be used to compute the number of ways to juggle  $b$  balls with capacity  $k$  for beats  $1, \dots, \ell$  without any restrictions on the initial and final states of the balls.
- In order to enumerate periodic multiplex juggling patterns we need to consider the number  $J_0(b, k, \ell)$  of  $\ell$ -cards  $(C_1, \dots, C_\ell) \in \mathcal{J}(b, k, \ell)$  such that the left side of  $C_1$  is equal to the right side of  $C_\ell$ .

## Real multiplex juggling patterns

- Note that  $J(b, k, \ell)$  can be used to compute the number of ways to juggle  $b$  balls with capacity  $k$  for beats  $1, \dots, \ell$  without any restrictions on the initial and final states of the balls.
- In order to enumerate periodic multiplex juggling patterns we need to consider the number  $J_0(b, k, \ell)$  of  $\ell$ -cards  $(C_1, \dots, C_\ell) \in \mathcal{J}(b, k, \ell)$  such that the left side of  $C_1$  is equal to the right side of  $C_\ell$ .

### Problem

For fixed  $k$  and  $\ell$ , find a formula for

$$\sum_{b \geq 0} J_0(b, k, \ell) x^b.$$

# Real multiplex juggling patterns

- Note that  $J(b, k, \ell)$  can be used to compute the number of ways to juggle  $b$  balls with capacity  $k$  for beats  $1, \dots, \ell$  without any restrictions on the initial and final states of the balls.
- In order to enumerate periodic multiplex juggling patterns we need to consider the number  $J_0(b, k, \ell)$  of  $\ell$ -cards  $(C_1, \dots, C_\ell) \in \mathcal{J}(b, k, \ell)$  such that the left side of  $C_1$  is equal to the right side of  $C_\ell$ .

## Problem

*For fixed  $k$  and  $\ell$ , find a formula for*

$$\sum_{b \geq 0} J_0(b, k, \ell) x^b.$$

## Problem

*Determine whether the following is a rational function in  $x, y, z$ :*

$$\sum_{b \geq 0} \sum_{k \geq 0} \sum_{\ell \geq 0} J_0(b, k, \ell) x^b y^k z^\ell.$$

Thank you!