

Lecture hall graphs and the Askey scheme

2023. 10. 28

Jang Soo Kim (Sungkyunkwan Univ.)

Joint work with

Sylvie Corteel (Univ. Paris Cité)

Bhargavi Jonnaula (Oxford Univ.)

Jon Keating (Oxford Univ.)

Outline

- ① Basics on orthogonal polynomials.
- ② Lecture hall partitions and little q -Jacoby poly.
- ③ Lecture hall graph.
- ④ Askey-Wilson polynomials
- ⑤ Main results

Orthogonal polynomials

Def) $\{P_n(x)\}_{n \geq 0}$ is a sequence of **orthogonal polynomials** (OPS) with respect to a linear functional \mathcal{L} if

$$\textcircled{1} \quad \deg P_n(x) = n$$

$$\textcircled{2} \quad \mathcal{L}(P_n(x)P_m(x)) = \begin{cases} 0 & \text{if } n \neq m \\ \text{nonzero} & \text{if } n = m. \end{cases}$$

Def) The ***n*th moment** μ_n of OP is $\mathcal{L}(x^n)$.

Favard's thm $\{P_n(x)\}_{n \geq 0}$ is a monic OPS iff

$$P_{n+1}(x) = (x - b_n)P_n(x) - \lambda_n P_{n-1}(x). \quad (\lambda_n \neq 0).$$

$$-b_0 -\lambda_2 \propto \propto -b_5 \propto -\lambda_8$$


Cor $P_n(x) = \sum$ 

Thm (Viennot, 1983)

Let $\{P_n(x)\}$ be a monic OPS satisfying

$$P_{n+1}(x) = (x - b_n) P_n(x) - \lambda_n P_{n-1}(x).$$

$$\Rightarrow \mu_n = \sum_{\pi \in \text{Motz}_n} \text{wt}(\pi).$$



$$\text{wt}(\pi) = b_0 b_1 b_2 \lambda_1^2 \lambda_2 \lambda_3$$

Motzkin path

ex). Hermite polynomial

$$\mu_n = (2n-1)!! = 1 \cdot 3 \cdot \dots \cdot (2n-1) = \# \text{ perfect matchings}$$

ex) Charlier polynomial

$$\mu_n = \# \text{ set partitions on } [n] = \{1, 2, \dots, n\}.$$

ex) Laguerre polynomial

$$\mu_n = n! = \# \text{ permutations.}$$

Another way to define moments.

$P_n(x)$: OPS w.r.t. \mathcal{L} . ($\mathcal{L}(1)=1$, $P_0(x)=1$)

Let $x^n = \sum_{k=0}^n \mu_{n,k} P_k(x)$.

Then $\mathcal{L}(x^n) = \sum_{k=0}^n \mu_{n,k} \underbrace{\mathcal{L}(P_k(x))}_{=\mathcal{L}(P_k(x)P_0(x))} = \mu_{n,0}$.

Def). $P_n(x)$: OPS

let $x^n = \sum_{k=0}^n \mu_{n,k} P_k(x)$, $P_n(x) = \sum_{k=0}^n v_{n,k} x^k$.

$\mu_{n,k}$ is the *mixed moment*.

$v_{n,k}$ is the *dual mixed moment*.

(These are the entries of change of bases.)

Little q -Jacobi polynomials $p_n(x; a, b; q)$

$$p_n(x; a, b; q) = \frac{(ab; q)_n}{(-1)^n q^{-\binom{n}{2}} (abq^{n+1}; q)_n} {}_2\phi_1 \left(\begin{matrix} q^{-n}, abq^{n+1} \\ ab \end{matrix}; q, qx \right),$$

$${}_2\phi_1 \left(\begin{matrix} A, B \\ C \end{matrix}; q, z \right) = \sum_{i \geq 0} \frac{(A; q)_i (B; q)_i}{(q; q)_i (C; q)_i} z^i,$$

$$(A; q)_i = (1-A)(1-Aq)\cdots(1-Aq^{i-1})$$

Corteel, K. 2020


There are combinatorial models for $\nu_{n,k}$, $\mu_{n,k}$.
using lecture hall partitions.

Def) $\lambda \vdash n \iff \lambda$ is a partition of n
 $\iff \lambda = (\lambda_1, \dots, \lambda_k), \quad \lambda_1 + \dots + \lambda_k = n$
 $\lambda_1 \geq \dots \geq \lambda_k > 0$

Each $\lambda_i > 0$ is called a part.

ex) $\lambda = (4, 3, 1, 1)$ is a partition of $9 = 4+3+1+1$.

- Young diagram of $\lambda = (4, 3, 1)$ is






Lecture hall partitions

$\lambda = (\lambda_1, \dots, \lambda_n)$ is a **lecture hall partition** if

$$\frac{\lambda_1}{n} \geq \frac{\lambda_2}{n-1} \geq \dots \geq \frac{\lambda_n}{1} \geq 0$$

ex) $(5, 3, 1)$ is LHP

$(5, 3, 2)$ is NOT LHP



Thm (Bousquet-Mélou, Eriksson, 1997)


$$\sum_{\lambda \in L_n} q^{|\lambda|} = \prod_{i=1}^n \frac{1}{1-q^{2i-1}}$$

Anti-lecture hall compositions

$\alpha = (\alpha_1, \dots, \alpha_n)$ is an anti-lecture hall composition
(or a planetarium composition)

if $\frac{\alpha_1}{1} \geq \frac{\alpha_2}{2} \geq \dots \geq \frac{\alpha_n}{n} \geq 0$.

ex)



$(4, 5, 3, 3, 2, 1)$

Thm (Corteel, Savage, 2003)

$$\sum_{\alpha \in A_n} q^{|\alpha|} = \prod_{i=1}^n \frac{1+q^i}{1-q^{i+1}}$$

Thm (Corteel, K., 2020)


Letting $a = -uv$, $b = -u/v$ for little q -Jacobi,

$$M_{n,k} = \sum_{\frac{\alpha_1}{k+1} \geq \dots \geq \frac{\alpha_{n-k}}{n} \geq 0} u^{\lfloor L\alpha \rfloor} v^{o(L\alpha)} q^{|\alpha|}$$

$$v_{n,k} = \sum_{\frac{\lambda_1}{n} > \dots > \frac{\lambda_{n-k}}{k+1} > 0} u^{\lfloor L\lambda \rfloor} v^{o(L\lambda)} q^{|\lambda|}$$

$\Rightarrow M_{n,k}$ = generating function for antilecture hall compositions

$v_{n,k} = \dots \quad \text{"} \quad \text{"} \quad \text{"} \quad \text{"} \quad \text{lecture hall partitions.}$



Askey-Wilson polynomials

$$p_n(x; a, b, c, d | q) = \frac{(ab, ac, ad; q)_n}{2^n a^n (abcdq^{n-1}; q)_n} {}_4\phi_3 \left(\begin{matrix} q^{-n}, abcdq^{n-1}, ae^{i\theta}, ae^{-i\theta} \\ ab, ac, ad \end{matrix}; q, q \right),$$

where $x = \cos \theta = (e^{i\theta} + e^{-i\theta})/2$.

$$P_{n+1} = (x - b_n) P_n - \lambda_n P_{n-1}$$

$$b_n = \frac{1}{2} (a + a^l - A_n - C_n), \quad \lambda_n = \frac{1}{4} A_{n-1} C_n$$


$$A_n = \frac{(1 - abq^n)(1 - acq^n)(1 - adq^n)(1 - abcdq^{n-1})}{a(1 - abcdq^{2n-1})(1 - abcdq^{2n})},$$

$$C_n = \frac{a(1 - q^n)(1 - bcq^{n-1})(1 - bdq^{n-1})(1 - cdq^{n-1})}{(1 - abcdq^{2n-2})(1 - abcdq^{2n-1})}.$$

Fact : $p_n(x; a, b, c, d | q)$ is symmetric in a, b, c, d .

Thm (Corteel, Williams, 2011)

The Askey-Wilson moment μ_n is a partition function for ASEP.




state diagram of ASEP (asymmetric simple exclusion process)

Thm (Corteel, Stanley, Stanton, Williams. 2012)

$$2^n(abcd; q)_n \mu_n(a, b, c, d; q)$$

$$\begin{aligned} &= i^{-n} \sum_{T \in \mathcal{T}(n)} (-1)^{b(T)} (1-q)^{A(T)+B(T)+C(T)+D(T)-n} q^{E(T)} \\ &\times (ac)^{C(T)} (bd)^{D(T)} ((1+ai)(1+ci))^{n-A(T)-C(T)} ((1-bi)(1-di))^{n-B(T)-D(T)}. \end{aligned}$$



After rescaling & reparametrization
 μ_n is a polynomial in $\alpha, \beta, \gamma, \delta$.

$$\begin{aligned} \alpha &= \frac{1-q}{(1+ai)(1+ci)}, & \beta &= \frac{1-q}{(1-bi)(1-di)}, \\ \gamma &= \frac{ac(1-q)}{(1+ai)(1+ci)}, & \delta &= \frac{bd(1-q)}{(1-bi)(1-di)}. \end{aligned}$$

staircase tableaux


Motivation

- ① Find a lecture hall partition model for Askey-Wilson.
- ② Find a combinatorial proof of the symmetry in a, b, c, d .


Q. Why can't we generalize Corneel-Kim result
on little q -Jacobi to Askey-Wilson?

→ We need a new approach to "guess" a combinatorial model.

Lecture hall graph (Introduced by Corteel - Kim 2020
Further studied by Corteel - Keating - Nicoletti 2021)



(k, ∞)



$(n, 0)$

$$\frac{4}{2} \geq \frac{5}{3} \geq \frac{3}{4} \geq \frac{2}{5} \quad (k=1, n=5).$$

$$\frac{\lambda_{k+1}}{k+1} \geq \frac{\lambda_{k+2}}{k+2} \geq \dots \geq \frac{\lambda_n}{n} \quad \leftrightarrow \text{ path from } (k, \infty) \text{ to } (n, 0)$$

Thm (Corneel, K., 2020)

little q -Jacobi mixed moment

$$M_{n,K} = \sum_{p:(K,\infty) \rightarrow (n,0)} \text{wt}(p)$$

(K,∞)

		⋮	⋮	⋮
4	$a^2b^2q^4$	$a^2b^2q^8$	$a^2b^2q^{12}$	
3	$-a^2bq^3$	$-a^2bq^7$	$-a^2bq^{11}$...
	$-a^2bq^6$	$-a^2bq^{10}$		
2	abq^5	abq^8		...
	abq^2	abq^4	abq^6	
1	$-aq^3$	$-aq^5$...
	$-aq^2$	$-aq^4$		
0	q	q^2		...
	1	1	1	

(n,0)

Thm (CJKKK, 2023)

big q -Jacobi

$$P_n(x; a, b, c | q) = \frac{(aq, cq; q)_n}{(abq^{n+1}; q)_n} {}_3\phi_2 \left(\begin{matrix} q^{-n}, abq^{n+1}\alpha \\ aq, cq \end{matrix} ; q, q \right)$$

$$\mu_{n,k} = \sum_{p: (K, \infty) \rightarrow (n, 0)} \text{wt}(p)$$

Cor Let $\tilde{\mu}_{n,k} = (-1)^{n-k} \mu_{n,k}(-a, -b, -c)$.

$\Rightarrow (\tilde{\mu}_{n,k})_{n,k=0}^\infty$ is totally positive.

Equivalently multivariate big q -Jacobi
is Schur positive.

		⋮	⋮
4	$abcq^3$	$abcq^5$	$abcq^7$
	$-abq^4$	$-abq^6$	$-abq^8$
	$-abq^2$	$-abq^3$	$-abq^4$
	$-acq^4$	$-acq^6$	$-acq^8$
3	$-acq^2$	$-acq^3$	$-acq^4$
	$-abq^4$	$-abq^5$	$-abq^6$
	$-abq^3$	$-abq^4$	$-abq^5$
	$-abq$	$-ab$	$-a$
2	acq^2	aq^2	aq^3
	acq	aq	aq^2
	abq	ab	a
	ab	a	1
1	cq^2	cq^3	cq^4
	cq	cq	cq^2
	c	c	c
	0	1	2

Thm (CJKK, 2023)

Askey-Wilson mixed moment

$$\mu_{n,k} = \sum_{p:(k,\infty) \rightarrow (n,0)} \text{wt}(p)$$


$$\nu_{n,k} = \sum_{\substack{p:(k,0) \rightarrow (n,\infty) \\ \text{no consecutive east step}}} \text{wt}(p)$$

$$f(x) = x + \bar{x}$$

9	$-bcd/2$	$-bcdq/2$	$-bcdq^2/2$	$-bcdq^3/2$
	$a^2bcdq^3/2$	$a^2bcdq^6/2$	$a^2bcdq^9/2$	
8	$a^2bcd/2$	$a^2bcdq/2$	$a^2bcdq^4/2$	$a^2bcdq^7/2$
	$-abcq^2/2$	$-abcq^4/2$	$-abcq^6/2$	
7	$-abc/2$	$-abcq/2$	$-abcq^2/2$	$-abcq^3/2$
	$-abdq^2/2$	$-abdq^4/2$	$-abdq^6/2$	
6	$-abd/2$	$-abdq/2$	$-abdq^2/2$	$-abdq^3/2$
	$bq/2$	$bq^2/2$	$bq^3/2$	
5	$b/2$	$b/2$	$b/2$	
	$-acdq^2/2$	$-acdq^4/2$	$-acdq^6/2$	
4	$-acd/2$	$-acdq/2$	$-acdq^2/2$	$-acdq^3/2$
	$cq/2$	$cq^2/2$	$cq^3/2$	
3	$c/2$	$c/2$	$c/2$	
	$dq/2$	$dq^2/2$	$dq^3/2$	
2	$d/2$	$d/2$	$d/2$	
	$-1/(2a)$	$-1/(2a)$	$-1/(2a)$	
1	$-1/(2a)$	$-1/(2aq)$	$-1/(2aq)$	
	$(aq+)/2$	$(aq^2+)/2$	$(aq^3+)/2$	
0	$(a+)/2$	$(a+)/2$	$(a+)/2$	


Thm (CJKK, 2023).

There is a lecture hall graph model for $M_{n,k}$ and $V_{n,k}$
for every orthogonal polynomial in Askey scheme



Idea of proof

Two bootstrapping methods



Thm (CJKK)

continuous q -Hermite

$$\mu_{n,k} = \sum_{p:(k,3) \rightarrow (n,0)} w(p)$$

3		$-1/2$	$-1/2$	$-1/2$	$-q/2$	$-q^2/2$	\dots
2	$-1/2$	$-q/2$	$-q^2/2$	$-q^3/2$	$-1/2$	$-1/(2q)$	\dots
1	$-1/2$	$-1/(2q)$	$-1/(2q^2)$	$-1/(2q^3)$	$(q^2+)/2$	$(q^3+)/2$	\dots
0	$(1+)/2$	$(1+)/2$	$(1+)/2$	$(1+)/2$	$(q+)/2$	$(q^2+)/2$	\dots
	0	1	2	3	4		

Thm (CJKK)

mixed moment of AW
with respect to q-Hermite

$$H_n(x) = \sum_k \tilde{\mu}_{n,k} P_k^{AW}(x)$$

$$\Rightarrow \frac{2^{n-k}}{i} \tilde{\mu}_{n,k}(a_i, b_i, c_i, d_i)$$

$$= \sum_{p: (k, \infty) \rightarrow (n, 0)} \text{wt}(p)$$

Cor Askey-Wilson is symmetric
in a, b, c, d .

9	$abcd^2$	$abcd^2q^2$	$abcd^2q^4$	$abcd^2q^6$
		$bcdq^2$	$bcdq^4$	$bcdq^6$
			$bcdq^3$	$bcdq^5$
8	bcd	$bcdq$	$bcdq^2$	$bcdq^3$
			a^2bcdq^3	a^2bcdq^6
			a^2bcdq^5	a^2bcdq^8
7	a^2bcd	a^2bcdq^2	a^2bcdq^4	a^2bcdq^6
			$acdq^2$	$acdq^4$
6	acd	$acdq$	$acdq^2$	$acdq^3$
			$abdq^2$	$abdq^4$
5	abd	$abdq$	$abdq^2$	$abdq^3$
			dq	dq^3
4	d	d	d	d
			$abcq^2$	$abcq^4$
3	abc	$abcq$	$abcq^3$	$abcq^5$
			$abcq^2$	$abcq^4$
2	c	c	c	c
			bq	bq^3
1	b	b	b	b
			aq	aq^3
0	a	a	a	a

Advantages of lecture hall graph models.

- ① a model for $M_{n,k}$ gives a model for $V_{n,k}$.
- ② multivariate moments can be obtained by
Lindström–Gessel–Viennot Lemma
- ③ Total positivity
- ④ Applicable to all OPS in Askey scheme.

Thank You!

Moments of multivariate
orthogonal polynomials

Symmetric polynomials

Def) $F(x_1, \dots, x_n)$ is symmetric if

$$F(x_{\sigma(1)}, \dots, x_{\sigma(n)}) = F(x_1, \dots, x_n) \text{ for all } \sigma \in S_n.$$

When we have a basis $\{P_m(x)\}_{m \geq 0}$ of univariate poly,
we can construct a basis $\{P_\lambda(x)\}_\lambda$ of sym. poly.

by $P_\lambda(x) = \frac{\det(P_{\lambda_i+n-i}(x_j))}{\det(P_{n-i}(x_j))}.$ $X = (x_1, \dots, x_n)$

Def) The Schur function $S_\lambda(x_1, \dots, x_n)$ is defined
in this way with basis $\{x^m\}_{m \geq 0}:$

$$S_\lambda(x) = \frac{\det(x_j^{\lambda_i+n-i})}{\det(x_j^{n-i})}.$$

Multivariate little q -Jacobi poly.

$$\text{Def)} \quad P_{\lambda}(x; a, b; q) = \frac{\det(P_{\lambda_i+n-i}(x_i; a, b; q))}{\det(P_{n-i}(x_i; a, b; q))}.$$


Since both P_λ and S_λ are bases for the space of symmetric polynomials, we can expand

Multivariate little q -Jacobi polynomials
& Lecture hall tableaux

Basic definitions

- $\lambda = (\lambda_1, \dots, \lambda_L)$ is a **partition** of n
if $\lambda_1 \geq \dots \geq \lambda_L > 0$ and $\lambda_1 + \dots + \lambda_L = n$

- Young diagram of $\lambda = (4, 3, 1)$ is



- Standard Young tableau

1	3	4	7
2	5	8	
6			


\leq

1	1	2	2
2	3	3	
4			

- Semistandard Young tableau

$$\mu = \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \subseteq \lambda = \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array}$$

skew shape $\lambda/\mu =$



- Schur function

$$s_{\lambda/\mu}(x) = \sum_{T \in \text{SSYT}(\lambda/\mu)} x^T,$$

$$T = \begin{array}{|c|c|c|} \hline 1 & 3 \\ \hline 1 & 2 \\ \hline 2 \\ \hline \end{array}$$

$$x^T = x_1^2 x_2^2 x_3^1$$

Lecture hall tableaux

T : a filling of λ with nonnegative integers

T is a **lecture hall tableau** of type $(n, \geq, >)$ if

$$\frac{T(i,j)}{n+c(i,j)} \geq \frac{T(i,j+1)}{n+c(i,j+1)}$$

∨

$$\frac{T(i+1,j)}{n+c(i+1,j)}$$

$$c(i,j) = j - i \quad (\text{content})$$

ex)

8	9	10	9	4	3	
4	5	6	4	3	1	
2	2	1	0			
1	0	0				

$T =$

LHT of type $(5, \geq, >)$

$8/5$	$9/6$	$10/7$	$9/8$	$4/9$	$3/10$	
$4/4$	$5/5$	$6/6$	$4/7$	$3/8$	$1/9$	
$2/3$	$2/4$	$1/5$	$0/6$			
$1/2$	$0/3$	$0/4$				

∨

If $n \rightarrow \infty$, LHT is \sim SSYT.

(reverse)

The multivariate little q -Jacobi polynomials (with $t=q$)

$$P_\lambda(x) = P_\lambda(x_1, \dots, x_n; a, b; q)$$

$$S_\lambda(x) = \sum_{\mu \leq \lambda} M_{\lambda, \mu} P_\mu(x),$$

$$P_\lambda(x) = \sum_{\mu \leq \lambda} N_{\lambda, \mu} S_\mu(x)$$

Thm (Corneel, K., 2018)

$M_{\lambda, \mu}$ = g.f. for LHT of shape λ/μ of type $(n, \geq, >)$

$N_{\lambda, \mu} = \dots \quad \dots \quad \dots \quad \dots \quad (n, <, \leq)$

The "moment" $M_{\lambda, \phi}$ has a product formula.

The "dual moment" $N_{\lambda, \phi}$ " .