

# STOR 320.1

## Tidy Data

# Introduction


- Read Chapter 9
- Functions From tidyr Package

```
>library(tidyr)
```

- pivot\_longer()
- pivot\_wider()
- separate()
- unite()
- complete()

# Tidy Data Definition

- For Tidy Data:
  - Each Variable Must Have Its Own Column
  - Each Observation Must Have Its Own Row
  - Each Value Must Have Its Own Cell



# Problem

- Most Data is Not Tidy
- Reason: Data Collectors Often Don't Know How Data Should Be Recorded Since They Don't Analyze the Data
- Common Problems
  - A Variable Spread Across Multiple Columns
  - A Observation is Spread Across Multiple Rows

“Tidy datasets are all alike, but every messy dataset is messy in its own way.” — Hadley Wickham

# Untidy Data Example 1

- Multiple Columns for One Variable

```
untidy1=tribble(  
  ~subject, ~sex, ~control, ~cond1, ~cond2,  
  1, "M", 7.9, 12.3, 10.7,  
  2, "F", 6.3, 10.6, 11.1,  
  3, "F", 9.5, 13.1, 13.8,  
  4, "M", 11.5, 13.4, 12.9  
)  
untidy1
```

```
## # A tibble: 4 x 5  
##   subject sex    control cond1 cond2  
##   <dbl> <chr>    <dbl> <dbl> <dbl>  
## 1       1 M        7.9   12.3  10.7  
## 2       2 F        6.3   10.6  11.1  
## 3       3 F        9.5   13.1  13.8  
## 4       4 M       11.5   13.4  12.9
```

# Problem

```
## # A tibble: 4 x 5
##   subject sex    control cond1 cond2
##   <dbl> <chr>    <dbl> <dbl> <dbl>
## 1       1 M        7.9  12.3  10.7
## 2       2 F        6.3  10.6  11.1
## 3       3 F        9.5  13.1  13.8
## 4       4 M       11.5  13.4  12.9
```

- Multiple Treatment Data
- Variables “Control”, “Cond1”, and “Cond2” are Measuring the Same Thing Under Different Treatments
- The Name of the Variable Whose Values Form the Column Names Can Be Called “Treatment”
- The Name of the Variable Whose Values are Spread Over the Cells Can Be Called “Outcome”

# Longer

```
```{r}
tidy1a=untidy1 %>%
  pivot_longer(control:cond2, names_to = "Treatment",
values_to = "Outcome")
tidy1a
```
```

| subject<br><dbl> | sex<br><chr> | Treatment<br><chr> | Outcome<br><dbl> |
|------------------|--------------|--------------------|------------------|
| 1                | M            | control            | 7.9              |
| 1                | M            | cond1              | 12.3             |
| 1                | M            | cond2              | 10.7             |
| 2                | F            | control            | 6.3              |
| 2                | F            | cond1              | 10.6             |
| 2                | F            | cond2              | 11.1             |
| 3                | F            | control            | 9.5              |
| 3                | F            | cond1              | 13.1             |
| 3                | F            | cond2              | 13.8             |
| 4                | M            | control            | 11.5             |
| 4                | M            | cond1              | 13.4             |
| 4                | M            | cond2              | 12.9             |

# Longer by index

```
```{r}
tidy1b=untidy1 %>%
  pivot_longer(3:5, names_to="Treatment", values_to="Outcome")
tidy1b
```
```

| subject<br><dbl> | sex<br><chr> | Treatment<br><chr> | Outcome<br><dbl> |
|------------------|--------------|--------------------|------------------|
| 1                | M            | control            | 7.9              |
| 1                | M            | cond1              | 12.3             |
| 1                | M            | cond2              | 10.7             |
| 2                | F            | control            | 6.3              |
| 2                | F            | cond1              | 10.6             |
| 2                | F            | cond2              | 11.1             |
| 3                | F            | control            | 9.5              |
| 3                | F            | cond1              | 13.1             |
| 3                | F            | cond2              | 13.8             |
| 4                | M            | control            | 11.5             |
| 4                | M            | cond1              | 13.4             |
| 4                | M            | cond2              | 12.9             |



# Process


```
## # A tibble: 4 x 5
##   subject sex    control cond1 cond2
##   <dbl> <chr>    <dbl> <dbl> <dbl>
## 1      1 M      7.9    12.3    10.7
## 2      2 F      6.3    10.6    11.1
## 3      3 F      9.5    13.1    13.8
## 4      4 M     11.5    13.4    12.9
```

| subject<br><dbl> | sex<br><chr> | Treatment<br><chr> | Outcome<br><dbl> |
|------------------|--------------|--------------------|------------------|
| 1                | M            | control            | 7.9              |
| 1                | M            | cond1              | 12.3             |
| 1                | M            | cond2              | 10.7             |
| 2                | F            | control            | 6.3              |
| 2                | F            | cond1              | 10.6             |
| 2                | F            | cond2              | 11.1             |
| 3                | F            | control            | 9.5              |
| 3                | F            | cond1              | 13.1             |
| 3                | F            | cond2              | 13.8             |
| 4                | M            | control            | 11.5             |
| 4                | M            | cond1              | 13.4             |
| 4                | M            | cond2              | 12.9             |

- pivot\_longer vs. gather()

# Longer

```
```{r}  
ggplot(tidy1b)+  
  geom_point(aes(x=Treatment,y=Outcome,color=sex),size=4) +  
  theme_minimal()  
```
```



# Untidy Data Example 2

```
untidy2=tribble(  
  ~subject, ~sex, ~`0.3`, ~`0.6`, ~`0.8`,  
  1, "M", 7.9, 12.3, 10.7,  
  2, "F", 6.3, 10.6, 11.1,  
  3, "F", 9.5, 13.1, 13.8,  
  4, "M", 11.5, 13.4, 12.9  
)  
untidy2
```

```
## # A tibble: 4 x 5  
##   subject sex    `0.3` `0.6` `0.8`  
##   <dbl> <chr> <dbl> <dbl> <dbl>  
## 1     1 M      7.9   12.3  10.7  
## 2     2 F      6.3   10.6  11.1  
## 3     3 F      9.5   13.1  13.8  
## 4     4 M     11.5   13.4  12.9
```

# Problem

```
## # A tibble: 4 x 5
##   subject sex    `0.3` `0.6` `0.8`
##   <dbl> <chr> <dbl> <dbl> <dbl>
## 1       1 M      7.9  12.3  10.7
## 2       2 F      6.3  10.6  11.1
## 3       3 F      9.5  13.1  13.8
## 4       4 M     11.5  13.4  12.9
```

- Repeated Measures Data
- Variables “0.3”, “0.6”, and “0.8” are Measuring the Same Thing Under Different Drug Strengths
- The Name of the Variable Whose Values Form the Column Names Can Be Called “Dosage”
- The Name of the Variable Whose Values are Spread Over the Cells Can Be Called “Outcome”

# Longer

```
```{r}
tidy2a=untidy2 %>%
  pivot_longer(`0.3`:`0.8`,names_to="Dosage",values_to="Outcome")
glimpse(tidy2a)
```
```

Rows: 12  
Columns: 4

|            |       |   |
|------------|-------|---|
| \$ subject | <dbl> | 1, 1, 1, 2, 2, 2, 3, 3, 3, 4, 4, 4                          |
| \$ sex     | <chr> | "M", "M", "M", "F", "F", "F", "F", "F", "F", "M", "M", ...  |
| \$ Dosage  | <chr> | "0.3", "0.6", "0.8", "0.3", "0.6", "0.8", "0.3", "0.6", ... |
| \$ Outcome | <dbl> | 7.9, 12.3, 10.7, 6.3, 10.6, 11.1, 9.5, 13.1, 13.8, 11.5...  |

```
```{r}
tidy2b=untidy2 %>%
  pivot_longer(3:5,names_to="Dosage_ch",values_to="Outcome") %>%
  mutate(Dosage=as.numeric(Dosage_ch)) %>%
  select(-Dosage_ch)
glimpse(tidy2b)
```
```


Rows: 12  
Columns: 4

|            |       |  |
|------------|-------|--|
| \$ subject | <dbl> | 1, 1, 1, 2, 2, 2, 3, 3, 3, 4, 4, 4                         |
| \$ sex     | <chr> | "M", "M", "M", "F", "F", "F", "F", "F", "F", "M", "M", ... |
| \$ Outcome | <dbl> | 7.9, 12.3, 10.7, 6.3, 10.6, 11.1, 9.5, 13.1, 13.8, 11.5... |
| \$ Dosage  | <dbl> | 0.3, 0.6, 0.8, 0.3, 0.6, 0.8, 0.3, 0.6, 0.8, 0.3, 0.6, ... |

```
tidy2b=untidy2 %>%
  gather(`0.3`:`0.8`,key="Dosage",value="Outcome",convert=T)
glimpse(tidy2b)
```

# Longer

```
```{r}  
ggplot(tidy2b) +  
  geom_line(aes(x=Dosage,y=Outcome,color=as.character(subject))) +  
  theme_minimal()  
```
```



# Untidy Data Example 3

- Multiple rows

```
untidy3=tribble(  
  ~Pack, ~Type, ~Measure, ~Value,  
  1, "Regular", "Count", 15,  
  1, "Regular", "Percent Blue", 0.2,  
  2, "Peanut", "Count", 12,  
  2, "Peanut", "Percent Blue", 0.3,  
)  
untidy3
```

```
## # A tibble: 4 x 4  
##   Pack Type Measure Value  
##   <dbl> <chr> <chr> <dbl>  
## 1     1 Regular Count    15  
## 2     1 Regular Percent Blue 0.2  
## 3     2 Peanut Count    12  
## 4     2 Peanut Percent Blue 0.3
```

# Problem

- Less Common
- Column “Measures” Contains Variable Names
- Column “Value” Contains the Output of the Different Variables
- Notice Values are of Different Units (Count vs Percentage)
- Wider Does the Opposite of Longer

```
## # A tibble: 4 x 4
##   Pack Type Measure Value
##   <dbl> <chr> <chr> <dbl>
## 1     1 Regular Count    15
## 2     1 Regular Percent Blue 0.2
## 3     2 Peanut Count    12
## 4     2 Peanut Percent Blue 0.3
```




# Wider

```
```{r}
tidy3=untidy3 %>%
  pivot_wider(names_from=Measure,values_from=Value)
tidy3
```
```

```
## # A tibble: 2 x 4
##   Pack Type      Count `Percent Blue`
##   <dbl> <chr>    <dbl>         <dbl>
## 1     1 1 Regular      15           0.2
## 2     2 2 Peanut      12           0.3
```

# Process

```
## # A tibble: 4 x 4
##   Pack Type Measure      Value
##   <dbl> <chr>  <chr>    <dbl>
## 1     1  1 Regular Count      15
## 2     1  1 Regular Percent Blue  0.2
## 3     2  2 Peanut  Count      12
## 4     2  2 Peanut  Percent Blue  0.3
```



```
## # A tibble: 2 x 4
##   Pack Type Count `Percent Blue`
##   <dbl> <chr>  <dbl>    <dbl>
## 1     1  1 Regular    15      0.2
## 2     2  2 Peanut    12      0.3
```

# Wider

```
tidy3 %>%  
  mutate(nBlue=Count*`Percent Blue`) %>%  
  select(-Count,-`Percent Blue`)
```

```
## # A tibble: 2 x 3  
##   Pack Type    nBlue  
##   <dbl> <chr>    <dbl>  
## 1     1 Regular     3  
## 2     2 Peanut    3.6
```

# Untidy Data Example 4

```
untidy4=tribble(  
  ~Pack, ~Type, ~PropBlue, ~Date,  
  1, "Regular", "3/15", "9-28-2018",  
  2, "Regular", "2/15", "9-30-2018",  
  3, "Peanut", "4/12", "9-28-2018",  
  4, "Peanut", "5/13", "9-30-2018",  
)  
untidy4
```

```
## # A tibble: 4 x 4  
##   Pack Type   PropBlue Date  
##   <dbl> <chr>   <chr>   <chr>  
## 1     1 Regular 3/15     9-28-2018  
## 2     2 Regular 2/15     9-30-2018  
## 3     3 Peanut 4/12     9-28-2018  
## 4     4 Peanut 5/13     9-30-2018
```

# Problem

```
## # A tibble: 4 x 4
##   Pack Type   PropBlue Date
##   <dbl> <chr>   <chr>   <chr>
## 1     1     1 Regular 3/15     9-28-2018
## 2     2     2 Regular 2/15     9-30-2018
## 3     3     3 Peanut 4/12     9-28-2018
## 4     4     4 Peanut 5/13     9-30-2018
```

- Very Uncommon
- The Variable “PropBlue” Contains Two Numeric Variables
- The Variable “Date” Contains Three Numeric Variables
- We Must Separate Both of These Variables Into Multiple Columns

# Separating


```
```{r}
tidy4a=untidy4 %>%
  separate(PropBlue, into=c("nBlue","Total"),sep="/") %>%
  separate(Date, into=c("M","D","Y"),sep="-")
glimpse(tidy4a)
```
```

```
Rows: 4
Columns: 7
$ Pack  <dbl> 1, 2, 3, 4
$ Type  <chr> "Regular", "Regular", "Peanut", "Peanut"
$ nBlue <chr> "3", "2", "4", "5"
$ Total <chr> "15", "15", "12", "13"
$ M     <chr> "9", "9", "9", "9"
$ D     <chr> "28", "30", "28", "30"
$ Y     <chr> "2018", "2018", "2018", "2018"
```

```
```{r}
tidy4b=untidy4 %>%
  separate(PropBlue, into=c("nBlue","Total"),convert=T) %>%
  separate(Date, into=c("M","D","Y"),convert=T)
glimpse(tidy4b)
```
```

```
Rows: 4
Columns: 7
$ Pack  <dbl> 1, 2, 3, 4
$ Type  <chr> "Regular", "Regular", "Peanut", "Peanut"
$ nBlue <int> 3, 2, 4, 5
$ Total <int> 15, 15, 12, 13
$ M     <int> 9, 9, 9, 9
$ D     <int> 28, 30, 28, 30
$ Y     <int> 2018, 2018, 2018, 2018
```

# Separating



# Untidy Data Example 5

```
untidy5=tribble(  
  ~Pack, ~Type, ~Day, ~Month,  
  1, "Regular", 1, 8,  
  2, "Regular", 2, 8,  
  3, "Regular", 3, 9,  
  4, "Regular", 4, 9,  
)  
untidy5
```

```
## # A tibble: 4 x 4  
##   Pack Type      Day Month  
##   <dbl> <chr>   <dbl> <dbl>  
## 1     1 Regular     1     8  
## 2     2 Regular     2     8  
## 3     3 Regular     3     9  
## 4     4 Regular     4     9
```



# Uniting

- Absolutely Silly
- Uniting Does the Opposite of Separating

```
tidy5=untidy5 %>%  
  unite(swap, Day, Month, sep=": ")  
tidy5
```

```
## # A tibble: 4 x 3  
##   Pack Type      swap  
##   <dbl> <chr>    <chr>  
## 1     1 Regular 1: (8  
## 2     2 Regular 2: (8  
## 3     3 Regular 3: (9  
## 4     4 Regular 4: (9
```

# Missing Values

- Two Ways
  - Explicitly: Defined to Be Missing Using NA
  - Implicitly: Absent From Data
- There is not a Uniform Way to Handle Either of These Problems
- Rule: Either Convert All Explicitly Missing to Implicitly Missing or Convert All Implicitly Missing to Explicitly Missing

# Example

```
## # A tibble: 14 x 3
##   year quarter wage
##   <dbl>   <dbl> <dbl>
## 1     1       1     1  10.5
## 2     2       1     2  10.5
## 3     3       1     3  10.5
## 4     4       1     4   11
## 5     5       2     2   11
## 6     6       2     3  11.2
## 7     7       3     1  11.2
## 8     8       3     2  11.2
## 9     9       3     3   12
## 10    10       3     4  NA
## 11    11       4     1   12
## 12    12       4     2  NA
## 13    13       4     3  13.0
## 14    14       4     4  13.0
```

# Missing Values

- Notice:

```
```{r}
missing %>%
  pivot_wider(names_from = year, values_from = wage)
```
```

| quarter | 1     | 2     | 3     | 4     |
|---------|-------|-------|-------|-------|
| <dbl>   | <dbl> | <dbl> | <dbl> | <dbl> |
| 1       | 10.5  | NA    | 11.23 | 12.00 |
| 2       | 10.5  | 11.00 | 11.23 | NA    |
| 3       | 10.5  | 11.23 | 12.00 | 13.04 |
| 4       | 11.0  | NA    | NA    | 13.04 |

```
```{r}
missing %>%
  pivot_wider(names_from=quarter, values_from=wage)
```
```

| year  | 1     | 2     | 3     | 4     |
|-------|-------|-------|-------|-------|
| <dbl> | <dbl> | <dbl> | <dbl> | <dbl> |
| 1     | 10.50 | 10.50 | 10.50 | 11.00 |
| 2     | NA    | 11.00 | 11.23 | NA    |
| 3     | 11.23 | 11.23 | 12.00 | NA    |
| 4     | 12.00 | NA    | 13.04 | 13.04 |

# Missing Values

```
```{r}
missing %>%
  pivot_wider(names_from=quarter, values_from=wage) %>%
  pivot_longer(2:5, names_to='quarter', values_to='wage')
```
```

- Implicit to Explicit

| year<br><dbl> | quarter<br><chr> | wage<br><dbl> |
|---------------|------------------|---------------|
| 1             | 1                | 10.50         |
| 1             | 2                | 10.50         |
| 1             | 3                | 10.50         |
| 1             | 4                | 11.00         |
| 2             | 1                | NA            |
| 2             | 2                | 11.00         |
| 2             | 3                | 11.23         |
| 2             | 4                | NA            |
| 3             | 1                | 11.23         |
| 3             | 2                | 11.23         |
| 3             | 3                | 12.00         |
| 3             | 4                | NA            |
| 4             | 1                | 12.00         |
| 4             | 2                | NA            |
| 4             | 3                | 13.04         |
| 4             | 4                | 13.04         |

# Missing Values

- Explicit to Implicit

```
```{r}
missing %>%
  pivot_wider(names_from=quarter, values_from=wage) %>%
  pivot_longer(2:5, names_to='quarter', values_to='wage', values_drop_na = T)
```
```

| <b>year</b> | <b>quarter</b> | <b>wage</b> |
|-------------|----------------|-------------|
| <dbl>       | <chr>          | <dbl>       |
| 1           | 1              | 10.50       |
| 3           | 1              | 11.23       |
| 4           | 1              | 12.00       |
| 1           | 2              | 10.50       |
| 2           | 2              | 11.00       |
| 3           | 2              | 11.23       |
| 1           | 3              | 10.50       |
| 2           | 3              | 11.23       |
| 3           | 3              | 12.00       |
| 4           | 3              | 13.04       |
| 1           | 4              | 11.00       |
| 4           | 4              | 13.04       |

# Missing Values

- Complete Function

```
missing %>%  
  complete(year, quarter)
```

```
## # A tibble: 16 x 3  
##   year quarter wage  
##   <dbl>   <dbl> <dbl>  
## 1     1     1     1  10.5  
## 2     1     2    10.5  
## 3     1     3    10.5  
## 4     1     4    11  
## 5     2     1    NA  
## 6     2     2    11  
## 7     2     3   11.2  
## 8     2     4    NA  
## 9     3     1   11.2  
## 10    3     2   11.2  
## 11    3     3    12  
## 12    3     4    NA  
## 13    4     1    12  
## 14    4     2    NA  
## 15    4     3   13.0  
## 16    4     4   13.0
```

# Tidy data Training

**Now, let us**

# **PRACTICE**

**Download the Rmd for Tutorial 5 to Your Computer from the Course Website and open the file in RStudio**