



Practicum - Consultation: 14.03.2024, 09-11

Please send your Group solutions to corneliafranziska.richter@vetpharm.uzh.ch on time.

What	File name	When
Group solutions	"02worksheet-Group-Name.zip"	18.03.2024 at 7 am.

Please upload all files to OLAT on time.

What	File name	When
Individual project	"02worksheet-Your-Name.zip"	18.03.2024 at 7 am.
Group contribution	"03contribution-Group-Name.zip"	19.03.2024 at 22 pm.

Individual tasks

Your 02worksheet-Your-Name.zip file contains reproducible code necessary to generate your results for both 2A and 2B parts and your report together with the resulting pdf-file.

Exercise 1 (Individual project (Part 2A))

Provide classical and Bayesian analyses for Secukinumab and placebo separately.

- Compute 95% confidence intervals (95%CI) for the true probability of response in the Secukinumab and in the placebo groups for the data provided in Table 1. Interpret 95%CI in each group separately.
- Plot both $\text{Beta}(0.5, 1)$ and the $\text{Beta}(11, 32)$ priors and summarize them with prior mean, median, and the equi-tailed 95% interval.
- Consider the Secukinumab group in Table 1: plot the $\text{Beta}(0.5, 1)$ prior, plot the posterior distribution of the response rate, and summarize the posterior distribution with posterior mean, median, and the equi-tailed 95% credible interval (95%CrI). Interpret the 95%CrI in the Secukinumab group.
- Consider the placebo group in Table 1: plot the $\text{Beta}(11, 32)$ prior, plot the posterior distribution of the response rate, and summarize the posterior distribution with posterior mean, median, and the equi-tailed 95%CrI. Interpret the 95%CrI in the placebo group.

Report your results.

Exercise 2 (Individual project (Part 2B))

Baeten et al. (2013) write in their supplementary material: "predictive distribution was



Table 1: ASAS20 responders at week 6: data and results provided explicitly and implicitly in Table 2 of Baeten et al. (2013). Priors: Beta(0.5, 1) in the Secukinumab group and Beta(11, 32) in the placebo group.

Group	n	Responders x (%)	Posterior response rate
Secukinumab	23	14 (60.9%)	59.2%
Placebo	6	1 (16.7%)	24.5%

approximated by a Beta density with matching mean and standard deviation". In this exercise you implement a function that computes α and β shape parameters of a Beta distribution with known mean and variance.

- Assume that sample mean and sample variance values of a Beta distribution are known. Derive analytically formulas that compute values of α and β shape parameters of the Beta distribution given its sample mean and sample variance (moments matching).
- Implement these formulas in a function in R. Input: sample mean and sample variance of a Beta distribution. Output: α and β shape parameters of the Beta distribution.
- Apply your function to mean= 0.255814 and variance= 0.004326663. What are the resulting values of α and β parameters of the Beta distribution?

Report your results.

Group tasks

Your o2worksheet-Group-Name.zip (one per group) file contains reproducible code necessary to generate your results and your report together with the resulting pdf-file, which can contain scans of your handwritten solutions. List the names of students who contributed to the solution of group tasks.

Exercise 3 (Conjugate Bayes: analytical derivation)

Assume that y_1, \dots, y_n are realizations (observations) generated by *iid* random variables which follow a $N(m, \kappa^{-1})$ distribution. Moreover, assume that the prior of m follows a $N(\mu, \lambda^{-1})$ distribution, where κ , μ and λ are fixed (known) constants. Derive analytically (with all constants!) the posterior distribution of m given that y_1, \dots, y_n have been observed.



Exercise 4 (Conjugate Bayesian analysis in practice)

Apply analytical formulas derived in Exercise 3 above to the vector of Height (cm) measurements 166, 168, 168, 177, 160, 170, 172, 159, 175, 164, 175, 167, 164 of 13 Swiss females. Assume that y_1, \dots, y_n are observations generated by $N(m, \kappa^{-1})$ distribution with $\kappa = 1/900$. Moreover, assume a $N(\mu, \lambda^{-1})$ prior for m with $\mu = 161$ and $\lambda = 1/70$.

- Apply descriptive graphs and summaries to the Height outcome. Compute a classical 95%CI for the true mean and interpret it.
- Plot the prior distribution of m , compute its expectation, standard deviation, median, and equi-tailed 95% interval. Estimate $P[m > 200]$ from the prior distribution.
- On the plot from task (b), plot the posterior distribution of $m|y_1, \dots, y_n$. Moreover, compute the posterior mean, standard deviation, and median. In addition, compute the equi-tailed 95%CrI and interpret it.
- Estimate $P[m > 200|y_1, \dots, y_n]$ under the posterior distribution.
- Discuss how much your opinion about moments of m and about $P[m > 200]$ have changed between the prior and the posterior.

Exercise 5 (Bayesian learning)

For an interim analysis after 12 subjects, 3 adverse events were observed. In the final study after seeing 64 subjects, 14 adverse events were seen.

Compare the results obtained for the following two Beta priors at each stage:

- $\alpha = \beta = 0.5$;
- $\alpha = 8, \beta = 24$.

How much evidence do you have for the probability of adverse events > 0.4 before seeing any data, at the stage of an intermediate study and for the final study?

Plot priors and posteriors at each stage.

What are the mean, median, and the equi-tailed 95%CrI of the distribution at each stage?

Group contributions to the "Big picture"

Exercise 6 (Group contributions for the lecture on 21.03.2024)



Please prepare a group contribution, which your group will present (ca. 5 min) during the next lecture.

- (3.1) Review asymptotical results in the classical statistics.
- (3.2) History of the Monte Carlo method.
- (3.3) Write a short program in R to estimate the area of the unit circle $x^2 + y^2 = 1$ by dropping randomly points (independent Monte Carlo sampling) on it.
- (3.4) An overview of random number generators used for independent Monte Carlo sampling in R and JAGS.

Make sure that the file `o3contribution-Group-Name.zip` (one per group) contains the file and the R code you want to present.

References

Baeten, D., X. Baraliakos, J. Braun, J. Sieper, P. Emery, D. van der Heijde, I. McInnes, J. van Laar, R. Landewé, P. Wordsworth, J. Wollenhaupt, H. Kellner, J. Paramarta, J. Wei, A. Brachat, S. Bek, D. Laurent, Y. Li, Y. Wang, A. Bertolino, S. Gsteiger, A. Wright, and W. Hueber (2013). Anti-interleukin-17A monoclonal antibody secukinumab in treatment of ankylosing spondylitis: a randomised, double-blind, placebo-controlled trial. *The Lancet* 382, 1705–1713.