

JAN VANHOVE

QUANTITATIVE METHODOLOGY

AN INTRODUCTION

AUTUMN 2026

UNIVERSITY OF FRIBOURG

Quantitative methodology: An introduction © 2020–2026 by Jan Vanhove is licensed under Creative Commons Attribution-NonCommercial 4.0 International.

To view a copy of this license, visit <https://creativecommons.org/licenses/by-nc/4.0/>.

jan.vanhove@unifr.ch

<https://janhove.github.io>

Preface

The principal goal of this class is to impart the fundamentals of quantitative research. Our focus will be on **drawing causal conclusions** from data—when are causal conclusions licensed, what are typical pitfalls when drawing causal conclusions, and how can you design and optimise studies that avoid these pitfalls? Throughout, it is imperative that you not only understand the recommendations given in this booklet but also the **logic** behind them. There are two reasons for this.

First, you need to know when the recommendations apply and when they don't. You might get away with just memorising the recommendations and their scope for now. But in a couple of months, you're bound to apply them where they don't make any sense. Understand the logic behind the recommendations, and you'll be better able to weigh your options.

Second, many researchers—even seasoned ones—operate on rules of thumb, so *they* inevitably end up applying recommendations they've picked up somewhere in situations where they don't apply. You need to be able to make an informed judgement about the research carried out by others and cogently argue for this judgement. A simple *I took this class that taught me that you shouldn't control for colliders* won't do. (You'll learn about colliders and why you shouldn't control for them soon enough.)

In this class, we'll cover the contents of this lecture script. In addition, we'll spend two sessions on questionnaire design. Furthermore, in the homework assignments, you'll learn how to visualise data using state-of-the-art software. These graphing assignments are available from <https://janhove.github.io/graphs>.

I occasionally refer to some blog entries I wrote. These links are clickable in the PDF version of this booklet, but in case you're reading this from paper, all blog entries can be found at <https://janhove.github.io>.

Jan Vanhove

jan.vanhove@unifr.ch

<https://janhove.github.io>

Contents

	<i>Preface</i>	3
1	<i>Association and causality</i>	7
1.1	<i>Definitions</i>	8
1.2	<i>Visualising causality: directed acyclic graphs (DAGs)</i>	9
1.3	<i>*Further reading</i>	18
2	<i>Constructing a control group</i>	19
2.1	<i>A made-up example</i>	19
2.2	<i>Critical questions</i>	20
2.3	<i>Increasing internal validity through randomisation</i>	22
3	<i>Alternative explanations</i>	31
3.1	<i>The roles of variables in research</i>	31
3.2	<i>Explanation 1: Expectancy effects</i>	32
3.3	<i>Explanation 2: Failed manipulation</i>	33
3.4	<i>Explanation 3: Chance</i>	35
4	<i>Inferential statistics 101</i>	37
4.1	<i>An example: Does alcohol intake affect fluency in a second language?</i>	37
4.2	<i>Chance as an alternative explanation</i>	38
4.3	<i>The null hypothesis</i>	39
4.4	<i>Testing the null hypothesis by exhaustive rerandomisation</i>	40
4.5	<i>On 'rejecting' null hypotheses</i>	43
4.6	<i>Monte Carlo rerandomisation p-values</i>	46
4.7	<i>Analytical shortcuts</i>	46
4.8	<i>Statistical power</i>	47
4.9	<i>*Examples in R</i>	48
4.10	<i>*Further reading</i>	48
5	<i>Increasing precision</i>	51

5.1	<i>Precision</i>	51
5.2	<i>Factors affecting precision</i>	52
5.3	<i>Matching and blocking</i>	53
5.4	<i>Leveraging control variables</i>	57
5.5	<i>*Examples in R</i>	61
6	<i>Pedagogical interventions</i>	63
6.1	<i>Sample mortality</i>	63
6.2	<i>Clustering</i>	64
6.3	<i>*Examples in R</i>	67
7	<i>Within-subjects experiments</i>	69
7.1	<i>Advantages and drawbacks</i>	69
7.2	<i>Minimising order effects</i>	70
7.3	<i>*Analysing data from within-subjects designs with two conditions</i>	73
7.4	<i>*Further reading</i>	75
7.5	<i>*Examples in R</i>	75
8	<i>Observational studies</i>	77
8.1	<i>Correlation coefficients</i>	78
8.2	<i>Statistical control using hierarchical regression</i>	80
8.3	<i>Caveats</i>	83
9	<i>Constructs and indicators</i>	85
9.1	<i>Systematic and random measurement error</i>	86
9.2	<i>Consequences of measurement error</i>	86
10	<i>Questionable research practices</i>	89
10.1	<i>A paradox</i>	89
10.2	<i>Hidden flexibility</i>	90
10.3	<i>*Further reading</i>	92
A	<i>Descriptive statistics</i>	93
B	<i>Reading difficult results sections</i>	95
C	<i>Reporting research transparently</i>	97
	<i>References</i>	99

1

Association and causality

Exercise 1.1 (Receptive multilingualism in Scandinavia). When talking their respective native languages, Danes understand Swedes better than the other way around. Furthermore, Danes like Swedish better than Swedes do Danish (see, e.g., [Delsing & Lundin Åkesson, 2005](#)).

1. Can we conclude that Danes understand Swedes better than the other way around *because* they like Swedish better than Swedes do Danish? If not, what are some plausible alternative explanations for the finding above?
2. Which additional finding would strengthen the claim that Danes understand Swedes better than the other way around because they like the language better? Several answers are possible.
3. Which additional finding would call this claim into question? Several answers are possible.



The diamond symbol signals where an example, exercise, remark, definition and such like ends. The few proofs included in this script end with an empty square symbol ($\square$).

Exercise 1.2 (Content and language integrated learning). Pupils in *Content and Language Integrated Learning* (CLIL) programmes in Andalusia perform better on English proficiency tests than other Andalusian pupils ([Lorenzo et al., 2010](#)).

1. Can we conclude that taking CLIL classes improves Andalusian pupils' English proficiency? If not, what are some plausible alternative explanations for the finding above?
2. Which additional finding would strengthen the claim that CLIL improves pupils' English proficiency? Several answers are possible.
3. Which additional finding would call this claim into question? Several answers are possible.



In both examples, an **association** of some sort is found in the data, and a **causal explanation** for this association is put forward: Not only do Danes both understand and like Swedish better than Swedes do Danish (association), it's suggested that one reason why they understand the other language better is that they like it better (explanation). Similarly, not only do CLIL pupils in Andalusia outperform non-CLIL pupils (association), it's suggested that they

outperform them *because* of the CLIL programme (explanation).

Uncovering associations and drawing causal conclusions from them is a key goal in empirical research. But it's also fraught with difficulty: after a moment's thought, you'll often be able to come up with alternative explanations for the findings. To the extent that there exist more, and more plausible, alternative explanations, the causal explanation proffered becomes more tenuous: The causal claim may still be correct, but in the presence of competing explanations, it can't be *shown* to be correct—that is, there isn't much **evidence** for the claim. A key goal when designing an empirical study is to reduce the number and the plausibility of such alternative explanations.

1.1 Definitions

Definition 1.3 (Association and independence). Two factors (or variables) are **associated** if knowing the value of one factor can help you hazard a more educated guess about the value of the other factor. If two factors are not associated, we say that they are **independent**. ◇

I use the terms *factor* and *variable* interchangeably. Sometimes, factors are constant rather than variable in the context of a study, but let's save our pedantic inclinations for other things.

Exercise 1.4. Explain why, according to Definition 1.3, the following are examples of associations:

- a person's height in centimetres and their height in inches;
- the time of day and the temperature outside;
- a person's height and their weight;
- a person's shoesize and the size of their vocabulary in their native language;
- a person's age and their father's age;
- a person's nationality and the colour of their eyes. ◇

From Exercise 1.4, we may draw a few conclusions:

- Associations work in both directions: knowing the time of day allows you to venture a more educated guess about the temperature outside than not knowing it, but also vice versa.
- Associations needn't be linear (e.g., the relation between weight and height levels off after a certain weight).
- Associations needn't be monotonic, e.g., the relationship between the two variables can go up and then down again (as in the time of day/temperature example).
- Associations needn't be perfect (e.g., there's a lot of variation about the general trend for taller people to be heavier).
- Associations can be found between variables that aren't typically expressed numerically (e.g., eye colour and nationality).

***Remark 1.5.** The concepts of association and independence can

Comments and sections prefaced with an asterisk are optional. You don't need to study them for the exam.

be made more precise mathematically, and in full-fledged statistics course, you would need the more precise definitions. For the purposes of this course, however, a conceptual understanding along the lines of the more informal Definition 1.3 suffices. ◇

Next, we define what we mean if we say that a variable D causally influences a variable Y .

Definition 1.6 (Causality). “We say that there is a **causal relationship** between [two variables] D and Y in a population if and only if there is at least one unit in that population for which intervening in the world to change D will change Y ” (Keele et al., 2019, p. 3) ◇

Definition 1.6 relies on a counterfactual and interventionist notion of causality: we are asked to imagine what the world *would have* looked like (‘counterfactual’) if we had *changed* D (‘interventionist’).

In Exercise 1.2, it is fairly easy to imagine that we could have arranged for some children who were in fact enrolled in a CLIL programme to have been enrolled in a non-CLIL programme instead, or vice versa, allowing us to track the development of their English proficiency in this parallel world we’ve created.

In Exercise 1.1, by contrast, it is less straightforward to imagine how we could intervene to directly change Danes’ attitudes towards Swedish, though perhaps we could imagine some intervention that is likely to induce a change in attitude. If we can’t imagine a plausible mechanism by which we could hypothetically change D , the causal question may be ill-defined (see Holland, 1986).

Three further remarks are in order:

- Saying that D causally influences Y doesn’t imply that D alone causally influences Y . (You can get lung cancer from smoking, but also from exposure to radon, air pollution or just genetic bad luck.)
- Saying that D causally influences Y doesn’t mean that changing D will result in a change in Y for *all* members of the population. (Some non-smokers get lung cancer, and not all smokers get it.)
- Saying that D causally influences Y doesn’t imply that changing D will result in a change in Y in all situations. (Smoldering cigarette stubs cause forest fires, but only during droughts. By the same token, droughts cause forest fires, but these need a spark to get started.)

The question of *whether* CLIL causally influences English proficiency seems trivial: for someone somewhere, it *has* to. The more interesting questions concern the strength of this causal effect and whether it is generally positive or negative.

1.2 Visualising causality: directed acyclic graphs (DAGs)

1.2.1 Why?

Research would be pretty easy if you could safely conclude that a causal link existed between two variables any time you observed an association between them. Fortunately for teachers of methodology




Figure 1.1: Source: <https://xkcd.com/552>.

courses who'd be on the dole otherwise, this isn't the case. But simply parroting back *Correlation is not causation* isn't too helpful (Figure 1.1). To help us figure out how associations between two variables can arise in the absence of a causal link between them, we turn to **directed acyclic graphs** (DAGs).

Definition 1.7 (Graphs). **Graphs** are mathematical objects in which nodes (also called vertices) can be connected by edges. In **directed** graphs, these edges point from one node to another, i.e., they're arrows. If, in a directed graph, it is impossible to start from some node and end up in the same node by following edges, then that graph is also **acyclic**.

A **path** is a sequence of distinct nodes that are linked by edges; the direction of the edges does not matter. A **directed path** from a node X to a node Y is a path in which you start at X and end at Y by following edges in the direction they are pointing in (i.e., $X \rightarrow W \rightarrow \dots \rightarrow Z \rightarrow Y$).

If, in a directed graph, there exists an edge from a node X to a node Y (i.e., $X \rightarrow Y$), then we call X a **parent** of Y and Y a **child** of X . If there exists a directed path from a node X to a node Y , then we call X an **ancestor** of Y and Y a **descendant** of X . In particular, parents are ancestors, and children are descendants. $\diamond$

Graphs are studied in mathematics and computer science for sundry purposes; here, we will use DAGs as a tool for visually representing the causal links between the factors at play in a study.

As we'll see below, when DAGs are used to represent causal links, they are subject to a number of rules that may appear cumbersome at first. However, when DAGs are properly specified, they allow researchers to figure out which factors they *should* control for, which factors they *can but needn't* control for and which factors they *must not* control for. Moreover, DAGs are useful for learning how associations in empirical data can occur both in the presence and in the absence of causal links between the variables of interest.

1.2.2 Some examples

Before spelling out the rules for drawing DAGs, let's look at a couple of possible DAGs for the Andalusian CLIL study.

Figure 1.2 is the simplest of DAGs. It represents the assumption that there is a direct causal influence from the **treatment** variable (CLIL) on the **outcome** variable. These variables are represented by nodes. There exists a directed edge (i.e., an arrow) between them that shows the assumed direction of the causal link.

The pupils' English proficiency won't be affected by their taking CLIL classes or not *alone* but by a host of other unobserved factors as well. In Figure 1.3, the unobserved factors are conveniently bundled and represented as 'U'. The U is circled to make it clear that these factors were not observed or measured. While this convention




Figure 1.2: DAG representing a causal influence of CLIL on English proficiency (ENG).




Figure 1.3: DAG representing a causal influence of CLIL and of unobserved factors on English proficiency.

isn't universal, it's useful and we'll adopt it here.

Important: If we don't draw an arrow between U and $CLIL$, this means that we assume that there is *no* direct causal relationship between these two factors. But presumably, some unobserved factors will also account for why some pupils are enrolled in $CLIL$ classes and others aren't; see Figure 1.4. As we'll discuss later, these unobserved factors, some of which may affect both the 'treatment' ($CLIL$) and the 'outcome' (English proficiency), **confound** the causal link of interest.

Figure 1.5 also features the unobserved factors as possible confounders, but this time there is no edge between $CLIL$ and ENG . Such a DAG can be useful for playing devil's advocate: We assume that there is no causal link between $CLIL$ and English proficiency and subsequently use the DAG to deduce if it is possible that $CLIL$ and English proficiency are nonetheless associated. If so, the mere presence of an association between $CLIL$ and English proficiency does not imply that there is some causal link between them.




Figure 1.4: Unobserved factors as confounders (1).




Figure 1.5: Unobserved factors as confounders (2).

1.2.3 Rules for drawing DAGs

1. The direction of the arrows shows the direction of the assumed causality (hence *directed*).
2. Bidirectional arrows are forbidden, i.e., no $A \leftrightarrow B$.
3. You're not allowed to draw graphs where you can end up at the same place where you started by just following arrows (hence *acyclic*). For instance, you're not allowed to draw a DAG like Figure 1.6. Mutually influencing factors can be represented in a DAG, however, but you need to break down the temporal structure. Figure 3 shows how you can break down the temporal structure implicit in Figure 1.6 to produce a legal DAG.




Figure 1.6: Not a DAG: A, B and C are allowed to influence themselves, making the graph cyclic rather than acyclic.




Figure 1.7: Reciprocal influences can be represented legally in a DAG if you break down the temporal structure. A_1, A_2, A_3 and A_4 represent the same variable measured at four points in time. The value of this variable at a given point in time is determined in part by its value at the previous point in time (e.g., A_3 is influenced by A_2) as well as by the value of another variable at the previous point in time (e.g., C_2 influences A_3).

4. Unobserved factors can, and often should, be drawn. By convention, we draw a circle around them to make it clear that they are not directly observed.
5. "DAGs insistently redirect the analyst's attention to justifying what arrows do not exist. Present arrows represent the analyst's ignorance. Missing arrows, by contrast, represent definitive

claims of knowledge.” (Elwert, 2013, p. 248)

6. A factor that isn’t of interest and that only affects one factor already in the DAG and/or is affected by only one factor already in the DAG doesn’t have to be drawn for you to be able to derive correct conclusions from the DAG. For instance, the U in Figure 1.3 doesn’t have to be drawn (it only affects one factor that was already in the DAG). However, the U in Figure 1.4 *does* have to be drawn since it affects *two* factors that were already in the DAG. That said, it can be difficult to decide if a variable should be included in a DAG or not, and we shouldn’t let perfect be the enemy of good. The purpose of DAGs is less to perfectly capture the true causal structure than making explicit one’s imperfect assumptions.

DAGs don’t specify *how* one variable causally influences another. For instance, if we draw $X \rightarrow Y$, this could mean that Y increases if we increase X , but it could also mean that Y decreases if we increase X . It’s even possible that Y first increases and then decreases, or that the pattern is another one altogether.

1.2.4 Chains, forks, and inverted forks

A DAG that is drawn by following the rules specified above is always built up out of at most three types of building blocks: chains, forks, and inverted forks.

Chains A chain is a sequence of causal links; in a DAG, chains show up as directed paths. In Figure 1.8, $A \rightarrow B \rightarrow C \rightarrow D$ forms a causal chain. Causality doesn’t flow ‘upstream’ against the direction of the arrows, so there is no causal chain from D back to A .

Chains may transmit genuine causal influences, that is, altering the values of (say) A may bring about a change in some values in (say) D . In other words, A may causally affect D , albeit indirectly through B and C . Since the causality is directional, altering the values of D won’t bring about any changes in the values of A , B or C .

Moreover, **chains may induce associations between the variables involved**. Based on the DAG in Figure 1.8, we wouldn’t be surprised to find some association between the values of A , B , C and D . The DAG doesn’t tell us what this association will look like, but we’ll encounter some common forms of association in the weeks to come.

It is possible that changes in A are not reflected in changes in D downstream. For instance, the effect that A has on B may be quite small, and perhaps only large changes in B affect C . This is why I wrote that changes in A *may* (rather than *will*) bring about changes in D .

If, for whatever reason, you want to prevent a chain from trans-




Figure 1.8: A chain.

mitting associations between two variables, the path between these variables has to be **blocked** somewhere. This is achieved by **controlling** for one (or several) of the variables along the path. A conceptually easy (if often practically arduous) way is to ensure that only people, words, etc. with the same value on that variable are included in the study. For instance, if for some reason you need to control for eye colour, you could include only green-eyed people in your study.

Example 1.8. Consider the chain $A \rightarrow B \rightarrow C$, where A, B, C can take on the values 0 and 1. The causal influence of A on B is such that there is a 75% probability that $B = 1$ if $A = 1$ but only a 25% probability that $B = 1$ if $A = 0$. Similarly, there is a 75% probability that $C = 1$ if $B = 1$ but only a 25% probability that $C = 1$ if $B = 0$, independently of the value of A .

As you can verify, the probability that $C = 1$ if $A = 1$ is

$$0.75 \cdot 0.75 + 0.25 \cdot 0.75 = 0.75,$$

that is, 75%, whereas the probability that $C = 1$ if $A = 0$ is $0.25 \cdot 0.75 + 0.75 \cdot 0.25 = 0.375$ (37.5%). Hence, A and C are associated.

We can control for B by fixing its value at one value. If we fix B at 0, then the probability that $C = 1$ is 25% regardless of whether $A = 0$ or $A = 1$, that is, A and C are no longer associated if we only look at those cases where $B = 0$. Similarly, if we fix B at 1, the probability that $C = 1$ is 75% regardless of the value of A . ◇

Forks When a single factor causally affects two or more other factors, a fork is formed; see Figure 1.9. In this example, A causally influences both B and C .

Forks themselves don't transmit causal influences between the prongs, that is, altering the values of B won't change the values of C and vice versa: Causality doesn't travel upstream. If you want to represent a causal link between B and C , you have to add it to the DAG.

Importantly, **forks may induce associations between the factors at the prongs**: Based on the DAG in Figure 1.9, we wouldn't be surprised to find some association between the values of B and C . This is not because of a causal link between them but because A influences both of them. A is also referred to as a **confounding variable** or **confounder**.

Example 1.9. To better appreciate the fact that causal forks can give rise to associations between the variables at the prongs, consider the fictitious example in Table 1.9. Here, A causally influences both B and C , and both B and C are additionally influenced by separate factors (U_B and U_C). The causal factors A , U_B and U_C can each take on two values (0, 1), and the outcomes of B and C are determined by simple equations. We assume that the probability of observing




Figure 1.9: A fork.

the values in a particular row is the same for all rows, i.e., that it is $1/8$.

A	U_B	U_C	$B := A + U_B$	$C := A + U_C$
0	0	0	0	0
0	0	1	0	1
0	1	0	1	0
0	1	1	1	1
1	0	0	1	1
1	0	1	1	2
1	1	0	2	1
1	1	1	2	2

Taking a closer look at this table, we see that B and C are associated: The overall probability that B is at least equal to 1 is $\mathbb{P}(B \geq 1) = 6/8 = 75\%$. But if you already know that an observation's value for C is 2, then you can be absolutely confident that its B value is at least 1: $\mathbb{P}(B \geq 1|C = 2) = 2/2 = 100\%$. By Definition 1.3, B and C are associated. By the same token, if you know that its C value is 0, you'd be less confident about this guess:

- $\mathbb{P}(B \geq 1|C = 0) = 1/2$.
- $\mathbb{P}(B \geq 1|C = 1) = 3/4$.
- $\mathbb{P}(B \geq 1|C = 2) = 2/2$.

If you want to prevent a fork from transmitting an association between the variables at the prongs, you can control for the confounder or otherwise block the path on which the confounder lies. To appreciate this fact, again consider Table 1.9. We've already established that $\mathbb{P}(B \geq 1|C = 0) \neq \mathbb{P}(B \geq 1|C = 1)$. But once we 'control for' A by fixing it at a specific value (e.g., $A = 0$), we find that the probability of observing $B \geq 1$ doesn't depend on C any more.

- $\mathbb{P}(B \geq 1|C = 0, A = 0) = 1/2$.
- $\mathbb{P}(B \geq 1|C = 1, A = 0) = 1/2$.

Similarly, we could fix A at 1 and vary C and observe the same phenomenon:

- $\mathbb{P}(B \geq 2|C = 1, A = 1) = 1/2$.
- $\mathbb{P}(B \geq 2|C = 2, A = 1) = 1/2$.

◇

Exercise 1.10. While I don't have the numbers handy, I'm confident that there is some positive association between the daily number of drownings in the River Aare and the daily revenue of Bernese ice-cream vendors. Why can I be pretty confident about this? ◇

Inverted forks Figure 1.10 shows an inverted fork where two variables both influence a third one. The 'handle' of an inverted fork is called a **collider** since the two causal arrows clash into each other in A .

Table 1.1: Illustration of how a causal fork can give rise to associations between the variables at the prongs.

$\mathbb{P}(B \geq 1|C = 2)$ reads as 'the probability that B will be at least 1 when C equals 2.' This kind of probability is referred to as a conditional probability. If the variables B, C are not associated, then $\mathbb{P}(B \geq 1|C = 2) = \mathbb{P}(B \geq 1)$. Hence, if $\mathbb{P}(B \geq 1|C = 2) \neq \mathbb{P}(B \geq 1)$, as is the case here, B, C are associated.

We can't fix A at 1 and evaluate this probability at $C = 0$ for the simple reason that there's no row in the table with $A = 1$ and $C = 0$.




Figure 1.10: An inverted fork.

Inverted forks don't transmit causal influences between the variables at the prongs, that is, there is no causal link between B and C (causality doesn't travel upstream). The intriguing thing about inverted forks is this, though: When the collider (i.e., A) is *not* controlled for, the variables at the prongs remain unassociated. However, **controlling for the collider may induce an association between the variables at the prongs even in the absence of a causal link between them**. Controlling for a descendant of a collider may likewise induce an association between the variables at the prongs.

The effects of controlling for a collider are not intuitive, so let's consider an example.

Example 1.11. University teachers can testify that there is some negative association between their students' intelligence and their diligence. This doesn't mean that the most intelligent students are *all* lazy and none of the most diligent students are particularly clever—just that there is some tendency for the most intelligent students to be less hard-working than the less clever ones. There is a simple and plausible causal explanation for this association: The most intelligent students quickly figure out that they don't need to work as hard in order to obtain their degree, so they shift down a gear.

But there is an equally plausible if less simple explanation: by only looking at university students, we've controlled for a collider without realising it; see Figure 1.11. Even if diligence and intelligence were completely unassociated in the human population, they are bound to be associated if we only look at university students. Figure 1.11 illustrates why: If we consider the population as a whole, it's possible that there is no (or hardly any) association between diligence and intelligence (left panel): If we know a person's degree of diligence, we can't make a more educated guess as to their intelligence than if we don't. But if we only consider university students (filled circles in the right panel), we're bound to find a negative association between diligence and intelligence: If we know that a university student is pretty lazy, we also know that they need to be pretty intelligent—otherwise they couldn't have made it into university.




Figure 1.11: To the extent that diligence (work) and intelligence (IQ) both determine if someone gets into university, some association between these two factors will be found if we only look at university students.




Figure 1.12: Collider bias in action: If you only look at the filled (or at the unfilled) circles, you'll discover an association between diligence and intelligence, even though there is no causal link between them.

By only looking at university students, we've unwittingly controlled for a collider, which by itself can explain the negative association between diligence and intelligence observed among university students. This doesn't mean that our first causal explanation is necessarily wrong. But it does illustrate that there is a non-obvious but plausible additional explanation that we need to reckon with. It's also possible that both explanations are simultaneously correct: There may be some (negative) causal influence of intelligence on diligence, but by only looking at university students, we would then end up overstating the strength of this causal effect.

In Figure 1.11, we've assumed—for ease of exposition—that there is a perfect deterministic relationship between diligence and intelligence on the one hand and university enrolment on the other hand (if the sum of both scores is above 10, enrolment is granted). In reality, this relationship won't be perfect (some highly intelligent and highly diligent people don't go to university), but even so, controlling for (or 'conditioning on') a collider can produce associations between two factors in the absence of a causal link between them. ◇

Other examples of this collider bias are only superficially different from the example we've just considered. The difficulty lies in figuring out what collider was unwittingly controlled for.

Exercise 1.12. There is a negative association between how easily accessible a restaurant is from a tourist resort and how good the food is. Come up with an explanation that does not assume any direct or indirect causal influence of food quality on location or vice versa. ◇

Exercise 1.13. People with a highly active dating life sometimes complain that their hottest dates tend to be comparatively boring. Come up with an explanation that does not assume any direct or indirect causal influence of attractiveness on interestingness. ◇

In sum, unbroken chains both transmit causality and induce associations; forks induce associations without causality unless measures are taken (e.g., controlling for the confounder); and inverted forks induce associations without causality if the collider (or one of its descendants) is controlled for.

Exercise 1.14. Consider the DAG in Figure 1.13.

- (a) Can A causally affect D ?
- (b) Can there be an association between A and D if no factor is controlled for? If so, via which path?
- (c) Can there be an association between A and D if C is controlled for? If so, via which path?
- (d) Can there be an association between A and D if F is controlled for? If so, via which paths (plural!)?




Figure 1.13: DAG for Exercise 1.14.

- (e) Can there be an association between A and D if G is controlled for? If so, via which paths (plural!)?
- (f) Can there be an association between A and D if B and C are controlled for? If so, via which path?
- (g) Can C causally affect E ?
- (h) Can there be an association between C and E if no factor is controlled for? If so, via which path?
- (i) Can there be an association between B and E if no factor is controlled for? If so, via which path?
- (j) Can there be an association between B and E if C is controlled for? If so, via which path?
- (k) Can there be an association between B and E if D is controlled for? If so, via which path?
- (l) Can there be an association between B and E if D and F are controlled for? If so, via which path?
- (m) Can there be an association between B and E if D and G are controlled for? If so, via which path? ◇

Definition 1.15 (Bias). The term **bias** refers to a systematic distortion of the results, e.g., due to confounding variables. A single **unbiased** study isn't guaranteed to estimate the size of the causal effect correctly, but roughly speaking, if we were to run the same study lots of times, the under- and overestimates would cancel each other out. If the under- and overestimates don't cancel each other out, then the study is **biased**. ◇

***Remark 1.16.** The more formal mathematical definition is that an estimate is unbiased if its expected value equals the parameter value it attempts to estimate, and biased otherwise. ◇

If we want to obtain an unbiased estimate of the total causal influence of a variable X on another variable Y , we need to ensure

- that all directed paths from X to Y are open (i.e., no intermediate variable is controlled for);
- and that X and Y cannot be associated via any other paths.

Satisfying both conditions doesn't strictly guarantee that an unbiased estimate will be obtained as this also depends on the statistical procedure to obtain the estimate.

Exercise 1.17. Consider the DAG in Figure 1.14 and answer the following questions:

- (a) List all paths via which X may causally affect Y if no variables are controlled for.
- (b) List all paths via which X and Y may be associated if no variables are controlled for.
- (c) Which variable(s) do you *need* to control for if you want to obtain an unbiased estimate of the causal effect of X on Y ? Also




Figure 1.14: DAG for Exercise 1.17.

quickly check if controlling for these variable(s) doesn't open up any new paths via which X and Y may be associated! ◇

Sometimes, different sets of variables can be controlled for to this end. But in this case, there is just a single correct solution.

1.3 *Further reading

Rohrer (2018) is an accessible introduction to DAGs. I don't recommend you read it right away, though, but save it in case you need a refresher from a different source in a couple of months or years.

2

Constructing a control group

2.1 A made-up example

Imagine that a new self-learning method for fostering Danish reading skills in speakers of German has been developed. You're tasked with finding out if this new method works better than the old one.

First attempt You find four students of German philology who want to learn Danish. You ask them to work autonomously with the new learning method half an hour a day for three weeks. After three weeks, you give them an article from a Danish newspaper, which they are to summarise orally in German. Two raters judge these summaries at their own discretion (20-point scale); the mean of the two ratings per learner counts as their reading comprehension score. The average group score is 11/20.

What can you conclude from this study?

The answer is 'pretty much nothing'. One of several problems with this study is that there is no baseline against which to compare the participants' average result: We don't know whether 11/20 indicates that the new learning method works better or worse than the old one, or whether the old and new learning methods are roughly equally effective. So we need a comparison or **control group** of people that didn't take part in the intervention.

Second attempt You convince four law students to also take part in the study. They're asked to work with the old learning method half an hour a day for three weeks. Then they take the same test as the German philology students. Their group mean is 8/20.

What can you conclude from this study?

2.2 Critical questions

The second attempt outlined above also falls short on a number of criteria. There are a couple of critical questions we can ask, and slightly modified versions of these questions can be asked about studies in general.

Internal validity Can the difference in test scores between the two groups be ascribed to the difference in learning methods, or do alternative explanations for it present themselves?

External validity Does the finding apply only to the present **sample** or can it be generalised to a larger **population**? To what population, exactly?

Ecological validity (Especially for applied research.) To what extent are the findings applicable to the world outside of the lab (e.g., teaching, policy)?

Reliability Are the raw data open for interpretation?

Inter-rater reliability: Would different observers agree on the measurements?

Intra-rater reliability: Would the *same* observers agree on the measurements if they had to redo them?

Interpretative consensus I use this non-standard term to refer to the possible problem captured by the following question: Confronted with the same data, would other researchers draw similar conclusions?

Replicability Can the results of this study be confirmed in an independent **replication**, that is, in a new but otherwise similar study?

Example 2.1 (Ecological validity). Neurolinguistic research may show that using a foreign language highlights different brain regions in early and late learners. This may be interesting to neurolinguists. But for language policy, what's more relevant is the extent to which these groups differ behaviourally on communicatively relevant tasks. ◇

Example 2.2 (Interpretative consensus). A study with considerable internal, external and ecological validity could show that beginning L2 French learners vastly prefer using a rising intonation over other strategies for formulating questions, but that more seasoned learners show a more balanced use of the different strategies (subject-verb inversion, *est-ce que*). Researchers may agree on these results, but are bound to disagree on which implications, if any, they ought to have for L2 French teaching.

Generally, empirical findings obtained in basic research cannot be used as evidence for nor against some teaching or learning method. Such evidence must stem from targeted studies comparing teaching or learning methods. ◇

The labels above aren't important; the questions behind them are. Further, the questions asked above can rarely be answered in absolute terms.

Exercise 2.3. The second attempt outlined on page 19 is deficient in several of these respects. Discuss a few problems. ◇

Anticipate and resolve problems related to lacking validity and reliability *before* collecting the data. This often involves making compromises or coming to the realisation that you can't satisfactorily answer all your questions in a single study. Do *not* assume that some statistical method will solve your problems.

Depending on your goals, some types of validity may not be as important as others. For instance, for most studies in psycholinguistics, university students are recruited as participants, and their results don't necessarily generalise to the population at large. But the purpose of these studies is often to demonstrate that some experimental manipulation *can* affect language use and processing, not that it will affect language use in all language users. From this perspective, these studies' lack of external validity isn't too damning (Mook, 1983).

Exercise 2.4. Skim the article by Offerman & Olson (2016) and answer the following questions.

1. The study features two **conditions**: an experimental group and a control group. Briefly describe the difference between these conditions in terms of what the participants were asked to do. (Sections 3.1 and 3.3.)
2. Consider the first research question on p. 48: "Does a visual feedback paradigm improve production of the segmental feature voice onset time?"

Based on your answer to the previous question, complete the following:

"Does a visual feedback paradigm improve production of the segmental feature voice onset time compared to [YOUR ANSWER]?"

What would a stronger baseline for the comparison look like, and why?

3. Who were the participants to this study? Do the authors report how the participants were assigned to the study's conditions? (Section 3.1.)

How does this affect our (i.e., the readers') ability to judge the study's internal validity? ◇

Exercise 2.5. De Wilde et al. (2020) found that Flemish children already knew a decent number of English vocabulary items even before they took their first English class. Especially English words

similar in both form and meaning to their Dutch translations (i.e., cognates) tended to be commonly known among these children. In their conclusion, the authors write that “[t]he study also suggests that (...) teachers do not need to pre-teach some cognate words, even for learners at an elementary level” (pp. 374–375).

De Wilde et al. (2020) didn’t empirically investigate the effects of implementing this suggestion. But can you think of an undesirable consequence of not ‘pre-teaching’ English words with true Dutch cognates but still teaching false friends, i.e., English words for which a Dutch word with a similar form but a different meaning exists (e.g., *room*, which means ‘cream’ in Dutch)? ◇

2.3 *Increasing internal validity through randomisation*

Of the three types of validity considered, internal validity is the most pressing one: If internal validity is low, external and ecological validity are both essentially irrelevant. Hence, our first priority is to maximise the study’s internal validity, that is, we want to maximise the chances that any association we find in the data can be ascribed to the factor of interest. Confounding in particular represents a substantial threat to internal validity: As we’ve seen in Chapter 1, confounding variables induce associations between the variables of interest even in the absence of a causal link between them. Moreover, even if a causal link does exist between the variables of interest, confounding variables can bias the association between them: The association may systematically under- or overestimate the strength of the causal link. Keeping confounding in check is therefore key.

Your first inclination may be to try to ensure that the intervention and control groups are identical in all respects save for the treatment itself. That way, any differences in the outcome variable can’t be explained by confounding due to pre-existing differences between the groups. However, it is practically impossible to assign a fixed number of participants to two groups in such a way that these groups are identical in all respects even in the utterly unrealistic case where all the relevant information is available beforehand. Clearly, it’s entirely impossible to do so when not all of the relevant information is available beforehand.

The solution is to assign the participants to the study’s conditions **at random**, i.e., to deliberately leave the allocation up to chance and chance alone. The DAGs in Figure 2.3 on the next page show what such randomisation achieves. When the participants themselves (or their parents, or their circumstances, etc.) decide or otherwise affect which condition they end up in (X), confounding is a genuine concern (left). However, when we assign the participants to the conditions at random, we *know* that there is no systematic link between pre-existing characteristics (U) and X , let alone a causal one. That is, randomisation prevents any causal arrows from entering




Figure 2.1: *Left*: The X - Y relationship is confounded by U : there are two paths from X to Y , but only one causal one. *Right*: Randomising the values of X prevents arrows from U entering X , which effectively closes the non-causal path via the confounder.

X (right)! The result of this is that the non-causal path between X and Y (via U) is broken and that the X - Y relationship is no longer confounded by U . To highlight that the values of X were specified at random, we use a subscript r .

Studies in which the participants are randomly assigned are called **true experiments**. Random allocation by itself doesn't guarantee that the results of the experiment can be trusted or interpreted at face value, but it does eliminate confounding as a threat to the study's internal validity.

Randomise wherever possible – unless you have a *very* good reason not to!

Tip 2.6. When reading empirical research articles, always look up if and how the participants (or classes, or whatever is of interest) were assigned to the study's conditions. And unless it explicitly says that the assignment was carried out at random, do not assume that it was. ◇

2.3.1 Why experiments?

1. "Experiments allow us to set up a **direct comparison** between the treatments of interest.
2. "We can design experiments to **minimize any bias** in the comparison. [especially randomisation]
3. "We can design experiments so that the **error** in the comparison is **small**. [see later chapters]
4. "Most important, we are **in control** of experiments, and having that control allows us to make stronger inferences about the nature of differences that we see in the experiment. Specifically, we may make **inferences about causation**." (Oehlert, 2010, p. 2, my emphasis)

What's meant by the first point is that we don't need to piece together snippets of evidence from different and often difficult-to-reconcile sources (e.g., in a literature review) to figure out which treatment (e.g., which learning method) works best in some situation. Instead, we set up an experiment that directly evaluates the efficacy of several treatments in the context we're interested in.

The second point alludes to techniques such as randomisation; the third to techniques that we will encounter in the chapters to come.

2.3.2 What does randomisation do?

1. “Randomization balances the population on average.”
2. “The beauty of randomization is that it helps prevent confounding, *even for factors that we do not know are important.*” (Oehlert, 2010, p. 15, my emphasis)

We’ve already discussed the second point, but the first point warrants some explanation. Some fundamental results from combinatorics are useful at this stage.

Lemma 2.7. If you have $n \geq 0$ distinct objects, there are

$$\begin{cases} n(n-1) \cdots 2 \cdot 1, & \text{if } n \geq 1, \\ 1, & \text{if } n = 0. \end{cases}$$

different ways to sort them (**‘permutations’**). $\diamond$

Proof. If $n = 0$, there is only one way to sort the objects: do nothing.

If $n \geq 1$: There are n candidates to fill the first slot. Once a candidate for the first slot has been picked, $n - 1$ candidates remain for the second slot, and so on, till there is only a single candidate left for the final slot. $\square$

Definition 2.8 (Factorial). For any natural number n , we define

$$n! := \begin{cases} n(n-1) \cdots 2 \cdot 1, & \text{if } n \geq 1, \\ 1, & \text{if } n = 0. \end{cases}$$

We say ‘ n factorial’ for ‘ $n!$ ’. $\diamond$

Lemma 2.9. There are

$$\frac{n!}{k!(n-k)!}$$

different ways to choose $k \geq 0$ objects from among $n \geq k$ distinct objects. $\diamond$

Proof. With Lemma 2.7, there are $n!$ ways to arrange the $n \geq 0$ distinct objects. For each possible rearrangement, pick the first k objects. This covers all possible choices of k from n objects. However, several rearrangements result in the same selection. Specifically, if we rearrange the first k elements and the last $n - k$ elements separately from each other, we end up with the same selection: the order among the first k and among the last $n - k$ doesn’t matter. Hence, again with Lemma 2.7, there are, for each of the $n!$ permutations, $k!(n - k)!$ rearrangements that result in the same selection. So there are $\frac{n!}{k!(n-k)!}$ different possible selections. $\square$

Definition 2.10 (Binomial coefficient). For natural numbers n and k with $n \geq k$, we define

$$\binom{n}{k} := \frac{n!}{k!(n-k)!}.$$

This script includes a few proofs. You don’t have to learn them as such (I won’t ask for proofs on the exam), but do try to understand them. The reason why I included them is that these proofs help to demystify what might otherwise seem to be arbitrary pronouncements. Further, these proofs clarify exactly which assumptions a claim depends on and how far it extends: a proof can show you that an argument holds regardless of some detail that you might otherwise have assumed was essential to it.

A term of this form is called a **binomial coefficient**; we say ‘ n choose k ’ for ‘ $\binom{n}{k}$ ’.



Let’s say that we have ten participants and we know both their sex and their IQ (Figure 2.2). We can pick five among them at random and assign them to the first condition; the others are assigned to the second condition. By the previous lemma, there are

$$\binom{10}{5} = \frac{10!}{5!(10-5)!} = \frac{3628800}{(120)^2} = 252$$

different choices for the participants in the first group; each of these choices has an equal probability of being our allocation. Six out of the 252 possible allocations are shown in Figure 2.10. Note that in none of them, the intervention and control groups are perfectly balanced with respect to both IQ and sex. So randomisation clearly does not generate balanced groups in any particular study. However, each participant is as likely to end up in the intervention group as they are to end up in the control group, so *on average*—across all 252 possible allocations—sex, IQ, as well as all unmeasured variables, are balanced between the two groups. For our present purposes, this means that randomisation is an equaliser: the result may not be two perfectly equal groups, but at least one group isn’t systematically given an advantage relative to the other. As we’ll see in Chapter 4, randomisation also justifies the use of some common statistical procedures.

The binomial coefficient is called so because these terms appear when expanding binomial expressions such as $(a + b)^n$. In German, one usually says ‘ n tief k ’ or ‘ k aus n ’ for ‘ $\binom{n}{k}$ ’; in French, ‘ k parmi n ’.




Figure 2.2: Ten participants sign up for a study. You measure their IQ and you also know their sex (represented here using circles and crosses).




Figure 2.3: Six possible random assignments (out of 252) of the ten participants from Figure 2.2. The dotted vertical lines show the mean IQ in each group.

Exercise 2.11. For each description, decide whether the participants were randomly assigned to the experiment’s conditions and, if not, explain how the lack of randomisation could result in confounding.

- (a) 60 participants trickle into the lab. The first 30 are assigned to the experimental condition, the final 30 are assigned to the

control group.

- (b) Experiment with a school class: Pupils whose last name starts with a letter between A and K are assigned to the control group, the others to the experimental group.
- (c) Participants come to the lab one by one. For each participant, the researcher throws a dice. If the dice comes up 1, 2, or 3, the participant is assigned to the experimental condition; if it comes up 4, 5, or 6, the participant is assigned to the control condition. After four weeks, no more participants sign up. The control group consists of 17 participants; the experimental group of 12.
- (d) Participants come to the lab one by one. The first participant is assigned to the experimental condition, the second to the control condition, the third to the experimental condition, the fourth to the control condition, etc. Of the 29 participants who signed up in total, 15 are assigned to the experimental condition and 14 to the control condition.
- (e) To investigate the effects of bilingualism on children's cognitive development, 20 bilingual 4-year-olds (10 girls, 10 boys) are recruited. 20 monolingual 4-year-olds (10 girls, 10 boys) serve as the control group.
- (f) 32 participants sign up for an experiment. The researcher enters their names into <http://www.random.org/lists/>, clicks Randomize and assigns the first 16 to the control group and the others to the experimental group. ◇

'Random' does not mean 'haphazard', 'arbitrary' or 'at the researcher's whim'.

Remark 2.12 (Randomisation and estimation). When participants are assigned randomly to the experiment's conditions, the results of the experiment are also subject to randomness. To appreciate this, consider an experiment for which sixteen participants sign up. The left-side panel in Figure 2.12 shows the score that each participant would have obtained in both conditions (often referred to as their **potential outcomes** in the literature on causal inference; see Definition 2.13 on the next page). The difference between both of these scores varies somewhat between participants, but on average, it amounts to one point. This is the true average intervention (or treatment) effect that is the target of the experiment.

Depending on the condition they are assigned to, we would only observe one of the scores for each participant, however. If we randomly assign nine of the sixteen participants to the intervention condition and seven to the control condition, we end up with one out of $\binom{16}{9} = 11,440$ possible allocations. To estimate the true average intervention effect, we can compute the mean difference between the scores of the participants assigned to the intervention

condition and those assigned to the control condition. The estimated intervention effect would then be one of the 11,440 values summarised in the histogram on the right-hand side. While the *average* estimated intervention effect coincides with the true intervention effect (i.e., the estimation is unbiased; see Theorem 2.14), depending on the actual allocation, the estimated intervention effect may differ considerably from the true one. In Chapter 5, we'll discuss techniques for increasing the precision of the estimation. ◇




Figure 2.4: *Left*: The scores that each of the sixteen participants would have obtained if they had been assigned to the control group (circles) or to the intervention group (crosses). The average difference between the intervention and the control scores, i.e., the average intervention effect, is one point. *Right*: The histogram shows the distribution of the mean difference between the intervention and control scores for each of the $\binom{16}{9} = 11440$ possible allocations of participants to the two conditions. Averaged over all 11,440 allocations, the mean difference is also one point. But for many individual allocations, the mean difference may differ considerably from the true average intervention effect of one point.

Definition 2.13 (Potential outcomes). Consider a study in which N units are each observed in one of K conditions (numbered $0, 1, \dots, K-1$). We denote by $Y_i(k)$ the value on the outcome variable that was observed for the i -th unit assigned to the k -th condition or that would have been observed for this unit had it been assigned to the k -th condition, $i = 1, \dots, N, k = 0, \dots, K-1$. These values, which may be unknown, are called **potential outcomes**. ◇

Theorem 2.14. When using random assignment, the mean difference in the outcome between the control and intervention groups averaged over all possible allocations equals the true average (mean) intervention effect. ◇

Proof. Assume an experiment with m units of which n_1 are randomly assigned to the intervention condition and $n_0 = m - n_1$ are assigned to the control condition; i.e., there are $\binom{m}{n_1} = \binom{m}{n_0}$ possible allocations. The true average intervention effect is defined as the mean of the m per-unit differences between the potential outcomes in the intervention and in the control condition. Equivalently, we can compute it as the difference between the mean of the m potential outcomes in the intervention condition and the mean of the m potential outcomes in the control condition.

The estimate we obtain from any particular random allocation is computed as the difference between the mean outcome among the n_1 units assigned to the intervention condition and the mean outcome among the n_0 units assigned to the control condition. Across all possible allocations, every unit is assigned to the intervention condition equally often; this works out to $\frac{n_1}{m} \cdot \binom{m}{n_1} = \binom{m-1}{n_1-1}$ times.

Similarly, every unit is assigned to the control condition equally often; this works out to $\frac{n_0}{m} \cdot \binom{m}{n_0} = \binom{m-1}{n_0-1}$ times.

The mean intervention outcome averaged over all $\binom{m}{n_1}$ allocations is therefore the sum of $\binom{m-1}{n_1-1}$ times each intervention potential outcome divided by $n_1 \cdot \binom{m}{n_1}$. You can verify that

$$\binom{m-1}{n_1-1} / \left(n_1 \cdot \binom{m}{n_1} \right) = \frac{1}{m}.$$

So the result is the mean potential intervention outcome over all m units. Similarly, the mean control outcome averaged over all $\binom{m}{n_0}$ allocations is the mean potential control outcome over all m units. Hence, averaged over all $\binom{m}{n_1} = \binom{m}{n_0}$ allocations, the mean difference between the groups equals the true average intervention effect. $\square$

Remark 2.15. Theorem 2.14 concerns the *mean* difference and the average (i.e., *mean*) intervention effect. The statement doesn't generalise to, say, medians. To see this, consider an experiment with three units whose potential outcomes in the intervention group are 0, 0, 3 and whose potential outcomes in the control group are 0, 0, 0. The median intervention effect for these units is 0. However, if we randomly assign one unit to the intervention group and two to the control group (three possibilities), the average median difference between the two groups is $\frac{0+0+3}{3} = 1 \neq 0$.

This counterexample demonstrates that the proof of Theorem 2.14 cannot be adapted to medians. In the proof, we first rewrote the mean of the intervention effects as the difference between two means; this doesn't work for the median. We also were able to compute the average mean intervention and control outcomes as averages over $n_1 \cdot \binom{m}{n_1}$ and $n_0 \cdot \binom{m}{n_0}$ potential outcomes, respectively. This doesn't work for the average median outcome. $\diamond$

2.3.3 How to randomise?

When collecting data using computers Have the computer randomly assign the participants to the conditions without your involvement. Frameworks for running experiments such as OpenSesame (<https://osdoc.cogsci.nl/>), PsychoPy (<https://www.psychopy.org/>) or jsPsych (<https://www.jspsych.org/>) all contain functions for allocating participants randomly.

When the data collection does not take place at the computer and you know who'll be participating beforehand Randomise the list of participants using <https://www.random.org/>. Assign the first half of the list to the experimental condition and the second half to the control condition. If the experiment features $k \geq 3$ conditions, similarly split up the list into k parts.

This procedure is known as **complete randomisation**. It guarantees that the number of experimental units is the same in each

condition (or at most one off if the number of units isn't divisible by the number of conditions).

When the data collection does not take place at the computer and you don't know who'll be participating beforehand Randomly assign each participant individually and with the same probability to a condition as they sign up. This procedure is known as **simple randomisation**. In contrast to complete randomisation, you're not guaranteed to end up with an equal number of units in each condition. This is usually of little concern, and in fact, simple randomisation arguably reduces the potential for the researchers' biases to affect the study's results (Kahan et al., 2015). Importantly, **there is nothing wrong with having unequal sample sizes**.

Humans make for poor randomisation devices. Always randomise mechanically (preferably with a computer).

More complex allocation procedures are possible. Some of these are common in medical research (Rosenberger & Lachin, 2016), but they are rarely applied in the social sciences.

Exercise 2.16 (Complete randomisation with several conditions). Consider an experiment with 60 participants who are assigned to three conditions using complete randomisation (20 participants per condition). How many possible assignments of participants to conditions are there under this scheme? ◇

Exercise 2.17 (Simple randomisation). Consider an experiment with 40 participants, each of which is assigned to one of the conditions using simple randomisation. How many possible assignments of participants to conditions are there under this scheme? Without computing the value of $\binom{40}{20}$, is the number you've obtained smaller or larger than $\binom{40}{20}$? ◇

The word 'experiment' can be used in a stricter or in a looser sense. The mere fact that a study is referred to as an 'experiment' does *not* mean that it's a *true experiment* (control group + randomisation): the use of the label doesn't automatically imply that confounding has been taken care of. Many quantitative studies in our field aren't experiments in the strict sense. Always look up how the participants were assigned to the conditions!

See blog entry *Causes and consequences of unequal sample sizes*.

Incidentally, medical researchers refer to simple randomisation as 'complete randomisation'. It's probably best to just spell out how the randomisation was carried out rather than to rely on short-hand terminology.

3

Alternative explanations

As we've seen in Chapter 1, causal explanations for observed associations become more tenuous to the extent that more, and more plausible, alternative explanations for the associations exist. In Chapter 2, the random assignment of units (e.g., participants) to conditions was introduced as a crucial technique for neutralising the danger that an association will come about through systematic confounding. The present chapter investigates some other plausible rival explanations.

3.1 The roles of variables in research

It is useful to clarify some terminology at this point.

Definition 3.1. An **outcome variable** (also called **dependent variable**) is a variable whose value we want to explain, predict, or otherwise characterise.

A predictor variable (also called **independent variable**) is a variable that is used to explain, predict, or otherwise characterise an outcome variable. In experiments, such variables are 'manipulated' by the researchers.

A **control variable** is a variable that is not of primary interest but that is collected in order for the researchers to be able to account for alternative explanations or to otherwise improve the study. We'll discuss the usefulness of control variables in a later chapter. ◇

Exercise 3.2. Skim Ludke et al. (2014) in the light of the questions below and briefly answer them. The answers to these questions can be found in the 'Method' section (pp. 43–47), so you don't have to read the rest.

- (a) What is or what are predictor variable(s) in Ludke et al.'s study?
- (b) What is or what are outcome variable(s) in Ludke et al.'s study?
- (c) How did the researchers assign the participants to the conditions? ◇

In the study by Ludke et al. (2014) (see Exercise 3.2), the question

about internal validity boils down to this: Can the differences in the outcomes between the conditions actually be ascribed to the difference between the conditions (control vs. intervention; singing vs. rhythmic speaking vs. speaking)? Or are there other explanations for it?

In Chapter 2, we focused on the threat that confounding poses to a study's internal validity and how this threat can be neutralised using randomisation. We saw that randomised ('true') experiments (probabilistically) negate the influence of confounding variables on the results: one group isn't systematically given an advantage compared to the other (e.g., higher motivation, greater affinity with a topic etc.). This increases the study's internal validity, but:

Even if confounding variables are taken into account, other systematic factors may give rise to a spurious difference between the experiment's conditions or may mask an existing effect of the conditions.

3.2 *Explanation 1: Expectancy effects*

Perhaps the researchers or their assistants (subconsciously) nudged the data in the hypothesised direction. This can happen even when the measurements seem perfectly objective. For instance, when you're counting the number of syllables in a snippet of speech, there are bound to be a number of close decisions (Does German *haben* [ha(b)m] have one or two syllables?). This isn't too big a problem in itself. But it does become a cause for concern if you tend to reach different decisions depending on which condition the participant was assigned to.

Relatedly, it's possible that the participants want to help (or thwart) the researchers achieve what they think are the researchers' goals. In this case, differences in the outcome variable between the conditions may arise not because of the intervention itself but because of unwanted changes in the participants' behaviour. Such changes in behaviour needn't come about consciously and are known as **expectancy effects**.

Expectancy effects can be countered by **blinding**. The term **single-blind experiment** is typically used to describe an experiment in which the participants don't know which condition they're assigned to, whereas the term **double-blind experiment** is used to describe experiments in which neither the participants nor the researchers themselves (at the time of collecting and preparing the data) know which condition the participants were assigned to. That said, these terms are often used in ambiguous ways so, when you use blinding, it's best to simply explain how you achieved it.

Blinding isn't always possible, and it may be immediately obvious to the participants what the intervention entails. But in stud-

ies with raters, it's usually easy to prevent them from knowing which condition the participants were assigned to. It is also possible (though rarely done in the social sciences) to apply blinding during the analysis of the data (Dutilh et al., 2021).

Example 3.3. Ludke et al. (2014) obtained the raw data from phonetic transcriptions of their participants' utterances without knowing which conditions the participants had been assigned to (p. 46, bottom right). This way, close decisions as to whether a response was good enough were less likely to contribute to a difference between the conditions. ◇

3.3 *Explanation 2: Failed manipulation*

A second class of alternative explanations is that the experiment didn't run quite as the researchers expected it to. For instance, the participants may have misunderstood, or failed to act on, the instructions, or the script used to run the experiment could have contained a crucial coding error.

One tool to uncover discrepancies between how the experimenters intended the instructions and how the participants interpreted them is to include **manipulation checks** in the study's design.

Example 3.4. The experimental manipulation in Ludke et al. (2014) involved asking the participants to practice Hungarian phrases while singing or rhythmically pronouncing them. In order to verify whether the participants followed these instructions, the authors recorded the participants as they practiced the phrases. ◇

Example 3.5. Lardiere (2006) had her (single) participant judge sentences in her second language for their grammaticality. The goal of the study was to figure out if this participant had acquired some syntactic property of the second language. To ensure that the participant rejected any sentences for the intended syntactic reason, she was also asked to correct any sentences she rejected. This turned out to be an excellent idea: The researcher found that the participant rejected a fair number of syntactically correct sentences, but that she did so for stylistic rather than strictly syntactic reasons. The researcher then correctly didn't draw the conclusion that the participant's syntactic knowledge was incomplete. ◇

Sometimes, the participants in a study don't really pay any attention to the stimuli or the instructions. For instance, questionnaire respondents may answer the questions in a zig-zag pattern (e.g., ABCDEDCBA...) rather than give their mind to each question, so that they can tell the researcher they've participated. Figure 3.5 provides another example of satisficing.

If you want to verify if your participants carefully read the instructions and you're able to run the study online or at the com-




Figure 3.1: A large number of raters were asked to each rate about 50 short texts on a 9-point scale. These two clearly lost interest at some point (Vanhove, 2017).

puter, Oppenheimer et al. (2009) introduces a neat and unintrusive way to do so.

Another useful technique to identify failed manipulations is the use of **positive controls**. A positive control is a condition, measure, or comparison that should show an effect if the study is functioning properly. Positive controls can therefore help identify failed manipulations.

Example 3.6. Let's say that a researcher wants to find out if knowledge of L2 verb morphology contributes to L2 French speakers' comprehension of temporally complex texts. To this end, she proposes an experiment that involves the intensive teaching of French verb morphology.

In this case, it would make sense for the researcher to check if the intervention does in fact succeed in improving the participants' command of French verb morphology. If it doesn't, then the results of the study can't speak to the link between knowledge of verb morphology and text comprehension. ◇

Example 3.7. In L2 research, the task given to the L2 speakers is often also given to a group of L1 speakers. This helps identify issues in the materials used. For example, if sentences meant to be grammatically correct are frequently rejected by L1 speakers and L2 speakers alike (perhaps due to awkward style), it suggests the problem lies with the stimuli, not necessarily with the L2 speakers' abilities. ◇

The term **negative control**, incidentally, refers to traditional control groups (of which we know that they shouldn't show an effect of the intervention).

The final tool is perhaps the most important one: **pilot studies**. A pilot study is a small-scale study conducted before the main study. Typical goals of a pilot study are

- (a) ensuring that the participants understand and act on the instructions;
- (b) identifying any remaining glitches in the experimental software or in the analysis pipeline;

- (c) if relevant, checking whether the responses obtained can be coded satisfactorily;
- (d) estimating how much time participants need to complete the study. This is useful for recruiting participants and scheduling;
- (e) detecting any **floor** or **ceiling effects**. A floor effect occurs when a task is so difficult that many participants score near the bottom; a ceiling effect occurs if a task is so easy that many participants score near the top. The problem with floor and ceiling effects is that participants may differ substantially in their ability yet obtain the same score.

3.4 *Explanation 3: Chance*

A third important possible non-causal explanation for one's results is that they're due to chance. The entire next chapter is devoted to an attempt to get a handle on this explanation.

4

Inferential statistics 101

If you've read a few quantitative research articles, you've probably encountered p -values. This chapter explains the basic logic behind p -value-based inferential statistics. It does so by explicitly linking the computation of p -values to the random assignment of participants to conditions in experimental research. If you have ever taken an introductory statistics class, chances are p -values were explained to you in a different fashion, presumably by making assumptions about how the observations in the sample were sampled from a larger population and by making reference to the Central Limit Theorem. For the explanation in this chapter, however, we're going to take a different tack and we will ignore the sampling method and the larger population. Instead, we're going to leverage what we know about how the observations, once collected, were *assigned* to the different conditions of an experiment. The advantages of this approach are that it connects the design of a study more explicitly to the analysis of its data and that it is less math-intensive while permitting one to illustrate several key concepts about inferential statistics.

The goal of this chapter is for you to *understand conceptually* what statistical tests attempt to achieve, not for you to be able to use them yourself. As a matter of personal opinion, statistical tests are overused (Vanhove, 2021). I think that, in your own research, your focus should be on describing your data (e.g., by means of appropriate graphs) rather than running umpteen significance tests. Analysing data and running statistical tests are not synonymous.

4.1 An example: Does alcohol intake affect fluency in a second language?

Research question Does moderate alcohol consumption increase verbal fluency in a foreign language (L2)?

Method Nine students (L1 German, L2 English) are randomly assigned to either the control (four) or the experimental condition (five); they don't know which condition they're assigned to. Participants in the experimental condition drink one pint of ordinary beer;

Nine participants is obviously a very low number of participants, but it keeps things more tractable here.

those in the control condition drink one pint of alcohol-free beer.

Afterwards, they watch a video clip and relate what happens in it in English. This description is taped, and two independent raters who don't know which condition the participants were assigned to count the number of syllables uttered by the participants during the first minute. The mean of these two counts serves as the verbal fluency/speech rate variable.

Results The measured speech rates are shown in Figure 4.1. On average (mean), the participants in the *with alcohol* condition uttered 4.42 syllables/second, compared to 3.60 syllables/second in the *without alcohol* condition. That is, the mean difference in favour of the *with alcohol* condition is 0.82 syllables/second.

Table 4.1 lists the precise values for good measure. This table introduces the shorthand $Y_i(0)$ to refer to the i -th participant's speech rate if this participant was assigned to the *without alcohol* condition (e.g., Lukas' speech rate is $Y_7(0) = 3.7$) and the shorthand $Y_i(1)$ if the participant was assigned to the *with alcohol* condition (e.g., Daniel's speech rate is $Y_5(1) = 3.8$).

i	Participant	Without alcohol ($Y_i(0)$)	With alcohol ($Y_i(1)$)
1	Sandra		5.6
2	Nicole		4.4
3	Maria		4.2
4	Yves		4.1
5	Daniel		3.8
6	Nadja	4.2	
7	Lukas	3.7	
8	Thomas	3.4	
9	Michael	3.1	




Figure 4.1: Individual results of a randomised experiment.

Table 4.1: The measured speech rates (syllables/second) in the fictitious alcohol-verbal fluency experiment.

4.2 Chance as an alternative explanation

The design of this fictitious experiment has deftly dealt with two major threats to internal validity: confounding has been neutralised by means of randomisation, and expectancy effects have been kept in check through blinding. But there is another threat to internal validity that we need to reckon with: While we observed a mean difference between the two conditions of 0.82 syllables/second in favour of the *with alcohol* condition, this difference could have come about through *chance*. We are, then, faced with two types of account for this mean difference:

- The result we've obtained is due *only* to chance.
- The result we've obtained is due to chance *and* systematic factors.

Frequentist inferential statistics seeks to quantify how surprising the results are if we assume that only chance is at play. To do so, it attempts to answer the following key question: **How likely**

is it that a difference at least this large would've come about if chance alone were at play?

If it's pretty unlikely that chance alone would give rise to at least the difference observed, then this can lead one to revisit the assumption that the results are due only to chance—perhaps some systematic factors are at play after all. By tradition, the threshold between 'pretty unlikely' and 'still too likely', usually written as α , is 5%, but there is nothing special about this number. If the result falls below this 5% threshold, the difference is said to be 'statistically significant'. This is just a phrase, however, and arguably a poorly chosen one: statistical 'significance' doesn't tell you anything about a result's practical or theoretical import.

Tip 4.1. Avoid using the words 'significance' and 'significant' in their non-technical sense when writing about quantitative research. ◇

4.3 The null hypothesis

The first type of account, namely that the result obtained is due solely to chance, is known as the **null hypothesis** (H_0); the second as the **alternative hypothesis** (H_A).

A particularly strong version of the first account is that experimental conditions do not differentially affect *any* of nine participants' speech rates. This version, which is due to Fisher (1935), is known as the **strong** (or **sharp**) **null hypothesis**.

Let us for a moment assume that this strong H_0 is correct. In this case, the participants' speech rates aren't affected at all by the condition (*without alcohol* vs *with alcohol*) that they were assigned to. For instance, Sandra was assigned to the *with alcohol* and had a speech rate of $Y_1(1) = 5.6$. But under the strong H_0 , had she been assigned to the *without alcohol* condition, her speech rate would have been $Y_1(0) = 5.6$, too. Similarly, under the strong H_0 , we can use the speech rate that Michael had in the *without alcohol* condition ($Y_9(0) = 3.1$) to reconstruct the speech rate that he would have had in the *with alcohol* condition, namely $Y_9(1) = 3.1$.

The strong null hypothesis allows us to fill out the empty cells in the results table; see Table 4.3. The putative results are known as potential outcomes; see Definition 2.13 on page 27.

Definition 4.2 (Strong null hypothesis). Under the **strong** (or **sharp**) **null hypothesis**, we have $Y_i(0) = Y_i(1) = \dots = Y_i(K-1)$ for $i = 1, \dots, N$. ◇

i	Participant	Without alcohol ($Y_i(0)$)	With alcohol ($Y_i(1)$)
1	Sandra	5.6	5.6
2	Nicole	4.4	4.4
3	Maria	4.2	4.2
4	Yves	4.1	4.1
5	Daniel	3.8	3.8
6	Nadja	4.2	4.2
7	Lukas	3.7	3.7
8	Thomas	3.4	3.4
9	Michael	3.1	3.1

Under the strong H_0 , we have $Y_i(0) = Y_i(1)$ for $i = 1, \dots, 10$. The difference between the means of the $Y_i(0)$ and of the $Y_i(1)$ values, then, is zero. Still assuming that the strong H_0 is correct, then, the difference in speech rate between the two conditions that we actually observed must be due solely to the random assignment of participants to conditions, due to which more fluent talkers ended up in the *with alcohol* condition. For instance, another roll of the dice could have assigned Sandra to the control condition instead of Michael, and under the strong null hypothesis, this would have produced a slower mean speech rate in the *with alcohol* condition than in the *without alcohol* one (3.9 vs. 4.2; see Figure 4.2).

4.4 Testing the null hypothesis by exhaustive rerandomisation

With 9 participants split up in a control group of 4 and an experimental group of 5 there were $\binom{9}{5} = 126$ possible assignments of participants to conditions, each of which was equally likely to occur. To see how easily a difference as least as large as the one observed (a mean difference in speech rate of 0.82 in favour of the *with alcohol* condition) could occur due to random assignment alone, we can re-arrange the participants' speech rates into each of these 126 combinations and see for each combination what the difference between the *with* and *without alcohol* condition means is (Figure 4.4).




Table 4.2: The potential outcomes in the fictitious alcohol-verbal fluency experiment under the strong null hypothesis




Figure 4.2: If only chance were at play, Michael's (3.1) and Sandra's (5.6) results would be unaffected by the experimental condition and the outcome might equally well have looked like this (swapping Michael and Sandra).

Figure 4.3: There are 126 different ways in which the 9 participants could have been split up into two groups so that there are five participants in the experimental group. These are the differences between the means for all 126 possibilities under the strong null hypothesis. The mean difference we actually observed (0.82) is represented by the dashed red vertical line.

In five out of 126 cases, the re-arrangement of participants to conditions produces a mean difference in speech rate in favour of the *with alcohol* group that is at least as large as the one we actually observed (0.82). In other words, under the strong null hypothesis, the probability with which we would observe a mean difference of at least 0.82 in favour of the *with alcohol* condition if chance alone (in the form of random assignment) were at play is $\frac{5}{126} \approx 0.0397$ (3.97%). This quantity is known as the **p-value**. The procedure outlined in this section to obtain such *p*-values is called **randomisation testing**.

Using the threshold $\alpha = 0.05$, we would find that $p = 0.0397 < \alpha$ and conclude that a difference of 0.82 or more in favour of the *with alcohol* condition is fairly unlikely to occur under the strong null hypothesis. This would provide some evidence that there are systematic factors at play in producing the mean difference of 0.82. Since the study's design was specifically intended to minimise major threats to internal validity, it could then plausibly be argued that moderate alcohol consumption increases verbal fluency in the L2 in these participants.

Remark 4.3 (Directional and non-directional tests). In the example above, the research question was if moderate alcohol consumption *increases* L2 verbal fluency. This is a **directional** research question, and the statistic that we used to compare the groups reflects the direction of this research question: we computed the mean difference in speech rate between the two conditions in favour of the *with alcohol* group.

How would we have proceeded if the research question had been if moderate alcohol consumption *decreases* L2 verbal fluency? In this case, we could have computed the mean difference in speech rate between the two conditions but this time in favour of the *without alcohol* group. Proceeding entirely analogously, we would have found that, under the strong null hypothesis, 123 out of 126 of the mean differences would have been larger than the actually observed mean difference of -0.82 ($p = \frac{123}{126} \approx 0.98$). This would have given us no evidence for an alcohol-induced decrease in L2 verbal fluency.

If we hadn't committed to a specific direction, the research question should have been formulated in a **non-directional** manner: Does moderate alcohol consumption *influence* (or affect, change, etc.) L2 verbal fluency? One suitable statistic to compute in this case would have been the *absolute* mean difference between the conditions. The analysis would then have shown that, under the strong null hypothesis, we would have obtained an absolute mean difference as least as large as the one actually obtained (namely 0.82) in 9 out of 126 cases ($p = \frac{9}{126} \approx 0.071$). Using the threshold $\alpha = 0.05$, we would then have surmised that the study's results are fairly compatible with the strong null hypothesis.

The choice of the statistic chosen to compare the conditions in a

statistical test, then, affects the outcome of this test. Moreover, as we will see in more detail in a moment, it is imperative that the choice of this statistic is settled in advance, i.e., before analysing the data. ◇

Exercise 4.4 (Other test statistics). In your own words, outline how you can compute a p -value for the difference between the condition medians in the example above. Assume that the research question still is if moderate alcohol intake increases L2 verbal fluency. ◇

Exercise 4.5 (More than two conditions). Let's say we want to conduct an experiment with twelve participants who are randomly allocated to three conditions with four participants each. (Again, the numbers are kept low to keep exhaustive rerandomisation feasible.) With three conditions, we could compute the three pairwise differences between the condition means. But if we want to capture the differences between these in a single number, we could instead compute, for instance, the standard deviation of the condition means (see Exercise A.1 on page 93). Standard deviations can't be negative. Under H_0 , you'd expect the condition means to be close to one another, so this standard deviation would be close to 0; under H_A , you'd expect it to be higher.

- (a) How many possible ways are there to split up twelve participants into three groups of four participants each?
- (b) Explain how you can compute an appropriate p -value using exhaustive rerandomisation in this setting.
- (c) Now assume that we're interested in the group medians rather than in the group means. What would you need to change in your answer to (b)? ◇

***Remark 4.6** (One-sided and two-sided tests). The content of Remark 4.3 is more typically phrased in terms of one-sided and two-sided tests and the resulting left-, right-, and two-sided p -values. In the present example, we first computed a right-sided test: in Figure 4.4, we counted the number of rerandomisations in which the mean difference was equal to or lay to the right of the mean difference that we actually observed. Denote the resulting p -value by p_r .

To compute the left-sided p -value, we would have computed the number of rerandomisations in which the mean difference was equal to or lay to the left of the mean difference that we actually observed. The result would have been a p -value of $p = \frac{123}{126} \approx 0.98$, exactly as we obtained when changing the direction of the mean difference in Remark 4.3. Denote the left-sided p -value by p_ℓ .

A two-sided p -value p_t could have been computed by using the absolute mean difference, as in Remark 4.3. A slightly better alternative is to compute it as

$$p_t := \min(2 \cdot \min(p_\ell, p_r), 1).$$

In 'nice' cases, both calculations yield the same result. But this isn't

true in general. This second way of obtaining a two-sided p -value seems a bit odd at first blush, and its slight advantage over the first method isn't worth the confusion it tends to instill in students in an introductory course such as this. $\diamond$

4.5 On 'rejecting' null hypotheses

Researchers will often say that they 'reject' the null hypothesis in favour of the alternative hypothesis if $p \leq \alpha$, where typically $\alpha = 0.05$. This practice is subject to often heated debate (see [McShane et al., 2019](#)), and it's important to realise that $p \leq \alpha$ is possible even if the null hypothesis is true. In theory, significance tests guarantee that $p \leq \alpha$ in at most $100\alpha\%$ of the studies in which H_0 actually is true (for all $\alpha \in (0, 1)$). This is true in particular for p -values computed using exhaustive rerandomisation under the strong null hypothesis.

Theorem 4.7. Under the strong null hypothesis, exhaustive rerandomisation results in a p -value P such that $\mathbb{P}(P \leq \alpha) \leq \alpha$ for each $\alpha \in (0, 1)$. $\diamond$

Proof. Assume there are M possible randomisations. Under the strong null hypothesis, we can compute the test statistic (e.g., a signed or absolute mean difference in our example) for each rerandomisation. Sort these test statistics in descending order, breaking ties randomly if necessary:

$$t_1 \geq t_2 \geq \cdots \geq t_{M-1} \geq t_M.$$

After this sorting, the p -value that the i -th randomisation would have resulted in (call it p_i) is at least i/M , $i = 1, \dots, M$: we have $t_1 \geq t_2 \geq \cdots \geq t_i$ for each i . (It is possible that $p_i > i/M$ if $t_{i+1} = t_i$.)

For any $\alpha \in (0, 1)$, now compute $\kappa := \lfloor \alpha M \rfloor$. ($\lfloor \cdot \rfloor$ rounds down the number to the nearest integer.) We were equally likely to have generated each of the M possible randomisations. Hence, the probability that we generated a randomisation resulting in a p -value P no larger than α is

$$\begin{aligned} \mathbb{P}(P \leq \alpha) &= \frac{\# \text{ randomisations with } p_i \leq \alpha}{M} \\ &\leq \frac{\# \text{ randomisations with } i/M \leq \alpha}{M} \\ &= \frac{\# \text{ rerandomisations with } i \leq \kappa}{M} \\ &= \frac{\kappa}{M} \\ &\leq \alpha. \end{aligned}$$

It is possible to generalise the procedure for computing p -values to cases in which the randomisations aren't all equally likely to occur, and the proof of this theorem can be adapted to take this into account. I see little value in considering such cases in the present course, however.

$\square$

Remark 4.8. Nothing in the proof of Theorem 4.7 depends on the choice of test statistic. In the running example, we used the mean

difference between the conditions as the test statistic. But the difference between the condition median or the probability of superiority (see Definition A.2 on page 93), or indeed any other statistic, can be used instead to obtain a valid p -value. But what you shouldn't do is try out several approaches and then picking the one that 'works best'!

◇

In principle, then, p -values can be used to bound the probability that we claim that something is going on in the data if in fact nothing is going on: We pick some value for α (typically 0.05). If H_0 is in fact true, then we'd only observe p -values lower than α with a probability of at most α . In practice, however, things aren't so simple. We'll return to this when discussing questionable research practices; but already see Exercise 4.11 and Example 4.12.

Furthermore, $p > \alpha$ can occur even if the alternative hypothesis is true. Consequently, researchers who are in the business of 'rejecting' null hypotheses can make two types of errors, depending on whether H_0 or H_A is actually true; see Table 4.8. Without additional information (e.g., in the form of converging evidence from other studies or logical reasoning), we can't really know whether ' $p \leq 0.05$ ' represents an error or a true finding. (!)

	H_0 is actually correct	H_A is actually correct
$p > \alpha$	Fine—we didn't reject H_0	Wrong conclusion
$p \leq \alpha$	Wrong conclusion	Fine—we rejected H_0

Furthermore, that the H_A stipulates that the results are due to a combination of chance and systematic factors. It doesn't stipulate *which* systematic factors, though. What we would like to conclude is that the systematic factor at play is our experimental manipulation. But expectancy effects, failed manipulations, confounding and collider bias are also systematic factors.

What is more, the experimental manipulation may exert a systematic effect on the results, but for different reasons from what we think it does. For instance, a systematic difference between the *with* and *without alcohol* conditions needn't be due to alcohol intake per se but may be related to the taste of the beers in question instead. Or maybe alcohol increases speech rate—not because the participants become more fluent per se, but because they use simpler syntactic constructions that they can produce more quickly. In other studies, different theoretical explanations may account for any given finding—in addition to more mundane reasons such as confounding, expectancy effects and the like.

Exercise 4.9. Assume that 10,000 experiments are carried out and analysed appropriately. Further assume that the null hypothesis is correct in all of these experiments.

- (a) Which of the four histograms shown in Figure 4.9 on the facing page could plausibly represent the distribution of the 10,000 p -values?

Table 4.3: If you're in the business of rejecting null hypotheses, there are two types of errors you can make. Incorrectly rejecting the H_0 is commonly referred to as a **Type-I error**; incorrectly not rejecting the H_0 is referred to as a **Type-II error**.

- (b) Which of the four histograms would you hope to find if the alternative hypothesis were correct in all of the 10,000 experiments? Why? ◇




Figure 4.4: What could the distribution of 10,000 p -values look like under the null hypothesis?

Exercise 4.10 (False discovery rate). Imagine that in a given academic field, 1,000 experiments are conducted over the course of a year. Assume that in 80% of these experiments, the null hypothesis is correct, whereas in 20%, the alternative hypothesis is correct. Further assume that if the null hypothesis is correct, there is a 5% chance that a statistically significant difference will be observed between the experiment's conditions; if the alternative hypothesis is correct, there is a 60% chance of observing a statistically significant difference.

For the following questions, you may use the following probabilistic fact: If n attempts each succeed with probability p , then the expected number of successful attempts out of the n attempts is np .

- In how many of the 1,000 experiments can we expect that the null hypothesis will be incorrectly rejected?
- In how many of the 1,000 experiments can we expect that the null hypothesis will be correctly rejected?
- Now consider the total number of experiments in which we expect to reject the null hypothesis. In what percentage of these will the null hypothesis actually be correct? ◇

Exercise 4.11 (Multiple testing). A research team is conducting an experiment in 30 school classes simultaneously. In each school class, the children are assigned to the control or experimental group using complete randomisation. That is, the researchers are in effect conducting 30 parallel and independent experiments.

Assume that in all of these classes, the null hypothesis is true. Further assume that, if the null hypothesis is true, there is exactly a 5% chance of observing a statistically significant difference between the control and experimental conditions.

What is the probability of obtaining a statistically significant difference in at least one of the 30 experiments? ◇

Example 4.12 ('Flexible' testing). Theorem 4.7 assumes that the

analyst decided on a suitable test statistic *before* looking at the data. To appreciate this, assume that the strong null hypothesis holds in our running example, that is, that the participants would have had the speech rates listed in Table 4.3.

If the analyst decided beforehand to test if moderate alcohol intake increases the speech rates and correspondingly used the mean difference in favour of the *with alcohol* group as the test statistic, then she would have obtained a p -value no larger than $\alpha = 0.05$ with 4.76% probability.

Similarly, if she had decided beforehand to test if moderate alcohol intake decreases the speech rates and had correspondingly used the mean difference in favour of the *without alcohol* condition as the test statistic, she would have obtained a p -value no larger than $\alpha = 0.05$ with 3.97% probability. In both cases, $\mathbb{P}(P \leq 0.05) \leq 0.05$ as guaranteed by Theorem 4.7.

However, imagine that the analyst had first looked at the data to see which condition had the faster average speech rate before deciding whether to test whether moderate alcohol intake increased or decreased verbal fluency. In this case, the probability of obtaining a p -value no larger than $\alpha = 0.05$ would have been 8.73%—larger than the 5% upper bound guaranteed by Theorem 4.7.

As we will see in Chapter 10, undisclosed flexibility when running significance test can render their results essentially meaningless. ◇

4.6 Monte Carlo rerandomisation p -values

Exhaustive rerandomisation is computationally impossible for larger samples and may be infeasible for more complex research designs. One alternative is to use a Monte Carlo simulation: Instead of generating all possible rerandomisations, we only generate, say, 20,000 of them, including the randomisation we actually observed. Then, we compute the proportion of the rerandomisations considered that resulted in test statistic values at least as extreme as the one we actually observed; this proportion serves as the Monte Carlo p -value. The guarantee encapsulated in Theorem 4.7 also holds for such Monte Carlo p -values. However, if you run the same analysis multiple times, the resulting p -value is bound to vary somewhat from run to run due to the inherent randomness in the simulation.

How many ways are there to split up 40 participants into two equal-sized groups? Assuming it takes 3 milliseconds to evaluate one rerandomisation, how many *years* would it take to run the test when using exhaustive rerandomisation?

4.7 Analytical shortcuts

Statistical inference predates computers, and analytical shortcuts were developed to circumvent the tedious computations required for randomisation tests. These shortcuts include the t -test, the χ^2 -test, and ANOVA, as well as their generalisations. They typically produce similar results to those produced by randomisation tests and are usually used instead. In the context of experiments with

random assignment, the p -values that these procedures return have the essentially same interpretation and are subject to the same caveats as those above. That said, they are based on a different set of assumptions from randomisation tests. Typically, the guarantee from Theorem 4.7 only holds approximately when using these shortcuts (i.e., $P(p \leq \alpha) \approx \alpha$ under H_0).

4.8 Statistical power

A study's statistical power is the probability with which its significance test will yield $p \leq \alpha$ under the alternative hypothesis, for some predetermined $\alpha \in (0, 1)$. In studies in which one group is compared to a different group, this probability depends on three factors apart from α itself (see Figure 4.8):

1. The size of the difference in the outcome between the groups that the systematic factors cause. Even if they don't cause any difference, it is possible to obtain a statistically significant difference due to chance.
2. The number of observations.
3. The variability in the outcome variable within each group.




Figure 4.5: These three graphs show how the statistical power of a study varies with the effect size (left), the number of observations per group (middle) and the variability in the outcome variable within each group (right).

The precise numbers along the y -axis in Figure 4.8 aren't important; what's relevant is the direction and the shape of the curves.

Exercise 4.13. Take a look at Figure 4.8 and answer the following questions:

- (a) How do the effect size, the number of observations and the within-group variability in the outcome affect the probability that a study will yield a statistically significant result?
- (b) Other things equal, what yields a greater improvement in a study's power: 10 additional participants per group when each group already consists of 10 participants, or 20 additional participants per group, when each group already consists of 50 participants?
- (c) How could researchers reduce the within-group variability in the outcome variable in order to increase their statistical power?



Exercise 4.14. In an experiment that was carried out and analysed

rigorously, we find that the mean difference between the control and intervention groups amounts to 5 points on a 100-point scale. This difference is “statistically significant”, with a p -value of 0.02.

Decide for each claim below whether it follows logically from the description above. Briefly explain your reasoning.

- (a) It’s unlikely that we would have found a difference of 5 points or larger between both groups if the null hypothesis were indeed true. More precisely, this probability would have only been 2%.
- (b) The null hypothesis is incorrect; the alternative hypothesis is correct.
- (c) It’s unlikely that the null hypothesis is indeed correct. More precisely, the probability that it is correct is only 2%.
- (d) It’s highly likely that the alternative hypothesis is correct. More precisely, the probability that it is correct is 98%.
- (e) A new but similar study would likely yield a low p -value as well. More precisely, there is a 98% probability that such a study would yield a significant p -value (i.e., $p \leq 0.05$).
- (f) If we concluded that the alternative hypothesis is correct, we would be wrong at most 2% of the time.
- (g) If we concluded that the alternative hypothesis is correct, we would be wrong at most 5% of the time. ◇

4.9 *Examples in R

See the file `randomisation_tests.html` in the tutorials directory on <https://github.com/janhove/QuantitativeMethodology>.

4.10 *Further reading

The blog post *The population model and the randomisation model of statistical inference* discusses why the randomisation-based approach to statistical inference used in this chapter differs from the population-based approach that is more commonly taught in introductory statistics courses. The blog entries *Explaining key concepts using permutation tests* and *A purely graphical explanation of p-values* may be of some further use.

Goodman (2008) discusses some common misinterpretations of p -values; his list is far from exhaustive.

Analysing quantitative data and running significance tests aren’t synonymous; see my booklet as well as Winter (2019) for introductions to statistics for linguists that don’t emphasise significance testing. Nonetheless, quantitative research in the social sciences, including in applied linguistics, has developed something of a significance fetish, with authors inundating their readers with significance tests and p -values that they themselves don’t seem to really

understand while giving them little insight into what the data actually look like. For further lamentations and some suggestions, see [Vanhove \(2021\)](#).

5

Increasing precision

5.1 Precision

Up till now, our chief goal has been to increase the study's internal validity:

- Bias introduced by confounding can be countered by randomly assigning the participants (or whatever units are being investigated) to the conditions. This won't always be possible, but randomisation remains the ideal.
- Bias introduced by expectancy effects, especially on the part of the researchers, can be reduced by blinding—for instance, by preventing raters from knowing which experimental condition the participant was assigned to.
- To decrease the chances that the results are affected by technical glitches or misunderstandings, the experiment should be piloted, and checks for comprehension and satisficing can be incorporated.

These three points concern bias—we want to prevent our study from systematically under- or overestimating the answer to the question we're interested in, which is often a causal question. But as we saw when discussing statistical tests, there is a random element to the results of any given study (also see Remark 2.12 on page 26). In a study with random assignment, the luck of the draw may produce an estimated effect that is larger or smaller than the actual effect—it's just that randomisation helps to prevent this estimate from being *systematically* too large or too small. Roughly speaking, randomisation is equally likely to yield overestimates as it is to yield underestimates.

But *an estimate obtained from an unbiased study can be completely off-target*. We've demonstrated this in Remark 2.12, but to better appreciate this, consider a six-sided dice. The average number of pips on a six-sided dice is $\frac{1+2+3+4+5+6}{6} = 3.5$. Let's pretend we didn't know this and we wanted to estimate this number (i.e., 3.5) by throwing the dice and jotting down the number of pips showing face-up.

- If we do this just once, we'll just obtain an integer between 1 and 6 with equal probability (six possibilities). If we obtain a 1, our estimate will be off by 2.5 pips; if we obtain a 2, we'll be off by 1.5 pips; ...; if we obtain a 6, we'll be off by 2.5 pips. Taking into account all six possibilities, our average absolute estimation error is 1.5 pips. Note that the estimation procedure itself is unbiased: underestimates and overestimates are equally likely to occur, and they're of the same size, so they will cancel each other out.
- If we throw the dice twice, we'll now observe one of $6^2 = 36$ possible outcomes. When we average the number of pips on both throws, we can still obtain an estimate of 1 (when you throw two 1s), but there's just a 1-in-36 probability of that happening. Six of the possible outcomes will be right on the mark (1+6, 6+1, 2+5, 5+2, 3+4, 4+3). Taking into account all 36 possibilities, our average absolute estimation error is 0.97 pips. Again, this estimation procedure is unbiased.
- If we throw the dice five times, we'll observe one of $6^5 = 7776$ possible outcomes. When we average the number of pips on the five throws, there's just a 1-in-7,776 probability that we'll end up estimating the average number of pips on the dice as 1. Taking into account all 7,776 possibilities, our average absolute estimation error is 0.62 pips. Again, this estimation procedure is unbiased.

So as we increase the number of throws (i.e., as we increase the sample size), the average of the observations tends to correspond more closely to the true average. Put differently, our estimate tends to become more **precise**. It's still *possible* to be completely off mark, but it's less *probable*. Clearly, the third 'design' (throwing the dice 5 times) is preferable to the first and second design—not because it's unbiased (all three attempts are unbiased), but because the estimate it yields is expected to be closer to the truth.

In a similar vein, even unbiased studies can often be improved upon by taking steps that increase their precision. The precision of an estimate obtained in a study can itself be estimated and is typically expressed by means of **standard errors**, **confidence intervals**, or **credible intervals**. We won't concern ourselves here with how these statistics are to be calculated and interpreted; a rough appreciation of precision along the lines of the dice example suffices.

5.2 Factors affecting precision

Precision is affected mainly by the following two factors:

- the number of data points. Other things equal, larger studies yield more precise estimates. As the dice example illustrates, the effect of increasing the sample size yields diminishing returns: the same number of added observations results in a greater increase in precision if the original sample is small compared to




Figure 5.1: If we throw a six-sided die once, we'll observe one of these 6 outcomes. The dashed vertical line highlights the true mean number of pips.




Figure 5.2: If we throw a six-sided die twice and take the mean number of pips observed, we'll obtain one of these 36 outcomes.




Figure 5.3: If we throw a six-sided die five times and take the mean number of pips observed, we'll obtain one of these 7,776 outcomes.

The phrase *other things equal* is crucial. There is often a trade-off between the sample size and the quality of the data.

when it is large.

- the variability of the data within each group. The more variable the data within the groups are, the less precise the estimates will be. (In the dice example, the absolute estimation error would be lower if our dice didn't have the values 1 and 6.) For the exercises in Section 4.8, you may have already identified a couple of ways to reduce the variability of the data within the groups (e.g., restricting the study to a more homogeneous group; using more precise measurements). But we can also reduce this variability through a combination of experimental design and statistics, see Sections 5.3 and 5.4.

Both of these factors also affect statistical power in the same way (see Figure 4.8 on page 47). Indeed, at least in the situations we'll be dealing with, design choices that increase precision also tend to increase statistical power.

5.3 Matching and blocking

Definition 5.1. Matching is a procedure in which researchers manually assign the participants (or whatever units are being investigated) to the different conditions in such a way that both conditions are comparable on one or a number of background variables. ◇

In the strict sense, matching seeks to equate exactly or approximately the distributions of such background variables between the conditions. For instance, for each participant in condition *A*, you try to find a similar participant (e.g., same age, sex, and L2 level) and assign this participant to condition *B*. This way, each participant has a counterpart in the other condition.

Many applied linguists, however, apply matching in a looser sense: They assign participants to the conditions *A*, *B* in such a way that the *average* age (etc.) is similar or identical in the conditions. But the individual participants themselves don't need to have any particular counterpart in the other group, nor does the shape of the age (etc.) distribution have to be the same in both conditions.

The rationale behind matching is that, by equating the conditions on one or a number of background variables, these variables can't act as confounding variables. However, matching is **not recommended** as a substitute for randomisation: It's possible that the researchers, while matching the participants on one background variable, inadvertently introduce a bias with respect to another background variable (see Chapter 8). Moreover, matching only allows you to equate those confounding variables that you matched for (see Figure 5.4). Randomisation also probabilistically equates *other* (and indeed unknown) confounding variables and is superior to matching.

Definition 5.2 (Blocking).




Figure 5.4: Matching the conditions (*X*) on *A* doesn't prevent confounding by other (perhaps unobserved) variables (*U*).

“[M]atching is no real help when used to overcome initial group differences. This is not to rule out matching *as an adjunct to randomization*, as when one *gains statistical precision* by assigning students to matched pairs, and then randomly assigning one member of each pair to the experimental group, the other to the control group. In the statistical literature this is known as ‘**blocking**.’” (Campbell & Stanley, 1963, p. 15; my emphasis)

◇

Example 5.3 (Blocking using pairs). Based on a pool of participants, we build pairs of participants of the same sex and age and with a similar IQ. From each pair of participants, we *randomly* (rather than arbitrarily) assign one participant to condition *A* and one to condition *B*. In doing so, we both equate the two groups in terms of age, sex and IQ, but, due to the randomised assignment, we also prevent confounding by unobserved factors.

This technique is rarely used in our line of research, possibly because the pool of participants is rarely known at the start of the experiment. However, you may want to consider using it when doing research in classes, where you presumably know the roster of pupils beforehand.

◇

Example 5.4. Ludke et al. (2014) randomly assigned one third of their female participants to each of the three conditions, and similarly for the male participants. The random assignment helps to prevent confounding variables from biasing the results, the added benefit is that the three conditions have the same sex distribution.

◇

That said, it is not clear to me why they chose to block on sex.

Blocking can increase a study’s statistical precision—provided it is taken into account during the analysis. The stronger the outcome is related to the blocking factors, the more powerful blocking is.

Furthermore, blocking takes place *before* the random assignment. You can’t block after the fact.

Remark 5.5 (Blocking and randomisation testing). If you incorporate blocking in the design of your study, but you don’t take this into account when analysing the data, you’re not reaping its full benefits.

If you’re testing some association using rerandomisation as discussed in the previous chapter, the idea is that you only generate randomisations that you could actually have obtained in your study. For instance, if you block your participants into k pairs based on sex, age, and IQ as in the first example above, and then randomly allocate the participants to the conditions within each pair, there are 2^k possible randomisations instead of $\binom{2k}{k}$ ones. You should then only generate randomisations in which participants belonging to the same block are assigned to different conditions.

Similarly, in Example 5.4, only the $\left(\binom{30}{10} \cdot \binom{20}{10}\right)^2 \approx 3.1 \cdot 10^{25}$ randomisations that split the participants of each sex equally among

the conditions should be considered rather than the $\binom{60}{20} \cdot \binom{40}{20} \approx 5.8 \cdot 10^{26}$ possibilities that unrestricted randomisation could have generated.

Further see [Vanhove \(2015\)](#) for some other options that don't involve randomisation testing. ◇

How does blocking increase precision? Let's say you're comparing the efficacy of two methods for learning Dutch as a foreign language. Six German-speaking and eight French-speaking learners sign up for your study, they work with one of the two learning methods for a while, and then take the same test at the end.

- The speakers of German can be expected to have an advantage because of the similarity between Dutch and German. Let's say this advantage corresponds to 3 points on a 20-point test scale. (Realistically, you wouldn't be able to peg this number down so precisely.)
- Within each language group, learners still vary in their ability to learn Dutch.
- Let's say that, unbeknownst to you, learning method *B* boosts test performance by two points on average relative to learning method *A*. Figure 5.5 shows what the learners' scores would have been if they each had been tested on both methods, i.e., their potential outcomes.

Again, the number of participants in this fictitious example is low to keep things tractable.





Figure 5.5: The test scores that each learner would obtain for each learning method. But we can't test all learners on one and the same method, so for half of the learners, we'll observe their scores for Method A, and for the other half, we'll observe their scores on Method B.

There are $\binom{14}{7} = 3432$ ways to randomly assign these 14 participants to two conditions (*A* and *B*) with 7 participants each. For the learners in Condition *A*, we'd observe the scores shown as circles in Figure 5.5; for the learners in Condition *B*, we'd observe the scores shown as crosses. If we then took the mean difference between these groups, we'd end up observing one of the 3,432 values shown in Figure 5.5 on the following page. Complete random assignment would yield a mean absolute estimation error of about 0.68 points.

However, we've identified a likely important source of variabil-

ity in the data: language group. It makes sense to block on this factor, i.e., restrict the random assignment in such a way that half of the learners in each language group are assigned to one learning method and the other half to the other learning method. Of the 3,432 total possible random allocations, only 1,400 feature four French speakers in one condition and four in the other, as well as three German speakers in one condition and three in the other. The average estimated difference between the two learning methods among these 1,400 allocations is still 2, but the average absolute estimation error is now only 0.59—a reduction of the average absolute error of about 13%; see Figure 5.5. What has happened is that, by restricting the randomisation in this fashion, we limited both the number of allocations that would have yielded large overestimates (when the Method B condition would've consisted mainly of advantaged German speakers) and those that would have yielded large underestimates (when the Method B condition would've consisted mainly of disadvantaged French speakers). In our example, complete randomisation resulted in 244 out of 3,432 (7%) allocations with an absolute estimation error of more than 1.5 points; blocked randomisation resulted in only 46 out of 1,400 (3%) such allocations. Also see Table 5.5.



There are $\binom{8}{4} = 70$ ways to split up the eight French speakers into two groups of four, and $\binom{6}{3} = 20$ ways of splitting up the German speakers into two groups of three. Combining these yields $70 \cdot 20 = 1400$ possibilities.

Figure 5.6: Across all 3,432 possible random assignments, the mean absolute estimation error of the difference between the two learning methods is about 0.67 points. The vertical line highlights the average estimate across all 3,432 possible random assignments.

Interval	Complete randomisation	Blocked randomisation
$(-0.5, 0.5]$	3.55	1.64
$(0.5, 1.5]$	24.94	24.00
$(1.5, 2.5]$	43.01	48.71
$(2.5, 3.5]$	24.94	24.00
$(3.5, 4.5]$	3.55	1.64

So blocking on influential factors prevents the randomisation from generating some of the ‘unlucky’ allocations, thereby reducing the study’s average absolute estimation error, i.e., increasing its

Table 5.1: In what percentage of randomisations did the mean difference end up in each interval? For blocked randomisation, a greater percentage of randomisations yield a mean difference close to the true difference between the learning methods (i.e., 2) than for complete randomisation.




Figure 5.7: Of the 3,432 possible random assignments, only 1,400 have an equal number of French speakers assigned to each learning method, as well as an equal number of German speakers assigned to each learning method. Across these 1,400 assignments, the mean absolute estimation error is just 0.59. The average estimate, highlighted by the vertical line, doesn't change.

precision.

The higher precision obtained by blocking is reflected in an increase in statistical power. In this particular example, the directional randomisation test presented from the previous chapter will return a p -value of 0.05 or less in 2,096 out of the 3,432 possible allocations when using complete randomisation, that is, it has 61% power. But when using blocking, such a significance test will return a p -value of 0.05 or less in 1,353 out of the 1,400 possible allocation, that is, it has 97% power.

5.4 Leveraging control variables

In randomised experiments, the added value of control variables (see Definition 3.1 on page 31) is mostly of a statistical nature: If control variables can account for differences in the outcome between participants, they can be used to statistically reduce the variability within the groups. Similarly to blocking, this yields greater power and precision. The use of a control variable to this end must be planned before the data are collected; you shouldn't try out a bunch of 'potential' control variables during the analysis to see which one works best!

Often, the most potent indicator of a participant's performance at the end of the experiment is their performance at the start of the experiment. A pre-intervention measure of their performance—be it in the form of a score on a full-fledged **pretest** or some rougher proxy—is therefore a useful control variable.

It's also possible to 'block' on pretest scores. To this end, sort the participants according to their pretest score and divide them up into pairs like so: (12)(34)(56)(78)... Within each pair, randomly assign one participant to the control group and one to the intervention group. You can similarly block on other continuous variables.

Contrary to common belief, including powerful control variables in the analysis is useful *even if* the groups are balanced with respect to these control variables. In fact, they're even more useful than when the groups aren't balanced.

On taking into account pretest results, see Vanhove (2015).

Pre- and post-tests don't have to look identically. Any measure of pre-experiment performance is better than no measure at all.

Of course, if the pre- and posttests aren't similar and can't be scored on the same scale, you won't be able to make any claims about how much the participants progressed in each condition. *But that's not important!* What's important in a pretest/posttest design is the comparison between the conditions on the posttest scores. The purpose of the pretest scores is to increase the precision of this comparison. This can be achieved by using them as you would any other blocking or control variable, so they don't have to be expressed on the same scale as or be otherwise comparable with the posttest.

In fairly small studies, blocking tends to increase precision a bit more than merely using control variables in the analysis, but in the vast majority of cases, either is a good idea compared to the alternative of not leveraging any prior information!

Even if you're conducting a randomised experiment, it pays to think about which factors are likely to strongly affect the outcome so that, if feasible, you can take these factors into account using blocking or by means of control variables.

We don't need to concern ourselves with the freak cases where blocking or using control variables reduces precision (namely tiny studies in which the blocking or control variables are uninformative with respect to the outcome; Imai et al., 2008.)

Don't go overboard with this, though. One or two strong blocking or control variables are likely to be helpful; umpteen variables that *might conceivably* bear some relation to the outcome aren't. Using several highly intercorrelated control variables doesn't hurt but it isn't too useful either: they will all tend to do the same work, which makes them mutually superfluous.

Exercise 5.6. Having taken this class, you've become the go-to expert on experimental methodology among your friends. So unavoidably, a friend of yours solicits your input on an experiment she wants to run with one experimental condition and one control condition in a Zurich-based school class with 24 pupils. For each pupil, your friend knows if they have Swiss-German as their main language and whether they receive special educational measures. She also knows the pupils' average school grades. These data are shown in Table 5.6 on the next page.

After some discussion with your friend, you realise that, ideally, she should block on language background, whether the pupils receive special educational measures as well as average grade when randomly allocating the pupils to the experiment's condition. An equal number of pupils is to be assigned to both conditions.

- (a) Explain, as you would to your friend, how to generate a suitable allocation of pupils to conditions. Also generate such a suitable allocation and comment the steps involved. List the

Pupil ID	Main language	Special measures?	Average grade
S01	Swiss-German	no	4.9
S02	Swiss-German	no	5.4
S03	Swiss-German	yes	4.2
S04	other	yes	3.5
S05	Swiss-German	no	4.0
S06	Swiss-German	no	4.6
S07	Swiss-German	no	6.0
S08	Swiss-German	no	3.9
S09	Swiss-German	no	4.7
S10	other	yes	5.0
S11	Swiss-German	yes	5.1
S12	other	no	4.2
S13	other	yes	3.9
S14	Swiss-German	no	4.5
S15	other	yes	4.2
S16	Swiss-German	yes	4.5
S17	other	yes	3.0
S18	other	no	4.2
S19	other	no	5.3
S20	other	yes	2.7
S21	other	no	4.6
S22	Swiss-German	no	5.3
S23	Swiss-German	yes	4.9
S24	Swiss-German	no	4.7


Table 5.2: Fictional pupil characteristics.

pupil IDs of the pupils that would be assigned to the control condition under the allocation you generated. Feel free to use your spreadsheet software or R for this part.

- (b) How many different allocations could your scheme generate? How many different allocations could complete randomisation without blocking have generated? Provide both the computations as well as the numeric results. ◇

Remark 5.7 (Don't control for post-treatment variables!). A fairly common error researchers make is controlling for variables that are themselves (directly or indirectly) affected by the treatment. The reason is that controlling for a descendant of a variable (when drawn in a DAG) is like controlling for the variable itself, only less strongly.

- If you 'controlled' for the treatment variable (e.g., throwing away data in order to keep it constant), you wouldn't be able to compare the outcome variable according to different values of the treatment variable (since there aren't any). Controlling for a descendant of the treatment variable (even by statistical means rather than by selecting observations) similarly amounts to throwing away data, just to a lesser extent. Rather than increase power and precision, you'll lose some.



- If you ‘controlled’ for the outcome variable, you wouldn’t be able to find any differences between the treatment groups even if the treatment produced some differences (since you fixed all outcome observations to the same value). Similarly, controlling for a descendant of the outcome variable (even by statistical means) typically amounts to artificially pulling the differences between the treatment groups towards zero.

See the DAGs in Figure 5.8 for these and two other cases. ◇

Example 5.8. Say you want to find out if a pedagogical intervention boosts learners’ conversational French skills. It may be a good idea to control for the learners’ vocabulary knowledge. But if you collect the measure of vocabulary knowledge *after* the intervention, it’s possible that this measure is also affected by the intervention. If you control for it, you could find yourself in one of the situations depicted in Figure 5.8: learners could conceivably pick up some vocabulary as they’re working on their conversational skills. ◇

A poorly chosen pretreatment control variable won’t be too helpful, but it won’t hurt your study either—other than in terms of time spent and money wasted. But controlling for a posttreatment variable can bias your results or decrease their precision. Luckily, in true experiments, there’s a simple solution:

Collect control variables at the outset of the study (before the intervention) so that you’re sure that the control variables aren’t themselves influenced by the intervention.

Exercise 5.9. Consider the DAG in Figure 5.9.

- If the goal is to obtain an unbiased estimate of the causal influence of X on Y , regardless of precision, which variable or which variables do you *need* to control for?
- Assume that all variables shown in the DAG have already been collected (i.e., there are no further costs involved in collecting them). If the goal is to obtain an unbiased and maximally precise estimate of the causal influence of X on Y , which variable or which variables would you *need* to control for? ◇

Figure 5.8: In all four DAGs, A is causally affected by X_r .

Top left: A is a collider. Controlling for it opens up a non-causal path between X_r and Y .

Bottom left: Controlling for A closes a causal path from X_r to Y . You’d be answering the question *What’s the causal effect of X_r on Y other than that mediated by A ?* This may be interesting in its own right, but you have to be aware that this question differs from the question *What’s the causal effect of X_r on Y ?*

Top right: Controlling for A is like controlling for Y , just less strongly. If you want to estimate the causal effect of X_r on Y , controlling for Y would be a dreadful idea, so controlling for A would only be a slightly less dreadful idea.

Bottom right: Controlling for A isn’t as terrible here as it is in the top right panel. But it’s unnecessary, and you’ll lose some precision.




Figure 5.9: DAG for Exercise 5.9.

5.5 **Examples in R*

See the file `blocking.html` in the tutorials directory on <https://github.com/janhove/QuantitativeMethodology>.

6

Pedagogical interventions

The following remarks are especially relevant to pedagogical interventions, but they apply to other studies, too.

6.1 *Sample mortality*

Sample mortality refers to research participants who become unavailable in the course of a study. In a school-based study, examples include children who can't participate in a follow-up test because of a dental appointment, because they switched schools, or quite simply, because they don't want to. Less lurid and more descriptive terms for sample mortality are **drop-outs**, **outmovers**, and **panel attrition**.

In addition to lowering the sample size, drop-outs may bias the results of the study. As Figure 6.1 shows, being a drop-out or not can be a post-treatment factor (see Remark 5.7), but one that has insidiously already been controlled for: the participants whose data goes into the analysis all have the same value on this variable.




Figure 6.1: The fact that some participants stayed in the study and others can be treated as a factor in its own right (D_s ; D for 'drop-out', s for making clear that selection took place).

Ideally, mortality doesn't depend on the condition (X), the participants' prior knowledge or other characteristics (U), or their progress (Y). When mortality does vary by condition, prior knowledge or learning progress, you have to take into account the selection effect when interpreting the results.

Figure 6.1 illustrates how mortality can bias the study's estimates. It is in fact possible that an observed 'treatment' effect is little more than a selection effect: If only gifted or highly motivated learners remain part of the treatment condition but the control condition isn't as selective, it's hardly surprising that at the end of the study, you'll find better scores in the treatment condition than in the control condition.

Example 6.1. [Slavin et al. \(2011\)](#) conducted a pretest–posttest randomised experiment to compare the effects of two school models




Figure 6.2: Example of the biasing effect of sample mortality. *Left*: Had it been possible to test all participants, we'd have found a mean difference of 5 points. *Right*: The boxplots are now only drawn based on the participants that didn't drop out. If the drop-out likelihood is itself affected by the intervention, we could end with a biased estimate.

for bilingual children on a series of language skills. In order to investigate whether sample mortality could explain their main findings, they first compared the school models in terms of the outmover rates (i.e., which proportion of children who took the pretest didn't take the posttest) and outmover type (i.e., whether the outmovers were simply absent on the day, e.g., due to a dental appointment, or whether they had changed school); see their Table 2. Furthermore, they compared the outmovers between the school models in terms of their pretest scores; see their Table 3. ◇

6.2 Clustering

In many pedagogical settings, the participants can't be randomly assigned to the experiment's conditions on an individual basis. Instead, entire intact groups of participants (e.g., entire classes, entire schools, entire school districts etc.) are assigned to the same condition. This induces **clustering**: Due to teacher/class/school etc. effects, participants belonging to the same cluster (class, school, etc.) tend to be somewhat more alike in their performance than participants belonging to different clusters.

Definition 6.2. A **cluster-randomised experiment** is an experiment that uses random assignment of units (e.g., children) to conditions in such a way that guarantees that all units in a certain set of units (e.g., classes) are assigned to the same condition. The sets that are kept intact are called **clusters**. ◇

Figure 6.2 schematically illustrates two kinds of research design that may be encountered in pedagogical settings. In a **between-school design** (or, similarly, a between-class, between-district, etc. design), all pupils in a school are assigned to the same experimental condition. In a **within-school design** (or, similarly, in a within-class, within-district, etc. design), by contrast, the school effect is neutralised using blocking, whereas the remaining possible confounding variables are ideally taken care of using randomisation. This increases precision relative to a between-school design.

One possible drawback of a within-school design is that the pupils in the control and treatment classes in the same school in-

fluence each other (e.g., by comparing notes or helping each other make sense of what's being taught), which may wash out an existing treatment effect.

Within-school								Between-school			
School 1		School 2		School 3		School 4		School 1		School 2	
I	C	I	C	I	C	I	C	I	C	I	C

Figure 6.3: Within- vs between-school designs. (I for intervention, C for control.)

Exercise 6.3. Let's say you want to run a pedagogical experiment (e.g., to compare two learning methods for French as a foreign language) in which randomisation has to take place at the class level rather than at the individual level. Other things equal (e.g., number of classes, number of pupils), what are the advantages and drawbacks of the following designs? What's the worst option, and why? What's the best, and why?

- All classes are taught by the same teacher.
- Each class is taught by a different teacher.
- One teacher teaches all classes in the control condition, and another teacher teaches all classes in the intervention condition.
- Each teacher teaches two classes: one in the control condition, and one in the intervention condition. ◇

Exercise 6.4. The article by [Slavin et al. \(2011\)](#) makes for pretty challenging reading, especially in terms of the analysis and the way the results are presented. But the introductory and methodological sections discuss some concepts that we've already discussed (especially p. 49).

First try to read the text in full, but skip the parts you find unintelligible. Then answer the following questions:

- What is the independent variable in this study? What are the dependent variables?
- How were the pupils assigned to the different groups?
- "Children were pretested ... on the English Peabody Picture Vocabulary Test (PPVT) and its Spanish equivalent, the Test de Vocabulario en Imagenes Peabody (TVIP)." (p. 51, right) Why did the researchers go to the bother of conducting such pretests? Try to find at least two reasons.
- Comparing Slavin et al.'s Tables 4 and 6, we observe that the cohorts' average TVIP (*Test de Vocabulario en Imagenes Peabody*) scores dropped from 99.85 and 90.19 to 92.86 and 85.64, respectively, over the course of two years.

Come up with the most plausible explanation for this result.

Hint: The children's Spanish vocabulary knowledge probably did not decline on average as they got older. ◇

While cluster-randomised experiments are commonly encountered in applied linguistics research, one crucial aspect is sadly often neglected:

In a cluster-randomised design, you need to take the cluster-randomisation into account when analysing the data.

To appreciate the need for taking clustering into account during the analysis, consider an experiment with 6 classes of 10 pupils each. There are over one-hundred *quadrillion* ways to split up 60 pupils into two groups of 30 ($\binom{60}{30} \approx 1.2 \times 10^{17}$). But there are only 20 ways to split up six classes into two groups of three classes each. The analysis needs to be based on the assumption that the allocation obtained is one of 20 possible ones, not one of a gazillion ones.

Remark 6.5 (Analysis of cluster-randomised designs). One simple and valid approach for analysing data obtained with a cluster-randomised design is to compute the mean (or some other kind of average) of each cluster and then analyse these cluster-level summaries using one of the methods covered in Chapter 4. That said, lots of alternative approaches exist; see [Vanhove \(2015\)](#), [Vanhove \(2020\)](#), and references therein. ◇

What you also need to know about cluster-randomisation is this:

Studies with one intact group as the experimental group and another intact group as the control group are useless.

Conceptually, the reason is that class, school and teacher effects can't be separated from the effect of the intervention if you just have one intact group per condition. If you use a randomisation test as per Remark 6.5 but you only have a single cluster per group, then there are only two possible allocations. As a result, the left- and right-sided p -value will both be at least 0.50, and the two-sided p -value will always be 1.

Remark 6.6. More complex research designs may combine aspects of between- and within-school (or -class, or -whatever) designs. For instance, researchers interested in L2 vocabulary learning could compare different methods for designing flash cards (i.e., cards with L2 words on one side and their L1 translations or some description on the other side). They may want to compare four methods:

- the learners design the physical cards themselves;
- the learners are given physical cards prepared by the researchers;
- the learners design digital cards themselves in a computer pro-

gram or smartphone app;

- the learners are given a digital card deck prepared by the researchers.

The researchers may then recruit multiple classes of learners to participate in the experiment.

For logistical reasons, it may be preferable to have all learners in the same class use either physical or digital cards. But within any given class, some learners can still design their own cards whereas others work with prepared cards. The resulting design would have two factors: (1) physical vs digital, and (2) prepared vs self-made. The first factor would be a between-class factor; the second factor a within-class factor.

Incidentally, the terms *between-class* (or between-whatever) and *within-class* (or within-whatever) refer to whether the comparison for a given factor can only be made between different classes or can also be made within the same class. ◇

6.3 *Examples in R

See the file `clusters.html` in the tutorials directory on <https://github.com/janhove/QuantitativeMethodology>.

7

Within-subjects experiments

Blocking increases power and precision by pairing up similar participants and randomly assigning one of each pair to each condition. In within-subjects designs, this idea is taken to an extreme.

Definition 7.1. In a **within-subjects experiment**, each unit (e.g., participant) is assigned to multiple (typically: all) conditions. ◇

We've already encountered within-school and within-class designs; within-subjects designs extend the same logic to the individual unit.

7.1 Advantages and drawbacks

One potential advantage of within-subjects designs is that it becomes easier to explore interindividual differences in the efficacy of the treatment. However, the *main* advantage of within-subjects designs is that they usually have greater statistical precision than an otherwise comparable experiment in which the units are assigned to the conditions at random. Indeed, a study's statistical precision depends on (a) the amount of data and (b) the variability in the data. Within-subjects designs easily account for a major source of variability in the data: pre-existing differences between participants.

How much more precise a within-subjects experiment is than a between-subjects experiment varies from case to case. [Quené \(2010\)](#) estimates that within-subjects designs have the statistical precision of between-subjects designs with four times as many participants. The precise factor depends on the extent to which the participants' performance in one condition correlates with their performance in the other condition: The stronger this correlation, the greater the added value of a within-subjects experiment. But even if you can't quantify this added value: Within-subjects designs typically offer greater statistical precision.

Let's now turn to potential drawbacks of within-subjects designs. First, in applied settings, you typically want the study to mimic the context in which its findings are to be implemented. But in such a context, people (e.g., pupils) won't be exposed to several conditions

(e.g., learning methods) but rather to just one. Hence, the within-subjects design may be criticised for a lack of ecological validity.

A second potential drawback is that, when participants are tested in several conditions, it's possible that they learn something in one condition that affects their performance in the other condition. This phenomenon is called a **carry-over effect**. It's also possible that their performance in the last condition differs from their performance in the first condition because they've grown accustomed to the setting or because they've grown tired of being tested. These phenomena are referred to as **learning effects**, **fatigue effects**, and more generally, **order effects**.

7.2 Minimising order effects

The first technique to prevent learning or fatigue effects from exerting a systematic effect on the results is to vary the order of the conditions between the participants. In **complete counterbalancing**, *all* possible orders are taken into account. If you have two within-subjects conditions, half of the participants first complete condition *A* and then *B*, and the other half first complete condition *B* and then *A*. If you have three conditions, there are $3! = 6$ possible orders; you can then randomly assign one sixth of the participants to first complete *A*, then *B*, then *C*; one sixth completes *A*, then *C*, then *B*, etc.:

A	A	B	B	C	C
B	C	A	C	A	B
C	B	C	A	B	A

Table 7.1: Complete counterbalancing for a within-subjects experiment with three conditions.

If a within-subjects experiment has lots of conditions, complete counterbalancing is impractical. For four conditions, for instance, there are already $4! = 24$ possible orders—we may not even have that many participants! The second technique, the **Latin square design**, lends itself to such cases. Latin squares are arrangements of symbols in a grid in which each of the symbols used occurs exactly once in each row and exactly once in each column. The grid below is a Latin square of size 4—one of the 576 possible arrangements of the symbols *A*, *B*, *C*, and *D* that form a Latin square.

A	B	C	D
B	C	D	A
C	D	A	B
D	A	B	C

'Latin' because the symbols used are typically letters of the Latin alphabet.

Can you come up with a couple of the other 575 ones?

Table 7.2: A Latin square for a within-subjects experiment with four conditions.

Let's say you picked the Latin square above for your study. You'd then randomly assign one quarter of the participants to the condition (or stimulus set) order *ABCD* (first row), one quarter to *BCDA* (second row), one quarter to *CDAB* (third row) and one quarter to *DABC* (fourth row). The conditions (or stimulus sets) are randomly assigned to one of the letters, too.

The third technique is useful when even a Latin square design

becomes impractical. For instance, let's say that our study involves us showing 50 pictures to our participants that they are to describe. If we also want to draw comparisons between the descriptions of different pictures, then we cannot just present them in the same order to all participants: order effects could plausibly bias the results. But $50! = 3 \cdot 10^{64}$ is an astronomical number, ruling out complete counterbalancing, and even just 50 different Latin square orders seem impractical. One possible solution is to present the images in a new random order for each participant. The drawback of doing this is that perhaps image 3 occurs much more often at the start than at the end of the data collection, but at least this imbalance isn't built into the design of the study. (Of course, it's possible that we just accept such 'bias' if we aren't interested in differences between the images, but just in differences between the participants or between the conditions. If that's the case, randomising the stimulus order is superfluous.)

In many psycholinguistic studies, participants need to react to several stimuli per condition (e.g., 12 stimuli per condition). The order of the stimuli in these studies are often randomised so that the conditions are mixed up (e.g., *ABAABBBAB* etc.).

Counterbalancing and Latin squares don't negate carry-over effects. Whether carry-over effects represent an acute danger to the study's validity needs to be judged on a case-by-case basis. That said, the possible danger of carry-over effects quite often isn't large enough to offset the certain gain in statistical precision.

If you have a genuine choice between a between-subjects and a within-subjects design for your own research, pick the within-subjects design. (Unless, of course, you have an excellent reason not to do so.)

Example 7.2. In [Vanhove \(2019\)](#), I examined the influence of metalinguistic knowledge about some structure in the L1 on the participants' intuitions about the corresponding structure in a foreign language. Metalinguistic knowledge was manipulated experimentally. In the control condition (call it *A*), the participants were given correct but irrelevant information about the L1's morphosyntax. In the second condition (*B*), they were given correct, relevant information. In the third condition (*C*), they were given correct, relevant information and they were additionally taught an algorithm for identifying the structure in question.

It's quite clear that this study couldn't be realised in a within-subjects design: Once you give the participants the relevant information (conditions *B* and *C*), you can't then expect them to suppress this information as they are tested in condition *A*. Similarly, once you teach the participants an algorithm for identifying a structure (condition *C*), you can't then expect them not to apply it in the other conditions. Consequently, I opted for a between-subjects

design. ◇

Remark 7.3 (Potential outcomes in within-subjects designs). We introduced the notion of potential outcomes when discussing true (between-subjects) experiments in Chapter 2. If each unit participates in both conditions of a two-condition within-subjects experiment, it may seem as all potential outcomes are, in fact, known. But this is not the case. For a unit assigned to the AB order, the known outcomes are how they performed in the A condition *if this condition is presented first* and how they performed in the B condition *if this condition is presented second*. But unknown potential outcomes are how they would have performed in the A condition if it had been presented second and in the B condition if it had been presented first. ◇

Exercise 7.4. From Ludke et al. (2014):

Participants were randomly assigned to one of three learning conditions: speaking, rhythmic speaking, and singing. The participants heard 20 paired-associate phrases in English and an unfamiliar language (Hungarian) (...). (...) The 15-min learning period was followed by a series of five different production, recall, recognition, and vocabulary tests for the English–Hungarian pairs.

Re-design this between-subjects experiment as a within-subjects experiment. What would this description look like? For the time being, ignore the *rhythmic speaking* condition.

In your design, you are constrained by the resources Ludke et al. had. This means that you cannot introduce stimuli and tests that were not already used by Ludke et al., and that you have to work with at most 30 men and 30 women as participants. ◇

Exercise 7.5. As in the previous exercise, but with all three conditions. ◇

Exercise 7.6 (Reading assignment). The study by Kang et al. (2013) serves as an example of a within-subjects design. Additionally, it uses some turns of phrase commonly found in research reports:

- (a) “The Hebrew nouns were learned in one of two training conditions – retrieval practice or imitation – that were manipulated within subjects across separate blocks and semantic categories.” (p. 1261)
 - i. What does “manipulated within subjects across separate blocks” mean?
 - ii. What does “manipulated within subjects across semantic categories” mean?
- (b) “The order of items in each test was randomized for each learner.” (p. 1262) Why? Would it have made sense to use the same fixed order of items for all learners?
- (c) “In Experiment 2, the order of both tests was counterbalanced across learners. . . .” (p. 1262). This merely means that half of

the learners first took Test A and then Test B, whereas the other half first took Test B and then Test A. Would it have made sense to administer the test in the same order to all learners? ◇

7.3 *Analysing data from within-subjects designs with two conditions

7.3.1 AB/BA cross-over design

Consider a within-subjects experiment with two conditions, creatively labelled A and B , in which participants are randomly assigned to the two possible orders AB and BA . This design is called an **AB/BA cross-over design**. We will consider condition A to be the intervention (or treatment) condition and B the control condition, but nothing hinges on these labels. Let's break down the systematic factors that contribute to the first and second measurements in both orders, see Table 7.3.1.

Order	First measurement	Second measurement
AB	$\beta + \tau$	$\beta + \omega + \kappa$
BA	β	$\beta + \tau + \omega$

Table 7.3: Systematic factors contributing to the measurements in a within-subjects experiment with two conditions. The meaning of the Greek symbols is explained in the running text.

- We assume that there is some baseline β common to all measurements. This common baseline is of no further interest.
- There may be a differential treatment effect τ that contributes to the measurements in condition A (i.e., the first measurement for order AB and the second for order BA), but not to those in condition B .
- There may be a differential order effect ω that contributes to the second measurements but not to the first measurements.
- There may be a differential carryover effect κ that affects the measurements in condition B but only if B follows A .

In any analysis of these data, we need to assume that there are no carryover effects, that is, $\kappa = 0$: If $\kappa \neq 0$, the analysis presented here is no longer appropriate, and a within-subjects design may not have been a good choice. Under the assumption that $\kappa = 0$, we may estimate the value of τ by first computing, for each participant, the period difference, that is, the difference between their first and their second measurement. For the participants in the AB order, and ignoring any non-systematic effects, we obtain

$$d_{AB} = (\beta + \tau) - (\beta + \omega + \kappa) = \tau - \omega - \kappa = \tau - \omega,$$

since $\kappa = 0$ by assumption. For the participants in the BA order, we similarly obtain

$$d_{BA} = \beta - (\beta + \tau + \omega) = -\tau - \omega.$$

Observe that

$$\frac{d_{AB} - d_{BA}}{2} = \frac{(\tau - \omega) - (-\tau - \omega)}{2} = \tau.$$

The consequence of this is that, under the assumption of no carryover effects, the treatment effect τ can be estimated as half the mean difference in the period differences between the two orders. A randomisation test or some analytical approximation thereof can be used to obtain a p -value for the null hypothesis that $\tau = 0$.

In practice, you'll encounter different approaches to analysing within-subjects designs when reading social science studies. Not all of these are grounded in sound statistical theory.

Remark 7.7. If we aren't willing to exclude the possibility that there is a carryover effect (i.e., that $\kappa \neq 0$), we can restrict the analysis to the first measurement and treat the data as stemming from a true (between-subjects) experiment. However, by doing so, we lose half of the data as well as any any improvement in statistical precision that we sought to obtain by adopting a within-subjects design. $\diamond$

7.3.2 Interleaved conditions

Consider again a within-subjects experiment with two conditions, A and B , in which n participants are shown a fixed list of m words $w_1, \dots, w_m$. For each participant, half of the words are shown in condition A , and half are shown in condition B ; the presentation software assigns the words to the conditions randomly and separately for each participant. The participants react to each word, and their response is expressed numerically in some fashion (e.g., accuracy, speed, ...).

For a design like this, there is no real sense in which one condition follows another condition for each participant. A sensible way to analyse these data is to compute, for each participant, the difference score between their average response to condition A and their average response to condition B , resulting in difference scores $d_1, \dots, d_n$. We then compute some average of these differences, $\bar{d}$ (e.g., the mean or the median). Under the null hypothesis of no difference, each d_i value could just as likely have had the opposite sign. To generate the distribution of $\bar{d}$ under the null hypothesis, we can randomly flip the signs of the observed d_i values and recompute their average.

Exhaustive sign-flipping would generate the full distribution of $\bar{d}$ under the null hypothesis, but it involves generating 2^n $\bar{d}$ values. A Monte Carlo version of this procedure would also work. Once the distribution of $\bar{d}$ has been generated or approximated using this sign-flipping, p -values can be computed as usual. An analytical short-cut to this procedure is the paired t -test on the condition averages per participant.

Remark 7.8 (Within-school/class designs). In within-school or within-class designs (see Figure 6.2 on page 65), the same procedure can be applied: For each school (or class), compute the average score obtained by the pupils in the intervention and the average

score obtained by the pupils in the control condition. Then analyse these averages as described above. ◇

7.4 **Further reading*

Latin-square designs are also used in studies other than within-subject experiments, for instance as a technique for blocking. See [Richardson \(2018\)](#) for an overview and some finer points that weren't discussed here; his article is geared towards educational researchers.

7.5 **Examples in R*

See the file `within.html` in the tutorials directory on <https://github.com/janhove/QuantitativeMethodology>.

8

Observational studies

Up till now, we've discussed designs that eliminate pre-existing factors as confounders by means of random allocation or by testing all participants in all conditions. We now turn to between-subjects studies without random allocation: **observational studies**.

Some observational studies superficially look like experiments in that they focus on a comparison between two or more groups and may even comprise pre- and posttests, but without any random allocation of units to conditions. Such observational studies are often referred to as **quasi-experiments**. Because no random assignment was used, we can't assume that the treatment variable is independent of pre-treatment variables—confounding is a real threat. Importantly, it doesn't matter whether the groups *could* have been constructed using random assignment, just whether they were.

Why carry out observational studies?

"But just because full experimental control *is* lacking, it becomes imperative that the researcher **be thoroughly aware of which specific variables his particular design fails to control**.

"The average student or potential researcher reading the previous section of this chapter probably ends up with more things to worry about in designing an experiment that he had in mind to begin with. This is all to the good if it leads to the **design and execution of better experiments and to more circumspection in drawing inferences from the results**. It is, however, an unwanted side effect if it creates a feeling of hopelessness with regard to achieving experimental control and leads to the abandonment of such efforts in favor of even more informal methods of investigation.

"[W]e shall . . . survey the strenghts and weaknesses of a heterogeneous collection of quasi-experimental designs, each deemed worthy of use *where better designs are not feasible*." (Campbell & Stanley, 1963, p. 34; their emphasis in italics, mine in bold-face)

The goal of observational studies (including quasi-experiments) is often to draw causal conclusions, but the findings—for better or for worse—tend to be couched in non-causal language (Grosz et al., 2020). The present course cannot hope to exhaustively cover the difficulties involved in drawing causal conclusions from observa-

tional studies. Instead, the following exercise highlights a few of the problems, and the remainder of this chapter introduces and then critiques a technique commonly used in applied linguistics to draw causal conclusions from observational data. Fuller treatments can be found in [Campbell & Stanley \(1963\)](#) and in the more technical books by [Imbens & Rubin \(2015\)](#) and [Ding \(2024\)](#).

Exercise 8.1 (Controlling for confounds is difficult). Consider the following description:

“There were 40 participants who composed two language groups and two age groups. Twenty of the participants were younger adults ranging in age from 30 to 54 years (mean age = 43.0 years), and 20 were older adults ranging in age from 60 to 88 years (mean age = 71.9 years). In each age group, half the participants were monolingual English speakers living in Canada, and the other half were Tamil-English bilinguals living in India. (...) All the participants in both groups had bachelor’s degrees ...” ([Bialystok et al., 2004](#), p. 44)

- (a) Did the authors control for level of education?
- (b) It is possible that the authors introduced additional bias into their study by only including participants with a Bachelor’s degree. How so?

Hint: If you’re stuck, consult <https://gpseducation.oecd.org/CountryProfile?primaryCountry=CAN&treshold=5&topic=E0> and look up similar data for India.



8.1 Correlation coefficients

Pearson correlation coefficients, typically just called correlation coefficients and abbreviated as r , express how closely the (X,Y) data points fall on a straight line.

- $r = 1$: All points fall exactly on an increasing line.
- $r = -1$: All points fall exactly on a decreasing line. Correlation coefficients of 1 or -1 (or close to it, e.g., $r = 0.99$) tend not to be too interesting: They typically indicate that the two variables express the same thing (e.g., body length in centimetres and in inches).
- $r = 0$: There’s no linear relation between the two variables whatsoever.

Correlation coefficients work in both directions: $r_{XY} = r_{YX}$.

Figure 8.1 shows eight examples of scatterplots and the correlation coefficients for the data presented in them. Importantly, a correlation coefficient close to zero doesn’t imply that there is no relation between them; correlation coefficients different from 1 or -1 don’t imply that the relation between two variables is imperfect; and it’s possible for a positive correlation coefficient to reflect a relationship that’s largely negative, and vice versa.




Figure 8.1: Examples of scatterplots and their associated correlation coefficients.

The same correlation coefficient can correspond to a multitude of relationships between two variables. Never *ever* compute a correlation coefficient without drawing a scatterplot first.

Tip 8.2. By the same token, don't put stock in conclusions that hinge crucially on correlation coefficients for which no scatterplots are provided. ◇

Exercise 8.3 (Honing intuitions about correlation coefficients). To hone your intuitions about correlation coefficients, you can use the `plot_r()` function from the `cannonball` package for R.

```
# Install the package if necessary
# install.packages("pak")
pak::pak("janhove/cannonball")

# Load the functions
library(cannonball)

# Draw 16 plots with 20 data points each and r = 0.6
plot_r(n = 20, r = 0.6)

# With 50 data points each and r = 0.0
plot_r(n = 50, r = 0.0)

# With 40 data points and r = -0.9
plot_r(n = 40, r = -0.9)
```

Type `?plot_r` at the R prompt to access the function's help page and read the text under 'Details'. ◇

8.2 Statistical control using hierarchical regression

Exercise 8.4 (Reading assignment). The next reading assignment concerns the study by [Slevc & Miyake \(2006\)](#). The “zero-order correlations” mentioned are correlation coefficients expressing the relationship between two measured variables. (“First-order correlations” would be correlation coefficients expressing the relationship between two variables from which the influence of a third variable was statistically ‘partialled out’.)

To help you make sense of Table 3:

- R^2 : The proportion of the variation (‘variance’) in the outcome variable that can be described using the predictor variables included in the regression model.
- ΔR^2 : The increase in R^2 compared to the previous *step* (i.e., the improvement in R^2 attributable to the current predictor).
- df, F : You can ignore this for this class.

- *Final β* : Expresses the form of the relationship between the predictor in question and the outcome.

Now to the questions:

- What was the most important goal that Slevc & Miyake (2006) set themselves?
- Why did they have this aim?
- Why did they collected the variables *age of arrival*, *length of residence*, *language use and exposure* and *phonological short-term memory*?
- Which conclusions do they draw on the basis of their results?
- Regardless of whether you agree with these conclusions: Try to find one or two alternative explanations for their results that call into question the claim that “musical skills may facilitate the acquisition of L2 sound structure” (abstract). ◇

In correlational studies, control variables are often used to adjust statistically for known confounders. One technique used to accomplish this is hierarchical regression; see Table 3 in Slevc & Miyake (2006) for an example. We will discuss this technique mainly so that you are better able to appreciate the shortcomings of this technique and ones similar to it.

Example 8.5. If you measure the shoe size and vocabulary knowledge of 4- to 16-year-olds, you’ll observe a positive correlation between the two. This isn’t surprising; see Figure 8.2. We’ll use this silly example to illustrate the principle behind hierarchical regression; see Figure 8.5 on the next page.

- Top left: Shoe size and vocabulary knowledge are positively correlated.
- Top right and middle left: Age—the confound—is correlated positively with both shoe size and vocabulary knowledge.
- Middle right: This plot shows the vertical distance between the points in the middle left panel and the regression line. This shows the remaining variation in vocabulary test scores once the linear association between age and vocabulary knowledge has been partialled out.
- Bottom left: The association between shoe size and the vocabulary test scores with the linear association of age partialled out is much less strong. In this simulated example, the fact that the remaining association isn’t exactly zero is due entirely to chance. ◇




Figure 8.2: Shoe size and vocabulary knowledge are correlated since age acts as a confound.




Figure 8.3: Hierarchical regression used to control for the age confound in the relationship between shoe size and vocabulary knowledge. The coloured circles in each panel show data belonging to the same three participants. The straight lines are regression lines. A regression line is the straight line that best captures the tendency in the cloud of data points, according to some definition of 'best'.

8.3 Caveats

You need to be hyper-aware of the following caveats concerning statistical control:

1. Controlling for a number of possible confounds doesn't rule out the possibility that there are even more confounds; Figure 8.4.
2. The methods typically used to account for confounding variables account for *linear* relationships between the confounds and the variables of interest. If these relationships aren't linear, the confound won't be fully accounted for. In DAG parlance, the path via the confound won't be fully closed.
3. The 'confound' may be a post-treatment variable. See Section 5.4 on page 57.
4. Statistical control may be imperfect because the confound was measured with some error. We'll treat this in more detail in Chapter 9.

The following excerpt makes the same points:

"When experimental designs are premature, impractical, or impossible, researchers must rely on statistical methods to adjust for potentially confounding effects. Such procedures, however, are quite fallible. We examine several errors that often follow the use of statistical adjustment.

"The first is inferring a factor is causal because it predicts an outcome even after "statistical control" for other factors. This inference is fallacious when (as usual) such control involves removing the linear contribution of imperfectly measured variables, or when some confounders remain unmeasured.

"The converse fallacy is inferring a factor is not causally important because its association with the outcome is attenuated or eliminated by the inclusion of covariates in the adjustment process. This attenuation may only reflect that the covariates treated as confounders are actually mediators (intermediates) and critical to the causal chain from the study factor to the study outcome.¹

"Other problems arise due to mismeasurement of the study factor or outcome, or because these study variables are only proxies for underlying constructs.

"Statistical adjustment serves a useful function, but it cannot transform observational studies into natural experiments, and involves far more subjective judgment than many users realize." (Christenfeld et al., 2004, abstract, my emphasis)

Large sample sizes don't solve these problems.

Also see the blog entry [Controlling for confounding variables in correlational research: Four caveats](#).

Remark 8.6 (On despair). To call back to the Campbell & Stanley (1963) quote at the start of this chapter, the point of the preceding discussion is not for budding researchers to throw their hands up in despair and avoid doing or reading observational studies.




Figure 8.4: Perfectly controlling for A and B closes the non-causal paths $X \leftarrow A \rightarrow Y$ and $X \leftarrow B \rightarrow Y$. But it leaves open the non-causal path via U .

¹ What's meant is a causal chain such as $A \rightarrow B \rightarrow C$. A is causally important, but if you control for B , you won't find any association between A and C .

Rather, it is to instill in these researchers a sense of modesty as well as the appreciation that observational studies, to a much greater extent than experimental studies, leave open a host of alternative explanations for any sort of finding that they produce and that more definitive conclusions can be drawn only by considering converging evidence from a wider range of studies. Each study may leave open plausible alternative accounts of the findings. But to the extent that the studies vary in terms of settings, participants, and methods, different alternative accounts are required, and a unifying causal account becomes ever more plausible. See [Hill \(1965\)](#) for a similar sentiment in the medical context. ◇

9

Constructs and indicators

We're faced with an inescapable fact of life:

Most measurements are imperfect.

Saying that a study's measurements aren't perfect isn't much of a criticism. But it's crucial to appreciate the consequences of imperfect measures—pointing out that a study's findings can plausibly be accounted for by the fact that its measurements are imperfect *is* a valid criticism.

Definition 9.1. A **construct** or **latent variable** is a characteristic that cannot be observed or measured directly.

An **indicator** or **manifest variable** is a measurable or directly observable variable on the basis of which the existence as well as the relative values about some construct are inferred. ◇

Table 9.1 lists some examples.

Construct	Example indicator
Intelligence	Result on an intelligence test
Working memory capacity	Length of a sequence of digits you can repeat in reversed order
Language aptitude	Result on the LLAMA-D test
L2 reading skills	Number of correctly answered items on a reading test
Attitudes towards Danish	Answer to the question 'How beautiful do you think Danish is?'
Socio-economic status	Father's occupational category

Table 9.1: Examples of constructs and indicators.

Even the best indicators are rarely perfect. Better indicators just have a smaller **measurement error**. Further, even variables that don't act as proxies for some cognitive or social construct are often measured with some error. Examples include body weight (bathroom scales are imperfect, and the result is rounded), blood

pressure (if you have a sphygmomanometer, check its manual), and age (invariably rounded down to the integer below when reported).

9.1 *Systematic and random measurement error*

Measurement error can include both a systematic and a random component.

The **systematic** component of an instrument's measurement error is the extent to which it tends to over- or underestimate what it's supposed to measure. For instance, a miscalibrated kitchen scale may overestimate weights by 10 g on average, and an overly harsh language test may tend to label learners' L2 skills one CEFR level below their actual proficiency on average.

It's possible for an instrument to systematically overestimate values on one part of the scale and to underestimate them on another part.

When there's no gold standard to which the measurements can be compared, it may be impossible to assess their systematic measurement error.

The **random** component of an instrument's measurement error is the extent to which the measured values differ from the true values + systematic error. Another way of putting this is: By how much will the measurements vary if the true values are the same? For instance, a kitchen scale may, on average, measure weights accurately (no systematic error), but the individual readings may be off by up to a couple of grams in either direction (random error).

As a second example, consider a group of 365 7-year-olds, all born on different days of the year. Just one of them actually is 7 years old on the day; the reported values of the others will be off by 1 day, 2 days, ..., 364 days. The reported age, then, systematically underestimates their true age by $\frac{0+1+2+\dots+364}{365} = 182$ days. The random component is 0, though, as children born on the same day will report the same age, even though this reported age will be lower than their actual age.

As a final example, consider a poorly calibrated bathroom scale. If you put a calibrated mass of precisely 60 kg on it on five different occasions, it returns readings of 61.1, 60.4, 60.4, 60.5 and 61.2. The mean observation for the same mass is $\frac{61.1+60.4+60.4+60.5+61.2}{5} = 60.78$, i.e., an overestimate of 0.78 kg. The mean absolute difference between the observations and their expected value (here: 60.78) is $\frac{|0.32|+|-0.38|+|-0.38|+|-0.28|+|0.72|}{5} = 0.42$.

9.2 *Consequences of measurement error*

The consequence of systematic measurement error is clear: Your data are biased. This isn't necessarily a problem: If you're comparing two groups for both of which you have data that are biased

to the same extent, the difference between them won't be biased. And for variables such as age, the systematic error (roughly 182 days) tends to be small relative to the variability of the true values, in which case it's probably inconsequential. (But see, for instance, [Helsen et al. \(2005\)](#) and [Sprietsma \(2010\)](#) on the consequences of 'relative age' (i.e., age differences within an age group, e.g., 15-year-olds) in sport and education.)

The consequences of random measurement error are much less intuitive and bear pointing out.

First, measurement error on the *outcome* variable will increase its variability. Since power and precision are lower when there's more variability in the outcome, measurement error on the outcome lowers power and precision.

Second, measurement error on a *control* variable means that controlling for this observed variable won't fully eradicate the confounding caused by the construct itself. The DAG in Figure 9.1 illustrates this.

"[F]allibility in a covariate usually implies that there would be more adjustment if the variable were measured without error." ([Huiteima, 2011](#), p. 569)

Controlling for A_{obs} is better than not controlling for it. But researchers routinely mistake controlling for an indicator with controlling for a construct, and their causal conclusions are overconfident as a result. A discussion of this problem can be found in [Westfall & Yarkoni \(2016\)](#), [Vanhove & Berthele \(2017\)](#) and [Berthele & Vanhove \(2020\)](#).

Third, when observations are due partly to skill or some underlying construct and partly to chance (e.g., measurement error), a second round of observations will likely show that the extreme scores have become less extreme, i.e., that they've **regressed to the mean**.

Example 9.2. First consider an example where the observations are purely due to luck, with no skill or construct involved: playing roulette. Playing roulette is a losing proposition: For every 100 francs bet, you stand to lose about 5 francs (= the mean). But on any given night, some players will luck out and make a killing, whereas other players get extraordinarily unlucky and lose much more than the expected 5 francs. Their winnings or losses are a dreadful measure of their skill level: they all have the same skill level, which corresponds to a loss of 5 francs.

The next night, however, the lucky players from the day before probably won't get as lucky again (their luck the day before was extraordinary), and similarly for the unlucky players—all again stand to lose about 5 francs. Some might get lucky or unlucky twice in a row, but they're more likely to end up somewhere near the 5-franc mark, i.e., most of the lucky and unlucky players will regress to the mean. ◇




Figure 9.1: The X - Y relationship is confounded by A . A , however, can't be observed directly. A proxy (indicator) A_{obs} can be controlled for instead, but this won't fully shut the non-causal path $X \leftarrow A \rightarrow Y$.

Example 9.3. The same principle is at play when the observations come about in part through skill (or some other construct) and in part through chance. For instance, the most successful stock picker of the year 2032 is likely not to perform as well in the year 2033—even if the conditions on the stock market are comparable and the stock picker didn't start to rest on his laurels. The reason could simply be that he had more than his fair share of luck in 2032—you need some luck to come out on top—and wasn't as lucky in 2033. As a result, his performance in the next year is likely to be closer to the average performance (i.e., he's regressed to the mean of stock picker performance). ◇

Example 9.4. If you administer a reading test to a group of learners one week and another reading test a couple of weeks later, you're likely to find that the very worst readers on the first test are still pretty poor readers on the second test (= the skill part), but their performance won't be as atrocious—it'll seem as though they've made some progress. Similarly, the best readers on the first test are likely to still be good readers on the second test, but their performance probably won't be as exceptional—it'll seem as though they've become worse.

But this pattern can be explained in terms of measurement error: Even if none of the learners actually learnt or unlearnt something, you're likely to find such a pattern. The reason is that, if you obtained a dismal score, you're likely to be a pretty poor reader *and* to have had some bad luck—perhaps the topic of the reading test just wasn't suited for you, or you were coming down with the flu. A couple of weeks later, you might encounter a topic you know a thing or two about or you might be in better physical shape. Similarly, if you scored exceptionally well on the first test, you may have had some luck with the test's topic or with other circumstances, and these may not be as favourable next time round. ◇

Exercise 9.5. A nationwide standardised maths test is administered to all 5th graders. It turns out that the classes with the highest mean test scores tend to be pretty small. One possible explanation is that small classes are more conducive to learning maths. Another explanation is that this finding is an artefact of measurement error. These explanations aren't mutually exclusive.

- (a) Explain how random measurement error can give rise to this finding.
- (b) How could you check if measurement error accounts (fully or in part) for this finding?

Hint: Which graph could you draw? ◇

Questionable research practices

Exercise 10.1 (Reading assignment). Read Chapter 2 in [Chambers \(2017\)](#). There are no guiding questions for this text; it should be intelligible enough. But by way of preparing for it, try to answer these questions.

- (a) You recruit 60 participants, aged 8–88. Half of them are assigned to the experimental group; the others to the control group (random assignment). You run a significance test comparing the mean age in both groups. What is, at most, the probability that you'll obtain a significant result (i.e., $p \leq 0.05$)?
- (b) Each of your participants throws a fair six-sided dice. You run another significance test to check if there's a mean difference in the number of pips obtained in the control and in the intervention groups. (Evidently, the intervention doesn't make you throw dice any better.) What is, at most, the probability that you'll obtain a significant result (i.e., $p \leq 0.05$)?
- (c) What is, at most, the probability of observing *either* a significant age difference between the two groups, *or* a significant difference in the mean number of pips obtained, *or* two significant differences? ◇

10.1 A paradox

[Sterling et al. \(1995\)](#) inspected 563 articles in psychology journals (published in 1986–1987) in which significance tests were used to answer the research question. In 538 of them (96%), the researchers reported a significant result that confirmed their own hypothesis. In medical journals, the figure was lower but still pretty high (270/316, 85%).

But at the same time, the sample sizes in psychological research are fairly small ([Marszalek et al., 2011](#); [Sedlmeier & Gigerenzer, 1989](#)): The average study in applied psychology published in 1995 only contained 22 participants per condition. This implies that many of these studies must have had fairly low statistical power (see Section 4.8 on page 47): Even if the null hypothesis hadn't been

Table 1. Outcomes of Tests of Significance for Four Psychology and Three Medical Research Journals

Journals	No. of articles reviewed in 1986–87	% articles reviewed that use tests in 1986–87	% articles using tests that reject H_0 in 1986–1987	No. of articles reviewed that used tests in 1958	% articles using tests that reject H_0 in 1958
<i>Experimental Psychology</i> (four journals)	165	92.73	93.46	106	99.06
<i>Comparative & Physiological Psychology</i> (two journals)	119	88.24	97.14	94	96.81
<i>Consulting & Clinical Psychology</i>	83	96.39	97.50	62	95.16
<i>Personality & Social Psychology</i>	230	97.83	95.56	32	96.88
Psychology Journals Total	597	94.30	95.56	294	97.28
<i>American Journal of Epidemiology</i>	141	81.56	80.87	N/A	N/A
<i>American Journal of Public Health</i>	97	43.30	88.10	N/A	N/A
<i>New England Journal of Medicine</i>	218	75.69	87.88	N/A	N/A
Medical Journals Total	456	69.25	85.40	N/A	N/A

Figure 10.1: Table 1 from Sterling et al. (1995).

correct in any of these studies, it'd have been impossible to reject it in 96% of cases.

For a long time (see already Sterling, 1959), it was believed that the reason for this discrepancy (low power, lots of significant results) was due to **publication bias**: Researchers prefer to write up the studies in which they obtained significant results, and editors and reviewers tend to reject studies with non-significant findings. The studies that were conducted but that produced non-significant findings were believed to languish in the researchers' file-drawers.

But while some studies never make it into print, the vast majority do. So where did the non-significant findings go?

10.2 Hidden flexibility

Eventually, scholars with an interest in meta-science (i.e., science about science) have come to realise that research projects afford a great deal of flexibility. Researchers can—consciously or subconsciously—leverage this flexibility to produce a steady stream of significant findings—even if the data are nothing but noise. (If the data aren't just noise, such flexibility will spuriously amplify the signal. For instance, even if A influences B , the literature as a whole will tend to overestimate the extent of this influence.) Simmons et al. (2011) call this flexibility **researcher degrees of freedom** and demonstrate how significant findings can be conjured from thin air if researchers afford themselves some leeway in analysing their data.

Sources of researcher degrees of freedom include:

- A researcher can run intermediate analyses and decide to stop or to continue collecting data based on the results. See Simmons et al. (2011) and Exercise 10.2 for the consequences of this.
- Sometimes, there are several ways in which a task or test can

be scored, or how some variable can be constructed. When one way yields a significant finding and the other doesn't, it's easy to convince yourself that the one that produced significance was obviously the right one. Relatedly, researchers routinely collect multiple outcome variables, but it's tempting to focus on the one that 'worked' (i.e., produced significance) rather than on those that didn't. See [Simmons et al. \(2011\)](#), [Gelman & Loken \(2013\)](#), and [von der Malsburg & Angele \(2017\)](#). For a discussion with a focus on bilingualism research, see [Poarch et al. \(2019\)](#).

- **HARKing** (hypothesizing after the results are known; [Kerr, 1998](#)): A largely exploratory analysis is reported as though it were planned all along. Inevitably, the researchers will find in the data what they claim to have anticipated.
- Convenient errors and biased debugging: Everyone makes mistakes, but you're more likely to catch your own mistakes when the results don't pan out than when they do. As a result, the mistakes that remain in the literature aren't distributed randomly but tend to favour the researchers' hypotheses.

Trying out several defensible analyses and glossing over the ones that didn't produce significance is referred to as **p-hacking**.

The practices listed above are examples of questionable research practices. Traditionally, these aren't viewed as outright fraud (which includes fabricating or manipulating data). But arguably, it should become increasingly difficult to invoke plausible deniability as professional researchers can be expected to know their consequences.

For possible solutions, see [Chambers \(2017\)](#).

Exercise 10.2 (False-positive psychology). The consequences of researcher degrees of freedom/p-hacking are best appreciated by seeing them. Do these exercises in order.

- Open the app at <https://janhove.github.io/qrp> and carefully read the description.
- Click 'Simulate!', leaving all settings at their default values. Describe what the two graphs (reproduced here as Figure 10.2 for your convenience) are showing.
- First try to answer the following questions by *thinking* about them. Once you've written down your answer, check it by running the simulation.
 - Increase the 'maximum number of additional participants in each group' to 30. Leave the other settings at their default values. How will the graphs change?
 - Leaving all other settings as they currently are, what will happen if instead of analysing the data after 10 new participants per condition, they're analysed after 5 new participants per condition? Or after just 2 new participants per

condition?

- iii. What will happen when the correlation between the two outcome variables becomes weaker (e.g., $r = 0.1$ instead of $r = 0.5$)? Why?
- iv. What will happen when the correlation between the two outcome variables becomes stronger (e.g., $r = 0.95$)? Why?
- v. For which combination of the different parameters will you obtain the highest Type-I error? Think before running the simulation!
- vi. For which combination of the different parameters will you find a Type-I error rate of about 5%? Are there any parameters that don't play a role? Think before running the simulation!




Figure 10.2: When you run the app using its default settings, you'll obtain two graphs similar to these. Your graphs won't be identical as they are based on simulations with random data.

10.3 *Further reading

Most studies referred to in this chapter are both accessible and short. If you read [Simmons et al. \(2011\)](#) (warmly recommended!), also read their short retrospective article ([Simmons et al., 2018](#)) lest you misinterpret the take-home message. [Peterson \(2016\)](#) presents an ethnographic study that gives you some insight into what questionable research practices look like in the field.

A highly accessible book-length treatment of these topics, and then some, which I cannot recommend highly enough, is [Ritchie \(2021\)](#). [Chambers \(2017\)](#) is also recommended.

A

Descriptive statistics

While this isn't a full-blown course in statistics, it is useful to know a few basic terms and concepts in descriptive statistics. Most of the terms we will need are covered in the reading assignment in Exercise [A.1](#).

Exercise A.1 (Descriptive statistics). Read [Johnson \(2013\)](#) with an eye towards explaining the following terms and concepts in a manner that you find intelligible by providing your own definition, clarifying example or illustration.

- (a) Continuous vs categorical variables (pp. 289–290 in [Johnson, 2013](#))
- (b) Histogram (pp. 292–293)
- (c) Bimodal distribution (p. 292)
- (d) Outliers (p. 292)
- (e) Normal distribution (pp. 293–294)
- (f) Arithmetic mean vs median vs mode (pp. 295–296)
- (g) The effect of outliers on the mean and median (p. 297)
- (h) Quantile, percentile, and quartile (p. 298)
- (i) Standard deviation and variance (p. 299)
- (j) Left- and right-skewed distributions (p. 301)
- (k) Ordinal vs nominal variables (p. 307)
- (l) Bar chart (pp. 307–308)
- (m) Contingency table (p. 311) ◇

Another useful concept is introduced in the next definition.

Definition A.2 (Probability of superiority). Let $\mathbf{a} = (a_1, \dots, a_m)$ and $\mathbf{b} = (b_1, \dots, b_n)$ be numeric vectors. Consider all $m \times n$ possible pairwise comparisons between an element from $\mathbf{a}$ and an element from $\mathbf{b}$. The proportion of such comparisons in which the element from $\mathbf{a}$ is larger than the element from $\mathbf{b}$ is called the **probability of superiority** of $\mathbf{a}$ over $\mathbf{b}$. Ties count as 'half-successes' towards this proportion. ◇

In psychology, the probability of superiority is also referred to as the **common-language effect size**.

Example A.3. Let $a = (1, 3, 5)$ and $b = (3, 4)$. There are six pairwise comparisons:

- $a_1 (1) < b_1 (3)$,
- $a_1 (1) < b_2 (4)$,
- $a_2 (3) = b_1 (3)$,
- $a_2 (3) < b_2 (4)$,
- $a_3 (5) > b_1 (3)$,
- $a_3 (5) > b_2 (4)$.

The element from a is strictly larger than the element from b in two out of six comparisons and equal in one comparison. Hence, the probability of superiority of a over b is $\frac{2+0.5}{6} \approx 42\%$. The superiority of b over a is thus about 58%. $\diamond$

You'll rarely explicitly encounter the probability of superiority when reading research articles in applied linguistics. But it's a statistic that can easily be interpreted, that can be used without any qualms when the data are ordinal and that is considerably less sensitive to outliers than is the arithmetic mean. It's also occasionally used implicitly in statistical tests (e.g., in the Mann–Whitney–Wilcoxon test).

B

Reading difficult results sections

Results sections in quantitative research reports can be daunting. Sometimes, the analyses are necessarily complex and require sophisticated knowledge about statistics and research design on the part of the reader. But too often, results sections are more difficult than they need to be (see [Vanhove, 2021](#)).

Don't allow yourself to be dazzled by complicated analyses and incomprehensible results sections—the complexity may be largely superficial.

Here are some tips for muddling through difficult results sections with minimal psychological damage, based on the blog entry [Surviving the ANOVA onslaught](#).)

1. Identify the central, genuine research questions and the corresponding hypotheses. Research papers in applied linguistics surprisingly often contain 'padding' research questions that are unrelated to the core goal of the study. When scanning the results section, you can usually leave aside the paragraphs about these uninteresting research questions. For example, in a report on a pretest/posttest experiment where participants were randomly assigned to conditions, you may find 'research' questions such as *Do participants in the treatment condition have different pretest scores from those in the control condition?* or *Do participants have higher scores on the posttest than on the pretest?* Both questions are uninteresting as they don't tell you whether the treatment actually worked.
2. **Draw a graph of the predictions.** (!) Having identified the key research questions and hypotheses, I often find it useful to sketch what the data would look like if the researchers' predictions panned out and what kind of data would, within reason, falsify their hypotheses. These graphs are usually simple hand-drawn line charts that illustrate the expected group differences. I find that they help me to better understand the logic behind the study and to focus on the important analyses in the Results section. You may find that several radically different patterns are in line

with the authors' stated predictions; this tells you something about how specific their predictions are. (It's good to have specific as opposed to very general hypotheses!) It can also be useful to draw some simple graphs of data that would *not* be consistent with the authors' predictions. This, too, can help you work out if the predictions are fairly specific (a good thing) or if pretty much any pattern in the data would be consistent with them (a bad thing).

3. Look for a graph in the paper. Ideally, the paper will contain a graph of the main results that you can then compare with the graphs you drew yourself. Do the results seem to confirm or disconfirm the researchers' predictions? Sometimes, a good graph will allow you to carry out the easiest of significance tests yourself: the **inter-ocular trauma test**—if the conclusion hits you between the eyes, it's significant. To be clear, this isn't a formal significance test. But it's a useful heuristic!

If the results are less clear cut, you'll need to scan the Results section for the more details, but by now, you should have a clearer idea of what you're looking for—and what you can ignore for now. If the paper doesn't contain a graph, you can often draw one yourself on the basis of the data provided in the tables.

4. Ignore tests unrelated to the central research questions. Results sections sometimes contain significance tests that are unrelated to the research questions the authors formulated up front (see [Vanhove, 2021](#)). Such tests include include balance tests in randomised experiments (e.g., "The control and intervention group did not differ in terms of SES ($t(36) = 0.8, p = 0.43$)."), tautological tests (e.g., "A one-way ANOVA confirmed that participants categorised as young, middle-aged and old differed in age ($F(2,57) = 168.2, p < 0.001$).") as well as some less obvious cases. By and large, these tests tend to add little to the study. In non-randomised experiments, systematic differences on background variables between the groups may represent confounds, but these can be assessed based on the descriptive statistics and don't need to be rubber-stamped with a significance test.

Evidently, you'll get better at this with practice, and it'll be helpful to educate yourself on basic statistics, too. The latter will help you to understand better what was done, but also it will also allow you to ask more critical questions, not least of which is *Are these analyses at all relevant?*.

C

Reporting research transparently

Many a research report leaves out information that is crucial for interpreting its findings correctly. And often, readers are implicitly asked to just take the authors word for it and trust that the analyses were run appropriately—even though reporting errors are common (Nuijten et al., 2016) and suboptimal or downright wrong analyses abound. Here are some tips to help ensure that your methods and findings are transparent to the readers.

1. Long reports detailing everything quickly become unreadable.

My own preference is to aim for a crisp main text that doesn't inundate the reader with numbers and numbing details. Instead, I try to communicate the findings mainly through plots and refer to copious online materials for the details (Vanhove, 2021, also contains further guidance for writing quantitative research reports and some useful references). Using these online materials, interested readers should at least be able to reproduce the results occurring in the report, and so the online materials that I make available minimally comprise the data sets and the computer code necessary for reproducing the plots and numbers in the main text. Materials such as stimulus lists, questionnaires, code for running the experiment itself etc. should in my view also be shared by default. For projects involving lots of tedious steps that will be of little interest to the average reader, I also like to make available a technical report that documents every little detail (e.g., Vanhove et al., 2019).

That said, **don't let perfect be the enemy of good**. If you're able to share your computer code but it's poorly documented, that's better than not sharing your code at all.

2. It's easier to share code, data, and materials if you made the decision to do so at the start of the project rather than a couple of weeks before handing in your report.
3. Nowadays I exclusively use <https://osf.io/> for making available online materials. See <https://osf.io/yxzfz/> for examples.
4. For further guidance, see Klein et al. (2018), Levenstein & Lyle (2018), and Soderberg (2018).

References

- Berthele, Raphael & Jan Vanhove. 2020. What would disprove interdependence? Lessons learned from a study on biliteracy in Portuguese heritage language speakers in Switzerland. *International Journal of Bilingual Education and Bilingualism* 23(5). 550–566. DOI: 10.1080/13670050.2017.1385590.
- Bialystok, Ellen, Fergus I. M. Craik, Raymond Klein & Mythili Viswanathan. 2004. Bilingualism, aging, and cognitive control: Evidence from the Simon task. *Psychology and Aging* 19(2). 290–303. DOI: 10.1037/0882-7974.19.2.290.
- Campbell, Donald T. & Julian C. Stanley. 1963. Experimental and quasi-experimental designs for research. In *Handbook of research on teaching*, Boston: Houghton Mifflin.
- Chambers, Chris. 2017. *The seven deadly sins of psychology: A manifesto for reforming the culture of scientific practice*. Princeton, NJ: Princeton University Press.
- Christenfeld, Nicholas J. S., Richard P. Sloan, Douglas Carroll & Sander Greenland. 2004. Risk factors, confounding, and the illusion of statistical control. *Psychosomatic Medicine* 66. 868–875. DOI: 10.1097/01.psy.0000140008.70959.41.
- De Wilde, Vanessa, Marc Brysbaert & June Eyckmans. 2020. Learning English through out-of-school exposure: How do word-related variables and proficiency influence receptive vocabulary learning? *Language Learning* 70(2). 349–381. DOI: 10.1111/lang.12380.
- Delsing, Lars-Olof & Katarina Lundin Åkesson. 2005. *Håller språket ihop Norden? En forskningsrapport om ungdomars förståelse av danska, svenska och norska*. Copenhagen: Nordiska ministerrådet.
- Ding, Peng. 2024. *A first course in causal inference*. CRC Press/Chapman & Hall. DOI: 10.120/9781003484080.
- Dutilh, Gilles, Alexandra Sarafoglou & Eric-Jan Wagenmakers. 2021. Flexible yet fair: blinding analyses in experimental psychology. *Synthese* 198(Suppl 23). S5745–S5772. DOI: 10.1007/s11229-019-02456-7.
- Elwert, Felix. 2013. Graphical causal models. In Stephen L. Morgan (ed.), *Handbook of causal analysis for social research*, 245–273. Dordrecht, The Netherlands: Springer. DOI: 10.1007/978-94-007-6094-3_13.

Fisher, R. A. 1935. *The design of experiments*. Edinburgh: Oliver and Boyd.

Gelman, Andrew & Eric Loken. 2013. The garden of forking paths: Why multiple comparisons can be a problem, even when there is no 'fishing expedition' or 'p-hacking' and the research hypothesis was posited ahead of time. URL http://www.stat.columbia.edu/~gelman/research/unpublished/p_hacking.pdf.

Goodman, Steven. 2008. A dirty dozen: Twelve *p*-value misconceptions. *Seminars in Hematology* 45. 135–140. DOI: 10.1053/j.seminhematol.2008.04.003.

Grosz, Michael P., Julia Rohrer & Felix Thoemmes. 2020. The taboo against explicit causal inference in nonexperimental psychology. *Perspectives on Psychological Science* 15(5). 1243–1255. DOI: 10.1177/1745691620921521.

Helsen, Werner F., Jan van Winckel & A. Mark Williams. 2005. The relative age effect in youth soccer across Europe. *Journal of Sports Science* 23(6). 629–636. DOI: 10.1080/02640410400021310.

Hill, Austin Bradford. 1965. The environment and disease: Association or causation? *Journal of the Royal Society of Medicine*.

Holland, Paul W. 1986. Statistics and causal inference. *Journal of the American Statistical Association* 81(396). 945–960. DOI: 10.2307/2289064.

Huitema, Bradley E. 2011. *The analysis of covariance and alternatives: Statistical methods for experiments, quasi-experiments, and single-case studies*. Hoboken, NJ: Wiley.

Imai, Kosuke, Gary King & Elizabeth A. Stuart. 2008. Misunderstandings between experimentalists and observationalists about causal inference. *Journal of the Royal Statistical Society: Series A (Statistics in Society)* 171. 481–502. DOI: 10.1111/j.1467-985X.2007.00527.x.

Imbens, Guido W. & Donald B. Rubin. 2015. *Causal inference for statistics, social, and biomedical sciences: An introduction*. Cambridge University Press. DOI: 10.1017/CBO9781139025751.

Johnson, Daniel Ezra. 2013. Descriptive statistics. In Robert J. Podesva & Devyani Sharma (eds.), *Research methods in linguistics*, 288–315. Cambridge: Cambridge University Press.

Kahan, Brennen C., Sunita Rehal & Suzie Cro. 2015. Risk of selection bias in randomised trials. *Trials* 16. 405. URL <http://www.trialsjournal.com/content/16/1/405>.

Kang, Sean H. K., Tamar H. Gollan & Harold Pashler. 2013. Don't just repeat after me: Retrieval practice is better than imitation for foreign vocabulary learning. *Psychonomic Bulletin & Review* 20. 1259–1265. DOI: 10.3758/s13423-013-0450-z.

Keele, Luke, Randolph T. Stevenson & Felix Elwert. 2019. The causal interpretation of estimated associations in regression

- models. *Political Science Research and Methods* 8(1). 1–13. DOI: 10.1017/psrm.2019.31.
- Kerr, Norbert L. 1998. HARKing: Hypothesizing after the results are known. *Personality and Social Psychology Review* 2(3). 196–217. DOI: 10.1207/s15327957pspr0203_4.
- Klein, Olivier, Tom E. Hardwicke, Frederik Aust, Johannes Breuer, Henrik Danielsson, Alicia Hofelich Mohr, Hans IJzerman, Gustav Nilsson, Wolf Vanpaemel & Michael C. Frank. 2018. A practical guide for transparency in psychological science. *Collabra: Psychology* 4(1). 20. DOI: 10.1525/collabra.158.
- Lardiere, Donna. 2006. Establishing ultimate attainment in a particular second language grammar. In ZhaoHong Han & Terrence Odlin (eds.), *Studies of fossilization in second language acquisition*, 35–55. Clevedon: Multilingual Matters.
- Levenstein, Margaret C. & Jared A. Lyle. 2018. Data: Sharing is caring. *Advances in Methods and Practices in Psychological Science* 1(1). 95–103. DOI: 10.1177/2515245918758319.
- Lorenzo, Francisco, Sonia Casal & Pat Moore. 2010. The effects of Content and Language Integrated Learning in European education: Key findings from the Andalusian Bilingual Sections Evaluation Project. *Applied Linguistics* 31(3). 418–442. DOI: 10.1093/applin/amp041.
- Ludke, Karen M., Fernanda Ferreira & Katie Overy. 2014. Singing can facilitate foreign language learning. *Memory & Cognition* 42. 41–52. DOI: 10.3758/s13421-013-0342-5.
- Marszalek, Jacob M., Carolyn Barber & Julie Kohlhart. 2011. Sample size in psychological research over the past 30 years. *Perceptual and Motor Skills* 112(2). 331–348. DOI: 10.2466/03.11.PMS.112.2.331-348.
- McShane, Blakeley B, David Gal, Andrew Gelman, Christian Robert & Jennifer L Tackett. 2019. Abandon statistical significance. *The American Statistician* 73(Sup1). 235–245. URL [10.1080/00031305.2018.1527253](https://doi.org/10.1080/00031305.2018.1527253).
- Mook, Douglas G. 1983. In defense of external invalidity. *American Psychologist* 38. 379–387. DOI: 10.1037/0003-066X.38.4.379.
- Nuijten, Michèle B., Chris H. J. Hartgerink, Marcel A. L. M. van Assen, Sacha Epskamp & Jelte M. Wicherts. 2016. The prevalence of statistical reporting errors in psychology (1985–2013). *Behavior Research Methods* 48(4). 1205–1226. DOI: 10.3758/s13428-015-0664-2.
- Oehlert, Gary W. 2010. *A first course in the design and analysis of experiments*. URL <http://users.stat.umn.edu/~gary/book/fcdae.pdf>.
- Offerman, Heather M. & Daniel J. Olson. 2016. Visual feedback and second language segmental production: The generalizability of pronunciation gains. *System* 59. 45–60. DOI:

10.1016/j.system.2016.03.003.

Oppenheimer, Daniel M., Tom Meyvis & Nicolas Davidenko. 2009. Instructional manipulation checks: Detecting satisficing to increase statistical power. *Journal of Experimental Social Psychology* 45. 867–872. DOI: 10.1016/j.jesp.2009.03.009.

Peterson, David. 2016. The baby factory: Difficult research objects, disciplinary standards, and the production of statistical significance. *Socius* 2. 1–10. DOI: 10.1177/2378023115625071.

Poarch, Gregory J., Jan Vanhove & Raphael Berthele. 2019. The effect of bidialectalism on executive function. *International Journal of Bilingualism* 23(2). 612–628. DOI: 10.1177/1367006918763132.

Quené, Hugo. 2010. How to design and analyze language acquisition studies. In Elma Blom & Sharon Unsworth (eds.), *Experimental methods in language acquisition research*, 269–284. Amsterdam: John Benjamins.

Richardson, John T. E. 2018. The use of Latin-square designs in educational and psychological research. *Educational Research Review* 24. 84–97. DOI: 10.1016/j.edurev.2018.03.003.

Ritchie, Stuart. 2021. *Science fictions*. Penguin.

Rohrer, Julia M. 2018. Thinking clearly about correlations and causation: Graphical causal models for observational data. *Advances in Methods and Practices in Psychological Science* 1(1). 27–42. DOI: 10.1177/2515245917745629.

Rosenberger, William F. & John M. Lachin. 2016. *Randomization in clinical trials: Theory and practice*. Wiley 2nd edn.

Sedlmeier, Peter & Gerd Gigerenzer. 1989. Do studies of statistical power have an effect on the power of studies? *Psychological Bulletin* 105. 309–316.

Simmons, Joseph P., Leif D. Nelson & Uri Simonsohn. 2011. False-positive psychology: Undisclosed flexibility in data collection and analysis allows presenting anything as significant. *Psychological Science* 22. 1359–1366. DOI: 10.1177/0956797611417632.

Simmons, Joseph P., Leif D. Nelson & Uri Simonsohn. 2018. False-positive citations. *Perspectives on Psychological Science* 13(2). 255–259. DOI: 10.1177/1745691617698146.

Slavin, Robert E., Nancy Madden, Malgarita Calderón, Anne Chamberlain & Megan Hennessy. 2011. Reading and language outcomes of a multiyear randomized evaluation of transitional bilingual education. *Educational Evaluation and Policy Analysis* 33(1). 47–58. DOI: 10.3102/0162373711398127.

Slevc, L. Robert & Akira Miyake. 2006. Individual differences in second language proficiency: Does musical ability matter? *Psychological Science* 17(8). 675–681. DOI: 10.1111/j.1467-9280.2006.01765.x.

Soderberg, Courtney K. 2018. Using OSF to share data: A step-

- by-step guide. *Advances in Methods and Practices in Psychological Science* 1(1). 115–120. DOI: 10.1177/2515245918757689.
- Sprietsma, Maresa. 2010. Effect of relative age in the first grade of primary school on long-term scholastic results: international comparative evidence using PISA 2003. *Education Economics* 18(1). 1–32. DOI: 10.1080/09645290802201961.
- Sterling, Theodore D. 1959. Publication decision and their possible effect on inferences drawn from tests of significance—or vice versa. *Journal of the American Statistical Association* 54. 30–34.
- Sterling, Theodore D., W. L. Rosenbaum & J. J. Weinkam. 1995. Publication decision revisited: The effect of the outcome of statistical tests on the decision to publish and vice versa. *The American Statistician* 49. 108–112. URL <http://www.jstor.org/stable/2684823>.
- Vanhove, Jan. 2015. Analyzing randomized controlled interventions: Three notes for applied linguists. *Studies in Second Language Learning and Teaching* 5. 135–152. DOI: 10.14746/ssllt.2015.5.1.7.
- Vanhove, Jan. 2017. Lexical richness of short French, German and Portuguese texts written by children (technical report). URL <https://osf.io/vw4pc/>.
- Vanhove, Jan. 2019. Metalinguistic knowledge about the native language and language transfer in gender assignment. *Studies in Second Language Learning and Teaching* 9(2). 397–419. DOI: 10.14746/ssllt.2019.9.2.7.
- Vanhove, Jan. 2020. Capitalising on covariates in cluster-randomised experiments. *PsyArXiv Preprints* DOI: 10.31234/osf.io/ef4zc.
- Vanhove, Jan. 2021. Towards simpler and more transparent quantitative research reports. *ITL - International Journal of Applied Linguistics* 172(1). 3–25. DOI: 10.1075/itl.20010.van.
- Vanhove, Jan & Raphael Berthele. 2017. Testing the interdependence of languages (HELASCOT project). In Raphael Berthele & Amelia Lambelet (eds.), *Heritage and school language literacy development in migrant children: Interdependence or independence?*, 97–118. Bristol: Multilingual Matters. DOI: 10.21832/9781783099054-007.
- Vanhove, Jan, Audrey Bonvin, Amelia Lambelet & Raphael Berthele. 2019. Predicting perceptions of the lexical richness of short French, German, and Portuguese texts using text-based indices. *Journal of Writing Research* 10(3). 499–525. DOI: 10.17239/jowr-2019.10.03.04.
- von der Malsburg, Titus & Bernhard Angele. 2017. False positives and other statistical errors in standard analyses of eye movements in reading. *Journal of Memory and Language* 94. 119–133. DOI: 10.1016/j.jml.2016.10.003.
- Westfall, Jacob & Tal Yarkoni. 2016. Statistically controlling for

confounding constructs is harder than you think. *PLOS ONE* 11(3). e0152719. DOI: 10.1371/journal.pone.0152719.

Winter, Bodo. 2019. *Statistics for linguists: An introduction using R*. Routledge. DOI: 10.4324/9781315165547.