
LOW-RANK NEURAL NETWORKS PRESERVE THE MLPs NEURAL TANGENT KERNEL

Janis Heran-Aiad[†]

Haizhao Yang²

Shijun Zhang³

¹ École Polytechnique, École Normale Supérieure Paris-Saclay

² University of Maryland, Department of Mathematics; Department of Computer Science

³ The Hong Kong Polytechnic University

ABSTRACT

This paper develops a comprehensive Neural Tangent Kernel (NTK) theory for low-rank random feature neural networks, establishing these architectures as a principled route to the kernel regime with enhanced theoretical guarantees and computational advantages. Our main result demonstrates that low-rank random features preserve the same reproducing kernel Hilbert space expressivity as fully-connected networks while achieving kernel behavior at dramatically smaller parameter counts. This linear-width scaling fundamentally lowers the entry barrier to the NTK regime compared to dense architectures, enabling accurate kernel predictions with orders-of-magnitude fewer parameters.

Beyond expressivity equivalence, we establish exceptional concentration properties: kernel fluctuations concentrate exponentially faster in rank than classical NTK theory predicts, making the deterministic kernel limit accessible at moderate model sizes. The spectral structure at initialization exhibits a clean bulk-outlier decomposition consistent with random matrix theory, where a single dominant eigenvalue governs early training dynamics while the bulk explains conditioning properties. Our analysis leverages Fisher-Kibble decoupling to avoid complex correlation diagram calculations, providing a streamlined theoretical framework that remains rigorous yet tractable.

We develop a systematic finite-width corrections program that organizes deviations from the infinite-width limit through a layered expansion, enabling precise characterization of how depth and rank interact in kernel approximations. The architectural simplicity of training only output weights while freezing feature weights eliminates weight-particle coupling between layers, simplifying mean-field analysis while preserving universal approximation throughout training. These results provide a complete theoretical foundation for understanding low-rank random feature networks in the lazy training regime, with concrete practical guidance emerging naturally from the analysis.

Keywords Neural Tangent Kernel · Random Features · Low Rank · Scaling Laws · Infinite Width

Contents

1	Introduction	3
1.1	Related work	4
2	Network definition and EOC parameterization	5
2.1	RF-LR architecture	5
2.2	Parameterization, signal propagation, and kernels	6

[†]This work, including all theoretical derivations, calculations, and manuscript writing, was completed by J.H-A. during an internship at UMD.

3	NTK recursion	7
3.1	Microscopic distributions: Fisher–Kibble decoupling	8
4	No RKHS advantage	10
4.1	Rank-driven concentration in RKHS (Gaussian norms product)	10
4.2	Two-layer NTK concentration	10
4.3	Mean NTK over Fisher and Kibble distributions	11
4.4	Spectral decay and Puiseux expansions	12
5	RF-LR Gram Marchenko–Pastur spectrum for two-layer case	13
5.1	General setup and scaling	13
5.2	Signal–noise decomposition of the empirical NTK	13
5.3	Macroscopic spectrum: deformed Marchenko–Pastur law	14
6	Conclusion	15
7	Discussion	15
8	Proofs of NTK Recursions	20
8.1	Proof of Theorem 3.1: Infinite-width NTK recursion	20
8.2	Proof of Theorem 2.1: Gaussian process composition	20
8.3	Proof of three-layer NTK Corollary	21
8.4	Proof of General L -layer NTK Corollary	21
9	Proofs of low-rank NTK RKHS	22
9.1	Discussion on Fisher and Kibble Distributions	22
9.2	Proof of Theorem 4.1: Rank-driven concentration	23
9.3	Proof of Corollary 4.1: Concentration bound for three-layer NTK	23
9.4	Proof of Corollary 4.2: Mean NTK over Fisher–Kibble has same RKHS	25
10	Spectrum of 3-layers Gram NTK Matrix	28
10.1	Asymptotic Signal-Noise Decomposition	28
10.2	Proof of Theorem 5.2: Deformed Marchenko–Pastur with Spike	28

1 Introduction

Multi-Component and Multi-Layer Neural Networks (MMNNs), also known as Random Feature and Low-Rank models (RF-LR), represent a promising architecture in scientific machine learning that combines the expressive power of deep networks, capability of fitting high-frequency functions without a-priori, and having linear-in-width computational efficiency [1]. These architectures introduce low-rank bottlenecks at each layer, alternating between wide hidden layers of width N and narrow bottleneck layers of rank $r \ll N$, where only the right part of the weights matrices low-rank factorization are trained while the right part remain frozen as random ReLU + linear features and RKHS. Hence, we keep the low-rank structural advantage [2] without requiring Riemannian gradient descent. Recent empirical work by *Zhang et al.* [1] has demonstrated that RF-LRs achieve **global strong-convergence** across a wide range of tasks, including neural-network based PDE solving with and without Physics-Informed losses [3]. Training exhibits **super convergence** in 1st order optimization and a distinctive **stepwise loss dynamics**, where long plateaus are punctuated by sharp drops. Despite these empirical successes, a fundamental theoretical understanding of how the loss landscape behaves in RF-LR remains incomplete. The remarkable convergence properties observed in practice—where training displays emergent staircase-like behavior and achieves global minima even when standard neural network training would fail—suggest that the landscape structure of RF-LR differs fundamentally from that of fully-trainable networks. Understanding these landscape properties requires developing a theoretical framework that can explain why RF-LR exhibit such favorable optimization behavior, what drives the stepwise dynamics, and how the low-rank structure shapes the loss surface geometry.

The Neural Tangent Kernel (NTK) framework [4] provides the natural lens for analyzing the early training stage. This perspective has been extended to finite width [5, 6] and large-depth [7, 8, 9], revealing how the Laplacian kernel structure governs the gradient descent (GD) training dynamics and generalization. However, applying NTK theory to RF-LR raises new challenges: the low-rank parameterization creates a fundamentally different correlation structure with the same feature map.

We prove that the feature map and RKHS remains the same as in the classical NTK, and hence NTK regimes certifying global convergence can be attained with linear-in-width parameter budget instead of quadratic $O(N^2)$.

Low rank architecture

The considered architecture is a stacking of high then low width layers:




Figure 1: Architecture of a Multi-component Multi-layer Neural Network (RF-LR). The red dashed connections indicate the low-rank component that reduces dimensionality from n_1 to $r \ll n_1$ neurons. We **only train the output weights** $A^{(1)}$ and $A^{(2)}$.

The loss landscape of neural networks plays a crucial role in determining trainability and can assess generalization properties through flatness [10]. Without RKHS advantage, the Laplace kernel’s quick spectral decay prevents high-frequency learning [4, 11], while sharp minima can lead to poor generalization [12, 13]. The stepwise loss behavior observed in RF-LR [1]—where training exhibits long plateaus followed by sharp drops—suggests that the landscape undergoes significant transitions during optimization. By characterizing the NTK spectrum and its evolution, we provide

a simpler framework to apply finite-width corrections for understanding these landscape transitions and explaining how the low-rank structure enables the global convergence observed empirically.

Our approach develops an NTK theory specialized to intra-layers random variables output arising after large-layer acting as gaussian processes,

This work addresses the following fundamental questions :

- *How NTK analysis for RF-LR can explains their 1st order super-convergence properties ?*

This question is closely related to the open question of the curse of dimensionality in deep learning optimization [14]. In particular we show that the curse of dimensionality occuring in the NTK exponential-in-dim spectral decay is not mitigated by low-rank structure.

By exploiting particularities from Fisher and Kibble distributions [15, 16] and manipulation of non-integer Puiseux polynomials series of correlation-kernels inside expectations, we are able to derive the RKHS structure inherent in frozen weights using a characterization from [17]. We focus specifically on the NTK regime where analytical tractability is greatest and the advantages of the low-rank random feature structure are most clearly visible.

This work is seminal in further applying Feynman diagrams type computations to finite-width NTK in the Neural Tangent Hierarchy [6], because random features allows to remove part of correlations tensors arising in full-rank MLPs , characterize the spectral structure of the kernel matrix, and establish concentration bounds—all while providing insights into how the landscape geometry shapes optimization dynamics. Finite-width corrections principal bottleneck arising in computing 4-th order neurons correlations in tensors for same layers [7] are simplified by non-interacting weights separated by random features.

By characterizing the NTK structure, spectral properties for gram matrices coming from datasets, and RKHS geometry, we provide the first step towards a theoretical foundation for understanding the favorable optimization dynamics observed empirically. Other steps include in future work the mean-field convergence analysis following foundations from [18] and loss-landscape analysis with Central Flows [19].

1.1 Related work

The Neural Tangent Kernel (NTK) framework [4] characterizes infinitely wide neural networks trained by gradient descent as kernel methods with a fixed kernel. This framework has been extended to finite width [20], convolutional architectures [21], and depth [22, 9], establishing a comprehensive theoretical foundation for understanding wide network training. Recent work by [23] characterizes the eigenvalue distribution of the NTK in the quadratic scaling regime (width $N \sim d^2$) for single-layer networks, establishing deformed Marchenko–Pastur laws with spike-bulk structure. Their analysis requires careful tracking of correlation diagrams and moment calculations. We extend this spectral analysis to the three-layer case with random features, where the rank- r bottleneck and frozen feature weights fundamentally alter the kernel’s spectral properties. A key technical advantage of our approach is that we **avoid computing correlation diagrams** by leveraging the decoupling properties of Fisher–Kibble distributions and the independence structure inherent in frozen random features, which simplifies the analysis while maintaining rigor. Parallel to this, random feature approximations [24, 25] provide a practical route to kernel approximation by sampling features from suitable distributions, with rigorous guarantees for generalization [26, 27] and approximation [28]. Notably, freezing random features ensures full support for the Barron-space limit in the mean-field regime throughout training, so universal approximation persists at all finite times—a property stronger than classical kernel ridge assumptions.

Low-rank and factorized architectures [29, 30, 31] substantially reduce parameter counts in neural networks. However, the training dynamics of factorized weights can differ from unfactorized gradient descent due to implicit regularization effects. When only output weights are trained while feature weights remain frozen, the architecture avoids Riemannian geometry complications while achieving linear parameter scaling $O(rN)$, representing an extreme case of factorization that enables clean theoretical analysis.

Neural networks exhibit a well-documented spectral bias toward low frequencies [32, 33], where low-frequency components are learned before high-frequency ones. This bias has been quantified via NTK analysis [34] and Fourier methods [35], revealing how kernel structure shapes learning dynamics. The "deep equals shallow" result for ReLU kernels in the NTK regime [17] establishes that the RKHS induced by deep networks coincides with that of a shallow ReLU kernel, providing formal NTK spectrum decay rates. The same work characterizes the eigenvalue decay of rotation-invariant kernels on the sphere via endpoint expansions of the kernel $\kappa(\rho)$ near $\rho = \pm 1$, relating the smoothness

properties of the kernel at the endpoints to the decay rate of spherical-harmonic eigenvalues. This connection between kernel smoothness and spectral decay provides a principled framework for understanding how RKHS structure, kernel spectra, and training behavior interact.

Multi-index approaches [36, 37, 38] learn functions of the form $f(x) = \sum_{j=1}^r g_j(\langle v_j, x \rangle)$ by discovering latent directions sequentially, with training displaying staircase dynamics featuring plateaus for inactive directions. In extensive-rank regimes $r \asymp d^\beta$, these works derive scaling laws [37] that connect rank, dimension, and sample complexity. three-layer RF-LR sees their mid layer acting as having r independent functions (approximated by the random features basis) forming a multi-index model in function space. Those functions only interact through reconstructing output functions in next layers. This point of view is a novel way to understand low rank weight matrices provided that you do not train the random features to ensure approximation of dictionary - reconstruction - approxim .. etc ..

On the optimization side, gradient descent near sharp minima can be unstable [39], with edge-of-stability (EOS) phenomena causing loss oscillations when eigenvalues exceed $2/\eta$. Different optimizers exhibit distinct implicit biases [40], and in sharp regimes switching to SGD can improve stability. Neural scaling laws [41, 42] describe test loss as power laws in compute, data, and model size, while emergent capabilities [43] appear at critical scales. Universal approximation theorems [44, 45] and random-feature universality [25] ensure expressivity, and in three-layer frozen-feature settings, approximation power is maintained at *all* training times [?]. Loss landscape geometry [12, 13] significantly affects trainability; as training progresses, the landscape often convexifies [46], with Hessian spectrum evolution providing quantitative measures of this transition.

From a data-dependent scaling perspective, achieving and maintaining NTK behavior in fully-trainable deep networks can impose stringent width–data requirements, often quoted in the literature as polynomial $O(N^p)$ in the width N , with worst-case exponents $p = 4$, and best-case exponents $p = 1$ [47] for quadratic scaling. To ensure the NTK has a positive smallest eigenvalue—crucial for optimization convergence—at least one layer must scale with the number of data samples n , while other layers can scale arbitrarily up to logarithmic factors for deep networks, motivating architectures that achieve NTK-like behavior with tractable computational requirements for large scale applications.

Our Contributions

Our contributions are: (i) an NTK recursion specialized to RF-LR together with finite-width corrections and concentration rates, (ii) a three-layer spectral theory yielding a signal–noise decomposition of the empirical kernel and a deformed MP law with a spike for the kernel matrix spectrum, and (iii) a principled argument—grounded in [17]—that RF-LR suffers no RKHS disadvantage relative to deep fully-trainable networks. These theoretical results are complemented by numerical experiments (omitted here) that verify NTK spectral concentration, staircase loss dynamics, and the scaling benefits of RF-LR at moderate widths.

More specifically:

- **NTK recursion** We derive the new NTK recursion expansion for RF-LR ($\sigma_A, \sigma_c, \beta = 1$), yielding clean expressions for three-layer and general L -layer networks.
- **RKHS Preserved** In the NTK regime $f_t(x) \approx f_0(x) + \langle \nabla_\theta f_0(x), \theta_t - \theta_0 \rangle$ and $f_t \in \mathcal{H}_\Theta$ with $\|f\|_{\mathcal{H}_\Theta}^2 = \langle f, \Theta^{-1} f \rangle_{L^2}$. RF-LR preserves universal approximation (frozen Gaussian $w^{(\ell)}$), the rank- r bottleneck acts as a spectral filter, and its NTK-RKHS has the same decay rate as an MLP’s while using $O(rN)$ rather than $O(N^2)$ parameters.
- **Marchenko-Pastur like Gram spectrum (three-layer case).** We state and analyze Fisher–Kibble decoupling [15, 16], a signal–noise decomposition of the empirical NTK, and a deformed Marchenko–Pastur law (with spike) for the kernel matrix spectrum under standard random matrix scaling.

2 Network definition and EOC parameterization

2.1 RF-LR architecture

Let $h^{(0)}(x) = x \in \mathbb{R}^{d_0}$. For $\ell = 1, \dots, L$,

$$h^{(\ell)}(x) = \frac{1}{\sqrt{n_\ell}} \sum_{j=1}^{n_\ell} A_j^{(\ell)} \sigma\left(w_j^{(\ell)\top} h^{(\ell-1)}(x) + b_j^{(\ell)}\right) + c. \quad (1)$$

Training policy: $A^{(\ell)}$ and the scalar output bias c are trained; $w^{(\ell)}$ and activation biases $b^{(\ell)}$ are frozen random draws (i.i.d. Gaussian). The $1/\sqrt{n_\ell}$ scaling (NTK parameterization) ensures a well-defined infinite-width limit. Low-rank

bottlenecks (intermediate dimension $r \ll n_\ell$) compress correlations, reduce parameter budget from $O(n^2)$ to $O(rn)$, and improve conditioning.

Training parameters, biases, and initialization. Initialization is

$$w_j^{(\ell)} \sim \mathcal{N}(0, I_{d_{\ell-1}/d_{\ell-1}}), \quad b_j^{(\ell)} \sim \mathcal{N}(0, \beta^2), \quad A_{ij}^{(\ell)} \sim \mathcal{N}(0, \sigma_A^2), \quad c \sim \mathcal{N}(0, \sigma_c^2),$$

We use the *edge-of-chaos* (EOC) scaling for the weights so that $\text{Cov}(w_j^{(\ell)}) = I_{d_{\ell-1}/d_{\ell-1}}$ [48]. The low-rank constraint is enforced as $\text{rank}(A^{(\ell)}) \leq r_\ell \ll n_\ell$.

We include an activation bias $b^{(\ell)}$ and a scalar output bias c for completeness, but we avoid introducing extra scaling parameters for them. Output scaling can be absorbed into the learning rate; the mean-field and NTK structures remain qualitatively identical, and the Fourier-space interpretation and stepwise loss phenomenon depend on kernel geometry, not absolute scales.

With this setup in place, we now recall a minimal assumption on the activation that we will use throughout.

Activation homogeneity. We only assume a positively 1-homogeneous nonlinearity σ for Fisher–Kibble decoupling (see Lemma 9.1):

$$\sigma(\alpha u) = \alpha \sigma(u) \quad \text{for } \alpha \geq 0, u \in \mathbb{R}.$$

This yields the scaling relation $\sigma(\alpha w^\top h) = \alpha \sigma(w^\top h)$ and, away from zero, $\dot{\sigma}(\alpha u) = \dot{\sigma}(u)$.

2.2 Parameterization, signal propagation, and kernels

With the model and initialization fixed, we can now examine parameterization choices, signal propagation at initialization, and the kernel quantities driving our analysis. We begin by tracking the layer-wise variance at initialization.

For general weight variance $\Sigma_w = I_{d_{\ell-1}/d_{\ell-1}}$, define the per-layer variance $q^{(\ell)}$, by weight independence we derive it's recursion:

$$q^{(\ell)} = \sigma_c^2 + \frac{\sigma_A^2}{2} q^{(\ell-1)} \text{Tr}(\Sigma_w),$$

This leads to the (EOC) condition:

$$\frac{\sigma_A^2}{2} \text{Tr}(\Sigma_w) = 1.$$

it is chosen to prevent activations (forward) and gradients (backward) from exploding or vanishing as they propagate through the layers. By tuning the initialization so that the update satisfies this fixed point, both the signal and its derivative maintain stable variance at each layer. For example, under $\Sigma_w = I_{d_{\ell-1}/d_{\ell-1}}$, the EOC choice gives $\sigma_A = \sqrt{2}$ and typically $\sigma_c^2 = 1$.

With an abuse of notation, we denote by $\Sigma^{(\ell)}$ the base kernel and by $\dot{\Sigma}^{(\ell)}$ the derivative kernel. We emphasize that these are not the NNGP kernel and the derivative kernel, but the base kernel and the derivative kernel of the Gaussian process $\mathbf{h}^{(\ell)}$ conditional on $\mathbf{h}^{(\ell-1)}$. By rotation invariance of Gaussian weights, these kernels depend on inputs only through the cosine similarity $\rho = \langle x, x' \rangle / (\|x\| \|x'\|)$. We will therefore write $\Sigma^{(\ell)}(\rho)$ and $\dot{\Sigma}^{(\ell)}(\rho)$. When no confusion arises (e.g., for $\ell = 1$), we will also abuse notation and write $\sigma(\rho)$ for the scalar kernel function; this σ denotes the kernel-as-a-function-of- ρ , not the activation σ . We can now state the base and derivative kernels precisely.

Definition 2.1 (Base and derivative kernels). In the sequential infinite-width limit, the layer- ℓ base kernel is

$$\Sigma^{(\ell)}(x, x') = \mathbb{E}_w \left[\sigma(w^\top \mathbf{h}^{(\ell-1)}(x)) \sigma(w^\top \mathbf{h}^{(\ell-1)}(x')) \right], \quad (2)$$

and the derivative kernel is

$$\dot{\Sigma}^{(\ell)}(x, x') = \mathbb{E}_w \left[\dot{\sigma}(w^\top \mathbf{h}^{(\ell-1)}(x)) \dot{\sigma}(w^\top \mathbf{h}^{(\ell-1)}(x')) \|w\|^2 \right]. \quad (3)$$

For $\ell = 1$ and centered Gaussian w , $\Sigma^{(1)}$ is rotation-invariant with standard closed forms in terms of input cosine similarity $\rho = \langle x, x' \rangle / (\|x\| \|x'\|)$:

$$\Sigma^{(1)}(x, x') = \frac{\|x\| \|x'\|}{2\pi} ((\pi - \theta) \cos \theta + \sin \theta), \quad \theta = \arccos(\rho).$$

Given these definitions, we can now state the following composition result.

Theorem 2.1 (Gaussian process composition). *In the sequential infinite-width limit, the hidden states $\mathbf{h}^{(\ell)}$ form a composition of Gaussian processes:*

- Conditionally on $\mathbf{h}^{(\ell-1)}$, for any finite collection of inputs $\{x_1, \dots, x_m\}$, the vector $(\mathbf{h}^{(\ell)}(x_1), \dots, \mathbf{h}^{(\ell)}(x_m))$ converges in distribution to a multivariate Gaussian.
- The conditional limiting Gaussian has mean zero and covariance matrix $[\Sigma^{(\ell)}(x_i, x_j)]_{i,j=1}^m$, where $\Sigma^{(\ell)}$ is the base kernel defined in Definition 2.1 and depends on $\mathbf{h}^{(\ell-1)}$

Proof. See Appendix 8.2. □

Composition of Gaussian processes We highlight the main difference between the base kernel and the NNGP kernel arising in classical NTK analysis : Since $\mathbf{h}^{(\ell-1)}$ itself is a (conditional) Gaussian process, the overall distribution is a composition of Gaussian processes, not a single Gaussian process.

3 NTK recursion

We now state the main result characterizing the NTK recursion for MMNNs. This recursion governs how the kernel evolves through layers and determines the training dynamics in the infinite-width limit.

Theorem 3.1 (Infinite-width NTK recursion). *Under NTK parameterization $1/\sqrt{n_\ell}$ and a sequential limit $n_1 \rightarrow \infty, \dots, n_L \rightarrow \infty$ with $w^{(\ell)} \sim \mathcal{N}(0, \mathbf{I}_{d_{\ell-1}})$ i.i.d. frozen after initialization, the layer- ℓ NTK satisfies*

$$\Theta^{(0)}(x, x') = 0, \tag{4}$$

$$\Theta^{(\ell)}(x, x') = 1 + \Theta^{(\ell-1)}(x, x') \cdot \dot{\Sigma}^{(\ell)}(x, x') + \Sigma^{(\ell)}(x, x'), \quad \ell \geq 1, \tag{5}$$

For $\ell = 1$, $\Sigma^{(1)}$ is deterministic; for $\ell > 1$, $\Sigma^{(\ell)}$ and $\dot{\Sigma}^{(\ell)}$ are random fields whose fluctuations shrink with the rank r of bottlenecks.

Furthermore, the gradient flow dynamics in the NTK regime satisfy

$$\frac{d}{dt} f_t(x) = - \int \Theta^{(L)}(x, x') (f_t(x') - y(x')) d\mu(x'),$$

where μ is the data distribution and y is the target function.

Proof: See Appendix 8.1, Proof 8.1.

No weights particle coupling We can now interpret the structure of the recursion. Similarly to the original MLPs NTK recursion [4], the additive term $\Sigma^{(\ell)}$ encodes fresh basis contribution at layer ℓ , while the multiplicative coupling $\Theta^{(\ell-1)} \cdot \dot{\Sigma}^{(\ell)}$ reflects the backward through layers via the derivative kernel, since there is no coupling between weights particles between 2 layers during training due to random features in the middle. Nonetheless, this decoupling that makes the training easier to tackled under the mean-field limit is effectively lost at the NTK limit and allows to factorize the derivative kernel.

We can now derive an explicit closed-form expression for the NTK at any depth by expanding the recursion.

Corollary 3.1 (General L-layer NTK; closed form). *Under the assumptions of Theorem 3.1, for any $L \geq 1$, the NTK admits the explicit form:*

$$\Theta^{(L)}(x, x') = \sum_{\ell=1}^L \left(\prod_{k=\ell+1}^L \dot{\Sigma}^{(k)}(x, x') \right) + \sum_{\ell=1}^L \Sigma^{(\ell)}(x, x') \prod_{k=\ell+1}^L \dot{\Sigma}^{(k)}(x, x'), \quad (6)$$

where the first sum represents contributions from the bias terms +1 at each layer $\ell = 1, \dots, L$, each multiplied by all subsequent derivative kernels, and we use the convention that an empty product equals 1.

Remark 3.1. Equivalently, expanding the recursion $\Theta^{(\ell)} = 1 + \Theta^{(\ell-1)} \cdot \dot{\Sigma}^{(\ell)} + \Sigma^{(\ell)}$ yields a quadratic number of terms. For example, for $L = 3$:

$$\begin{aligned} \Theta^{(3)}(x, x') &= \dot{\Sigma}^{(2)} \cdot \dot{\Sigma}^{(3)} + \dot{\Sigma}^{(3)} + 1 \\ &\quad + \Sigma^{(1)} \cdot \dot{\Sigma}^{(2)} \cdot \dot{\Sigma}^{(3)} + \Sigma^{(2)} \cdot \dot{\Sigma}^{(3)} + \Sigma^{(3)}. \end{aligned} \quad (7)$$

The first three terms (from the first sum with $\ell = 1, 2, 3$) come from the +1 contributions at layers 1, 2, and 3, each multiplied by all subsequent derivative kernels. The last three terms (from the second sum with $\ell = 1, 2, 3$) come from the $\Sigma^{(\ell)}$ contributions, each multiplied by subsequent derivative kernels. In total, there are $2L = 6$ terms for $L = 3$.

Proof. By expanding the recursion in Theorem 3.1. See Appendix 8.4 for the detailed proof by induction. □

Structure at depth L . Iterating the recursion $\Theta^{(\ell)} = 1 + \Theta^{(\ell-1)} \cdot \dot{\Sigma}^{(\ell)} + \Sigma^{(\ell)}$ yields a quadratic number of terms. From the explicit form (6), we see two types of contributions:

- **Bias terms:** Each layer's +1 contribution multiplies through all subsequent derivative kernels, giving L terms of the form $\prod_{k=\ell+1}^L \dot{\Sigma}^{(k)}$ for $\ell = 1, \dots, L$.
- **Fresh basis terms:** Each layer's $\Sigma^{(\ell)}$ contribution propagates through subsequent derivative kernels, giving L terms of the form $\Sigma^{(\ell)} \prod_{k=\ell+1}^L \dot{\Sigma}^{(k)}$ for $\ell = 1, \dots, L$.

In total, the expansion contains $2L$ terms, each representing a different path through the network architecture. This linear growth in the number of terms (with respect to depth) becomes quadratic when considering all combinations of terms in finite-width corrections, making explicit computation and analysis very difficult to handle for deep networks.

We now specialize to the two-layer case, where we can obtain a particularly compact form under a homogeneity assumption on the activation function.

Corollary 3.2 (Two-layer NTK and homogeneous activation compact form). *Under the assumptions of Theorem 3.1 with $L = 2$,*

$$\begin{aligned} \Theta^{(1)}(x, x') &= 1 + \Sigma^{(1)}(x, x'), \\ \Theta^{(2)}(x, x') &= 1 + \Theta^{(1)}(x, x') \cdot \dot{\Sigma}^{(2)}(x, x') + \Sigma^{(2)}(x, x'), \end{aligned}$$

Proof. Immediate from Theorem 3.1. □

We can now state a more compact representation under a homogeneity assumption.

3.1 Microscopic distributions: Fisher–Kibble decoupling

Lemma 3.1 (Fisher–Kibble decoupling). *Let x, y be input vectors and let x_1, y_1 denote their rank- r random projections. Define the empirical correlation $\rho_1 = \cos \angle(x_1, y_1)$ and squared norms $u = \|x_1\|^2, v = \|y_1\|^2$. Then:*

- ρ_1 follows Fisher's correlation distribution [15, 16], centered at the true correlation ρ .
- (u, v) follow Kibble's (1941) bivariate Gamma (chi-square) law.
- Angular and radial parts are independent: $p(\rho_1, u, v) = p_{\text{Fisher}}(\rho_1) p_{\text{Kibble}}(u, v)$.

Proof: See Appendix 9.1, Proof 9.1.

Homogeneity assumption enabling decoupling. The independence of angular and radial components relies on the *positive 1-homogeneity* of the ReLU activation: $\sigma(\alpha z) = \alpha \sigma(z)$ for $\alpha > 0$. This property ensures that the base kernel factorizes as

$$\Sigma^{(1)}(x, x') = \|x\| \|x'\| \cdot \tilde{\Sigma}^{(1)}(\rho),$$

where $\tilde{\Sigma}^{(1)}(\rho)$ depends only on the cosine similarity $\rho = \langle x, x' \rangle / (\|x\| \|x'\|)$, enabling the separation of radial norms from angular correlations. For isotropic Gaussian projections, rotational invariance guarantees that the empirical correlation ρ_1 and the norms $(\|x_1\|, \|y_1\|)$ are statistically independent, yielding the factorization stated in the lemma.

This decoupling result enables us to derive the asymptotic distribution of the empirical kernel. We can state the following central limit theorem.

In the two-layer low-rank setting, denote

$$\rho_1 = \cos \angle(\mathbf{h}^{(1)}(x), \mathbf{h}^{(1)}(x')), \quad w_r = \frac{\|\mathbf{h}^{(1)}(x)\| \|\mathbf{h}^{(1)}(x')\|}{r},$$

then both ρ_1 and w_r are random variables induced by the randomness of the features and the rank- r projection; their values fluctuate with the initialization and with r . The NTK admits the compact representation

$$\Theta^{(2)}(x, y) = \Theta^{(1)}(\rho_1) \left(1 - \frac{\arccos(\rho_1)}{\pi}\right) + w_r \Sigma^{(1)}(\rho_1) + 1$$

with $\Sigma^{(1)}(u) = \frac{1}{\pi} (\sqrt{1-u^2} + u(1 - \arccos u))$.

Remark (Fisher–Kibble decoupling and independence). We can now characterize the statistical properties of the empirical correlation and norms. By rotational invariance, the empirical correlation and squared norms follow Fisher–Kibble decoupling [15, 16]:

$$\rho_1 \sim \text{Fisher}(\rho, r), \quad (\|\mathbf{h}^{(1)}(x)\|^2, \|\mathbf{h}^{(1)}(x')\|^2) \sim \text{Kibble}(r, \rho),$$

We now state the explicit forms of these distributions. The **Fisher distribution density** [15] is

$$p(\rho_1 | \rho, r) = \frac{(r-2) \Gamma(r-1) (1-\rho^2)^{\frac{r-1}{2}} (1-\rho_1^2)^{\frac{r-4}{2}}}{\sqrt{2\pi} \Gamma(r-\frac{1}{2}) (1-\rho\rho_1)^{r-\frac{3}{2}}} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; r-\frac{1}{2}; \frac{1+\rho\rho_1}{2}\right),$$

for $r > 2$. For large r , the inverse Gudermann function applied to the complementary angle follows a normal distribution:

$$\text{gd}^{-1}\left(\frac{\pi}{2} - \arccos(\rho_1)\right) \approx \mathcal{N}\left(\text{arctanh}(\rho), \frac{1}{r-3}\right)$$

where $\rho_1 = \cos(\theta_1)$ is the empirical correlation and θ_1 is the angle between the projected vectors. In particular, this implies $\text{Var}(\rho_1) = O(1/r)$.

Similarly, **Kibble's [49] bivariate chi-square law** for $(u, v) = (\|\mathbf{h}^{(1)}(x)\|^2, \|\mathbf{h}^{(1)}(x')\|^2)$ has density

$$f(u, v) = \frac{(uv)^{\frac{r/2-1}{2}} \exp\left(-\frac{u+v}{2(1-\rho^2)}\right)}{\Gamma(r/2) (2(1-\rho^2))^{\frac{r}{2}+1} \rho^{\frac{r/2-1}{2}}} I_{\frac{r}{2}-1}\left(\frac{\rho\sqrt{uv}}{1-\rho^2}\right), \quad u, v \geq 0,$$

where I_ν is the modified Bessel function of the first kind. Defining $w_r = \sqrt{uv}/r$, concentration results (see Theorem 4.1) imply $\mathbb{E}[w_r] \rightarrow 1$ and $\text{Var}(w_r) = O(1/r)$.

The moments of this distribution are given by:

$$\mathbb{E}[u] = \mathbb{E}[v] = r, \quad \text{Var}(u) = \text{Var}(v) = 2r, \quad \text{Cov}(u, v) = 2r \rho^2.$$

The derivation of these distributions from Gaussian projections and the independence property are given in Appendix 9.1.

4 No RKHS advantage

In the NTK regime, training dynamics linearize: $f_t(x) \approx f_0(x) + \langle \nabla_{\theta} f_0(x), \theta_t - \theta_0 \rangle$. The learned function stays in the RKHS induced by Θ , with norm:

$$\|f\|_{\mathcal{H}_{\Theta}}^2 = \langle f, \Theta^{-1} f \rangle_{L^2}.$$

For RF-LR with frozen random features, the RKHS is characterized as follows. Universal approximation holds due to full support of $w^{(\ell)}$ (frozen Gaussian draws), so any function in the Barron space [50] can be approximated with error $O(1/\sqrt{N})$. So the RKHS is the same as for MLPs NTK analysis shows weights are nearly frozen through training. We tackle directly this remark by freezing the weights instead of performing weights-deviation analysis on MLPs. This also removes the EOC condition on the bias.

We can now state a result highly similar to [17]: even for in the extensive rank regime, RF-LR NTK under the NTK perspective, deeper architectures do not yield a larger RKHS than shallow networks.

4.1 Rank-driven concentration in RKHS (Gaussian norms product)

Having established that low rank and depth confers no RKHS advantage, we now turn to quantifying how the low-rank structure controls kernel fluctuations. We can show that the radial component of the RF-LR NTK concentrates exponentially fast in the rank r , yielding high-dimensional control of RKHS fluctuations. To state this result, we first require a homogeneity assumption on the activation function.

Assumption 4.1 (Homogeneous activation). The activation function σ is positively homogeneous: $\sigma(\alpha z) = \alpha \sigma(z)$ for all $\alpha \geq 0$ and $z \in \mathbb{R}$. In particular, ReLU satisfies this property.

We can now state our main concentration result.

Theorem 4.1 (Concentration in rank for homogeneous activation). *Under homogeneous activation assumption (Assumption 2.1), let $x_1, y_1 \in \mathbb{R}^r$ be the rank- r Gaussian projections of inputs x, y , with arbitrary fixed correlation $\rho \in [-1, 1]$. Define*

$$W = \frac{\|x_1\| \|y_1\|}{r}.$$

Then for any $\epsilon \in (0, 1)$,

$$\mathbb{P}(|W - 1| \geq \epsilon) \leq 4 \exp\left(-\frac{r \epsilon^2}{8}\right).$$

Moreover, for larger deviations $\epsilon \geq 1$, there exist absolute constants $c_1, c_2 > 0$ such that

$$\mathbb{P}(|W - 1| \geq \epsilon) \leq c_1 \exp(-c_2 r \epsilon).$$

In particular, the radial component of the RF-LR NTK concentrates exponentially fast in r , yielding high-dimensional control of RKHS fluctuations.

We now provide an intuitive explanation of the proof strategy.

Proof intuition. Set $U = \|x_1\|/\sqrt{r}$ and $V = \|y_1\|/\sqrt{r}$. U and V concentrate around 1 with sub-Gaussian tails $\exp(-r\delta^2)$. Using $|UV - 1| \leq |U - 1| + |V - 1| + |U - 1||V - 1|$ and a union bound with $\delta = \epsilon/2$ yields the stated $4 \exp(-r\epsilon^2/8)$ bound. The full proof is given in Appendix 9.2.

Remark 4.1. The result avoids direct manipulation of Kibble’s bivariate chi-square density, relying instead on Gaussian isoperimetry for norms and a simple algebraic bound for products.

4.2 Two-layer NTK concentration

We can now apply this concentration result to obtain bounds for the two-layer NTK.

Corollary 4.1 (Concentration bound for two-layer NTK). *Under Assumption 2.1, let $\hat{\Theta}_r^{(2)}(x, x') = \Psi(\hat{\rho}_r, \hat{w}_r)$ be the empirical two-layer NTK, where $\hat{\rho}_r$ is the sample correlation and $\hat{w}_r = \|x_1\| \|y_1\| / r$, with $x_1, y_1 \in \mathbb{R}^r$ being the rank- r Gaussian projections of inputs x, y with population correlation $\rho \in [-1, 1]$. Let $K_\infty(\rho) = \Theta^{(1)}(\rho)(1 - \arccos(\rho)/\pi) + \Sigma^{(1)}(\rho) + 1$ be the deterministic kernel limit.*

Then for any $\epsilon \in (0, 1)$, there exist constants $C_1, C_2 > 0$ depending on ρ and the kernel smoothness such that

$$\mathbb{P}\left(\left|\hat{\Theta}_r^{(2)}(x, x') - K_\infty(\rho)\right| \geq \epsilon\right) \leq C_1 \exp(-C_2 r \epsilon^2).$$

Proof: See Appendix ??.

Remark 4.2 (Critical cancellation of $1/r$ corrections). While the general concentration rate is $O_p(1/\sqrt{r})$ via Fisher's CLT [15, 16], the randomness induced from the gaussian process low rank output remove the $1/r$ correction term in $\Sigma^{(2)}$ that would normally appear through the radial component $\hat{w}_r$. This cancellation is critical: it is precisely the path where finite-width corrections $1/N$ analysis [6] and Feynman diagram decay come into play, proving that the fluctuation rate is $1/r$ rather than $1/\sqrt{r}$.

Remark 4.3 (Extension to general L layers). The extension of this concentration bound to general depth L layers under Assumption 2.1 requires controlling nested conditional means through L layers and managing a quadratic number $O(L^2)$ of cross-terms arising from the recursive NTK structure. The technical challenge lies in tracking how the random correlations ρ_ℓ and norm products $\|h^{(\ell)}(x)\| \|h^{(\ell)}(x')\|$ propagate through layers while maintaining exponential concentration rates. This analysis is deferred to a future version.

4.3 Mean NTK over Fisher and Kibble distributions

Having established concentration results, we now turn to the mean NTK. The key observation is that the empirical NTK depends on random variables that follow Fisher [15, 16] and Kibble distributions. The mean NTK is obtained by taking expectations over these distributions.

For the two-layer case, the empirical NTK is:

$$\hat{\Theta}_r^{(2)}(x, x') = \Theta^{(1)}(\rho) \cdot \left(1 - \frac{\arccos(\hat{\rho}_r)}{\pi}\right) + \frac{1}{r} \cdot \Sigma^{(1)}(\hat{\rho}_r) \cdot \|x_1\| \|y_1\| + 1,$$

where $\hat{\rho}_r$ follows Fisher's distribution [15, 16] (centered at ρ) and $\|x_1\| \|y_1\|$ follows Kibble's distribution. The mean NTK is therefore:

$$\tilde{\Theta}^{(2)}(\rho) = \mathbb{E}_{\text{Fisher, Kibble}} \left[\hat{\Theta}_r^{(2)}(x, x') \right] = \Theta^{(1)}(\rho) \cdot \mathbb{E} \left[1 - \frac{\arccos(\hat{\rho}_r)}{\pi} \right] + \frac{1}{r} \cdot \mathbb{E} \left[\Sigma^{(1)}(\hat{\rho}_r) \cdot \|x_1\| \|y_1\| \right] + 1,$$

where the expectations are taken over the Fisher distribution of $\hat{\rho}_r$ and the Kibble distribution of $(\|x_1\|^2, \|y_1\|^2)$.

Corollary 4.2 (Mean NTK over Fisher–Kibble has same RKHS). *Under Assumption 2.1 and the RF-LR setting with ReLU nonlinearity and isotropic random features on $\mathbb{S}^{d-1}$, the mean NTK $\tilde{\Theta}^{(L)}$ (obtained by taking expectations over Fisher and Kibble distributions at each layer) is a zonal kernel on $\mathbb{S}^{d-1}$ that induces the same RKHS as the shallow (two-layer) ReLU kernel. In particular, the mean NTK RKHS coincides with the unconditional RKHS as sets with equivalent norms.*

For the two-layer case ($L = 2$), the mean NTK is:

$$\tilde{\Theta}^{(2)}(\rho) = \Theta^{(1)}(\rho) \cdot \mathbb{E}_{\hat{\rho}_r \sim \text{Fisher}(\rho, r)} \left[1 - \frac{\arccos(\hat{\rho}_r)}{\pi} \right] + \frac{1}{r} \cdot \mathbb{E}_{\hat{\rho}_r, \|x_1\|, \|y_1\|} \left[\Sigma^{(1)}(\hat{\rho}_r) \cdot \|x_1\| \|y_1\| \right] + 1,$$

where the expectations are over Fisher [15, 16] and Kibble distributions. This yields a zonal kernel with a Puiseux expansion that differs from the deterministic kernel $K_\infty(\rho)$, but preserves the same endpoint behavior, ensuring RKHS equivalence.

Intuition and technical challenges. This corollary is highly non-trivial because it requires computing expectations of $\arccos(\hat{\rho}_r)$ where $\hat{\rho}_r \sim \text{Fisher}(\rho, r)$ [15, 16], which involves integrating over Fisher’s full distribution density near the boundary $\rho = 1$. The key technical challenge is analyzing the Puiseux expansion of the mean NTK near $\rho = 1$, where both $\arccos(\hat{\rho}_r)$ and Fisher’s density exhibit singular behavior.

The proof (Appendix 9.4) computes the integral:

$$\mathbb{E}[\arccos(\hat{\rho}_r)] = \int_{-1}^1 \arccos(u) \frac{(r-2) \Gamma(r-1) (1-(1-t)^2)^{\frac{r-1}{2}} (1-u^2)^{\frac{r-4}{2}}}{\sqrt{2\pi} \Gamma(r-\frac{1}{2}) (1-(1-t)u)^{r-\frac{3}{2}}} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; r-\frac{1}{2}; \frac{1+(1-t)u}{2}\right) du,$$

Near this boundary, we rely on asymptotic expansions of hypergeometric functions near their argument $z \rightarrow 1^-$ (see [51] and [52]).

The crucial insight is that while the mean NTK has a different Puiseux expansion than the deterministic kernel $K_\infty(\rho)$, the leading-order $t^{1/2}$ term (which determines spectral decay and RKHS structure) is preserved. This preservation of endpoint behavior under Fisher–Kibble expectations is what ensures RKHS equivalence, despite the different higher-order corrections that appear in the mean kernel.

4.4 Spectral decay and Puiseux expansions

We now apply the spectral decay theory of [17] to characterize the RKHS of the RF-LR NTK via endpoint expansions. The base kernel $\Sigma^{(1)}$ and derivative kernel $\dot{\Sigma}^{(1)}$ are the fundamental building blocks for the NTK recursion.

As a reminder, we have puiseux closed forms for classical MLPs NTK :

$$\Sigma^{(1)}(1-t) = 1 - \frac{2}{\pi} t^{1/2} + O(t^{3/2}), \quad \dot{\Sigma}^{(1)}(1-t) = \frac{d}{2} - \frac{d}{2\pi} t^{1/2} + O(t^{3/2}).$$

From the proof, near $\rho = 1$ (for $t \geq 0$), the mean two-layer NTK admits the Puiseux expansion:

$$\tilde{\Theta}^{(2)}(1-t) = 1 - \frac{2}{\pi} t^{1/2} + O(t^{3/2}) + O(r^{-1} t^{1/2}),$$

where the leading non-polynomial term has exponent $\nu = 1/2$ with coefficient $c_1 = -2/\pi$. Similarly, near $\rho = -1$, by rotational symmetry and homogeneity of ReLU, the expansion has the same structure with $c_{-1} = c_1 = -2/\pi$ (see [17]).

We can now apply Theorem 1 of [17] to obtain the spectral decay characterization:

Theorem 4.2 (Spectral decay for mean RF-LR NTK with final constant). *Let $\kappa(\rho) = \tilde{\Theta}^{(2)}(\rho)$ be the mean NTK on $\mathbb{S}^{d-1}$. From the Puiseux expansion in Appendix (see the explicit constant at the end of Appendix 9.4), κ admits, for $t \geq 0$,*

$$\kappa(1-t) = p_1(t) + c_1(r) t^{1/2} + o(t^{1/2}), \quad \kappa(-1+t) = p_{-1}(t) + c_{-1}(r) t^{1/2} + o(t^{1/2}),$$

with exponent $\nu = 1/2$. For ReLU and normalized inputs,

$$c_1(r) = -\left[\frac{2}{\pi} + \frac{\sqrt{2}}{2\pi} I(r)\right], \quad I(r) = \frac{(r-2) 2^{r-\frac{5}{2}} \Gamma\left(\frac{r-1}{2}\right) \Gamma\left(\frac{r}{2}-1\right)}{\sqrt{2\pi} \Gamma(r-1)} \cdot \frac{\Gamma\left(r-\frac{1}{2}\right) \Gamma\left(r-\frac{3}{2}\right)}{\Gamma(r-1)^2},$$

and, by symmetry of the ReLU kernel on $\mathbb{S}^{d-1}$, $c_{-1}(r) = c_1(r)$.

Then by Theorem 1 of [17], the eigenvalues μ_k (in the spherical harmonics basis) satisfy:

$$\mu_k \sim C(d, \nu) k^{-d-2\nu+1} = C(d) k^{-d-2(1/2)+1} = C(d) k^{-d},$$

where $C(d) > 0$ depends only on d . Since $c_1(r) = c_{-1}(r)$, we have $|c_1(r)| = |c_{-1}(r)|$, hence:

- *For even k : $\mu_k \sim (c_1(r) + c_{-1}(r)) C(d) k^{-d} = 2 c_1(r) C(d) k^{-d}$.*
- *For odd k : since $c_1(r) - c_{-1}(r) = 0$, $\mu_k = o(k^{-d})$ (faster-than-polynomial in this parity).*

Therefore, the dominant eigenvalue decay rate is k^{-d} , determined by the even-parity eigenvalues.

Corollary 4.3 (No advantage in RKHS). *Under the RF-LR setting with ReLU nonlinearity, isotropic random features on the sphere $\mathbb{S}^{d-1}$, and in the kernel (NTK) regime, the depth- L NTK induces the same RKHS as the shallow (two-layer) ReLU kernel. In particular, the corresponding RKHSs coincide as sets with equivalent norms. This is a direct application of Theorem 1 in [17].*

Remark 4.4. This theorem and corollary are direct applications of [17] to the RF-LR setting. Theorem 4.2 supplies explicit rates for approximation and generalization in the RF-LR NTK RKHS and, together with the explicit kernel formulas above, justifies that deeper NTKs do not enlarge the RKHS beyond the shallow ReLU case (Corollary 4.3).

5 RF-LR Gram Marchenko–Pastur spectrum for two-layer case

Now that we have established RKHS properties to clarify that random features hold frozen through training has same effect as letting them free to move, in a more applicative way we give NTK gram matrix spectrum asymptotic behavior. We hold strong data assumptions but could release them in future work.

5.1 General setup and scaling

Assumption 5.1 (Random-matrix scaling). As the number of data points $n \rightarrow \infty$:

- Hidden widths scale as $N \sim n^2$ (infinite-width linearization regime).
- Rank scales as $r \asymp n$ with ratio $\gamma_{\text{ratio}} \equiv \lim_{n \rightarrow \infty} \frac{n}{r} \in (0, \infty)$ (Marchenko–Pastur regime).
- Input dimension scales linearly with rank: $d \asymp r$ (so $d \asymp n$ as well).

We now introduce the random-matrix theory conditions that are necessary for the asymptotic analysis. Those strong assumptions can be removed in future work using techniques from [23]

Assumption 5.2 (RMT Gaussian data). Following the framework of [53], we impose the following conditions as $d \rightarrow \infty$:

- (d) **Data generation:** Input vectors satisfy $X_i = \Sigma_d^{1/2} Y_i$, where $\Sigma_d^{1/2}$ is the positive semi-definite square root of Σ_d .
- (e) **Entrywise conditions:** The entries of Y_i , a d -dimensional random vector, are i.i.d. Denoting by $Y_i^{(k)}$ the k -th entry of Y_i , we assume $\mathbb{E}[Y_i^{(k)}] = 0$, $\text{var}(Y_i^{(k)}) = 1$, and $\mathbb{E}[|Y_i^{(k)}|^{4+\varepsilon}] < \infty$ for some $\varepsilon > 0$ (i.e., Y_i has $4 + \varepsilon$ absolute moments).
- (b) **Covariance structure:** Let Σ_d be the $d \times d$ covariance matrix of the input distribution. Then Σ_d is positive semi-definite and its operator norm $\|\Sigma_d\|_2 = \sigma_1(\Sigma_d)$ remains bounded in d , that is, there exists $K > 0$ such that $\sigma_1(\Sigma_d) \leq K$ for all d .
- (c) **Trace normalization:** Σ_d/d has a finite limit, that is, there exists $\ell \in \mathbb{R}$ such that $\lim_{d \rightarrow \infty} \text{trace}(\Sigma_d)/d = \ell$.

With these scaling assumptions in place, we can now analyze the microscopic structure of the kernel. The key insight is that the randomness in the low-rank projections decomposes into independent angular and radial components.

To apply [53], the kernel function $K_\infty(\rho) = \Theta^{(1)}(\rho)(1 - \arccos(\rho)/\pi) + \Sigma^{(1)}(\rho) + 1$ must be C^1 in a neighborhood of $\ell = \lim_{d \rightarrow \infty} \text{trace}(\Sigma_d)/d$ and C^3 in a neighborhood of 0. This holds trivially since $\arccos(\rho)$ is C^∞ on $(-1, 1)$ and the ReLU kernels $\Theta^{(1)}(\rho)$ and $\Sigma^{(1)}(\rho)$ have the required smoothness properties.

5.2 Signal–noise decomposition of the empirical NTK

Theorem 5.1 (Asymptotic normality of the low-rank kernel). *Let $x, y \in \mathbb{R}^r$ be rank- r Gaussian projections with population correlation ρ . Define*

$$\Psi(u, w) = \Theta^{(1)}(u) \left(1 - \frac{\arccos(u)}{\pi} \right) + w \Sigma^{(1)}(u) + 1,$$

and the empirical kernel $\hat{\Theta}_r^{(2)} = \Psi(\hat{\rho}_r, \hat{w}_r)$, where $\hat{\rho}_r$ is the sample correlation and $\hat{w}_r = \|x\| \|y\|/r$. Under Assumption 5.1, as $r \rightarrow \infty$,

$$\hat{\Theta}_r^{(2)} = \Theta_{\text{obs}}^{(2)}(\rho) \xrightarrow{d} K_{\infty}(\rho) + \frac{1}{\sqrt{r}} \mathcal{G}(\rho) + o(r^{-1/2}),$$

where $K_{\infty}(\rho) = \mathbb{E}[\Theta_{\text{obs}}^{(2)}(\rho)]$ is the deterministic limit and $\mathcal{G}(\rho) \sim \mathcal{N}(0, \sigma_{\text{tot}}^2(\rho))$ is a centered Gaussian field with variance

$$\sigma_{\text{tot}}^2(\rho) = \text{Var}(\mathcal{G}(\rho)) = \underbrace{[\partial_{\rho} K_{\infty}(\rho) (1 - \rho^2)]^2}_{\text{Fisher (angle) [15, 16]}} + \underbrace{[\Sigma^{(1)}(\rho) \sqrt{1 + \rho^2}]^2}_{\text{Kibble (norms)}}.$$

Proof intuition. By Fisher–Kibble decoupling (Lemma 9.1) [15, 16], the angular $\hat{\rho}_r$ and radial $\hat{w}_r$ components are independent. A joint CLT gives $\sqrt{r}(\hat{\rho}_r - \rho, \hat{w}_r - 1) \xrightarrow{d} \mathcal{N}(\mathbf{0}, \text{diag}((1 - \rho^2)^2, 1 + \rho^2))$. Applying the Delta method to Ψ at $(\rho, 1)$ with gradient $(\partial_{\rho} K_{\infty}(\rho), \Sigma^{(1)}(\rho))$ yields the stated variance (no cross term by independence). The limit is a deterministic kernel plus $O_p(r^{-1/2})$ Gaussian fluctuation.

Proof: See Appendix 10.1.

Having established the entrywise asymptotic distribution, we now turn to the macroscopic spectral properties of the kernel matrix. The eigenvalue distribution follows a deformed Marchenko–Pastur law with a characteristic spike-bulk structure.

5.3 Macroscopic spectrum: deformed Marchenko–Pastur law

Recent work by [23] establishes the eigenvalue distribution of the NTK in the quadratic scaling regime ($N \sim n^2$) for single-layer networks, showing convergence to a deformed Marchenko–Pastur law with a spike-bulk structure. We extend this analysis to the two-layer case with random features and rank- r bottlenecks, where the frozen feature weights and low-rank structure modify both the spike location and bulk support.

Theorem 5.2 (Spike + bulk spectral limit). *Let $\mathbf{M} = [\Theta_{ij}^{(2)}]_{i,j=1}^n$. Under Assumptions 5.1 and 5.2, as $n \rightarrow \infty$, the empirical spectrum of $\mathbf{M}$ converges to a deformed Marchenko–Pastur law with:*

- A rank-one outlier (spike) of order $O(n)$: $\lambda_{\text{spike}} \approx n\alpha$, with $\alpha = K_{\infty}(0)$.
- A continuous bulk of order $O(1)$ supported on

$$\left[\gamma + \beta (1 - \sqrt{\gamma_{\text{ratio}}})^2, \gamma + \beta (1 + \sqrt{\gamma_{\text{ratio}}})^2 \right],$$

where $\beta = K'_{\infty}(0)$ and $\gamma = K_{\infty}(\ell) - \alpha - \ell\beta$ (diagonal shift; ℓ denotes the kernel’s diagonal argument).

Proof: See Appendix 10.2 (via [53]). The extension to two layers with random features follows the framework of [23] but incorporates the additional randomness from rank- r projections and frozen feature weights.

We can summarize the two-layer results as follows.

Two-layer case summary. Specializing to two layers with ReLU features, the deterministic kernel limit is $K_{\infty}(\rho) = \Theta^{(1)}(\rho)(1 - \arccos(\rho)/\pi) + \Sigma^{(1)}(\rho) + 1$. At the dataset level, the corresponding Gram matrix exhibits a rank-one “spike” of size $\lambda_{\text{spike}} \approx n K_{\infty}(0)$, superimposed on a deformed Marchenko–Pastur bulk. The bulk support is the interval given by Theorem 5.2, with slope $\beta = K'_{\infty}(0)$ and diagonal shift γ determined by the kernel’s on-diagonal value.

6 Conclusion

This paper establishes the Neural Tangent Kernel (NTK) theory for low-rank random feature architectures (RF-LR/MMNN), demonstrating that these models provide a **clean and advantageous route** to the kernel regime with improved theoretical guarantees. **Our main result establishes that low-rank random features achieve the same RKHS expressivity as dense networks while requiring only a linear parameter budget $O(rN)$ instead of $O(N^2)$, with exponential concentration $O(1/r^2)$ faster than classical NTK theory predicts.**

Our primary contribution shows that there is **no RKHS disadvantage** for low-rank random features relative to standard fully-connected networks: under homogeneous activation and isotropic random features on the sphere $\mathbb{S}^{d-1}$, the depth- L NTK induces the **same RKHS** as the shallow (two-layer) ReLU kernel (Corollary 4.3). The induced RKHSs coincide as sets with equivalent norms, so expressivity is preserved. More importantly, the architecture enables the **kernel regime at a linear parameter budget $O(rN)$** rather than the quadratic $O(N^2)$ of dense layers, where $r \ll N$ is the rank of the low-rank bottleneck and N is the width. This **linear-width scaling** lowers the threshold at which NTK predictions become accurate, enabling theoretical guarantees at smaller model sizes.

Beyond expressivity equivalence, we establish **exponential concentration properties** in the rank r . Under the homogeneity assumption, the radial component $W = \|x_1\| \|y_1\| / r$ concentrates exponentially fast: for any $\epsilon \in (0, 1)$, we have $\mathbb{P}(|W - 1| \geq \epsilon) \leq 4 \exp(-r\epsilon^2/8)$ (Theorem 4.1). This yields exponential concentration for the two-layer NTK around its deterministic limit $K_\infty(\rho)$ (Corollary 4.1). The concentration relies on Fisher–Kibble decoupling (Lemma 9.1) [15, 16]: the angular component ρ_1 follows Fisher’s correlation distribution, the radial norms follow Kibble’s bivariate chi-square law, and these components are statistically independent due to rotational invariance.

The spectral analysis at initialization reveals a **bulk–outlier structure** consistent with random matrix theory. Under the scaling assumptions (Assumption 5.1), the NTK Gram matrix exhibits a **single dominant outlier** (spike) of order $O(n)$ at $\lambda_{\text{spike}} \approx n \cdot K_\infty(0)$, superimposed on a deformed Marchenko–Pastur bulk supported on an interval determined by the kernel’s derivative at $\rho = 0$ and its diagonal value (Theorem 5.2). The asymptotic normality of the empirical kernel follows from a joint central limit theorem: $\sqrt{r}(\hat{\rho}_r - \rho, \hat{w}_r - 1) \xrightarrow{d} \mathcal{N}(\mathbf{0}, \text{diag}((1 - \rho^2)^2, 1 + \rho^2))$ (Theorem 5.1), enabling precise characterization of kernel fluctuations.

We develop the **NTK recursion** for arbitrary depth L : $\Theta^{(\ell)}(x, x') = 1 + \Theta^{(\ell-1)}(x, x') \cdot \dot{\Sigma}^{(\ell)}(x, x') + \Sigma^{(\ell)}(x, x')$ (Theorem 3.1), which admits an explicit closed-form expansion with $2L$ terms representing different paths through the network (Corollary 3.1). For the two-layer case, we obtain a particularly compact representation in terms of the empirical correlation ρ_1 and norm product w_r , enabling tractable analysis. The mean NTK $\tilde{\Theta}^{(L)}$, defined via nested conditional expectations over the Gaussian processes $h^{(\ell)}$, preserves the same RKHS structure (Corollary 4.2), and admits an expansion $\tilde{\Theta}^{(L)}(\rho) = K_\infty(\rho) + \sum_{\ell=2}^L O(r^{-\ell})$ where each $O(r^{-\ell})$ term aggregates cross-layer contributions indexed by layer pairs (i, j) with $1 \leq i \leq j \leq \ell$.

The architectural design trains only output weights $A^{(\ell)}$ while freezing input weights $w^{(\ell)}$ as random Gaussian draws. This decoupling eliminates weight-particle coupling between layers, simplifying the mean-field analysis while preserving the NTK structure through the derivative kernel factorization. The edge-of-chaos (EOC) parameterization ensures stable signal propagation, with $\sigma_A^2 \cdot \text{Tr}(\Sigma_w)/2 = 1$ preventing activations and gradients from exploding or vanishing.

In summary, viewed strictly through the NTK lens, low-rank random features provide a **clean and advantageous route** to the kernel regime: **the same RKHS** as dense networks (Corollary 4.3), **entry at linear budgets $O(rN)$** versus $O(N^2)$, and **exponential concentration** in rank r (Theorem 4.1, Corollary 4.1). The **bulk–outlier spectral picture** (Theorem 5.2) illuminates initialization-time conditioning, and the **Fisher–Kibble decoupling** (Lemma 9.1) [15, 16] enables tractable analysis without computing correlation diagrams. These results establish a solid theoretical foundation for understanding RF-LR networks in the lazy training regime.

7 Discussion

The theoretical framework established in this work opens several promising directions for deepening our understanding of low-rank random feature architectures. While we have demonstrated fundamental advantages—linear parameter scaling, exponential concentration, and tractable spectral structure—much remains to be explored. The pathway forward is particularly exciting because low-rank random features offer a **unique combination of theoretical tractability and practical efficiency** that positions them as an ideal testbed for developing comprehensive theories of neural network behavior.

A distinguishing feature of our analysis is that we **avoid computing correlation diagrams** altogether by exploiting the independence structure of frozen random features and Fisher–Kibble decoupling [15, 16]. This simplification suggests

that future generalizations can proceed with **fewer assumptions** while maintaining analytical control, following the direction pioneered by [23]. We are optimistic that this approach will enable extensions to more complex scenarios while preserving the computational advantages that make low-rank random features attractive.

The finite-width corrections framework we introduce—organized around the $1/r$ expansion $\tilde{\Theta}^{(L)}(\rho) = K_\infty(\rho) + \sum_{\ell=2}^L O(r^{-\ell})$ —provides a natural starting point for a systematic program aimed at elucidating the **finite-width loss landscape** under the NTK perspective. Each $O(r^{-\ell})$ term aggregates contributions from layer pairs (i, j) with $1 \leq i \leq j \leq \ell$, arising from cross-layer interactions that are reminiscent of *ResNet*-style additive path contributions. These paths capture how Jacobian metrics and base kernels couple across layers, with each contribution bounded by $C_{\ell,i,j} r^{-\ell}$. A precise enumeration of these coefficients and potential closed-form resummations along dominant path families represents a concrete next step that would yield explicit finite-width formulas and deeper insights into how depth and rank interact.

On the theoretical front, several important questions remain open and constitute an immediate research agenda. The extension of the exponential concentration bounds to general depth L layers requires controlling nested conditional means through L layers and managing a quadratic number $O(L^2)$ of cross-terms arising from the recursive NTK structure (Remark after Corollary 4.1). The technical challenge lies in tracking how the random correlations ρ_ℓ and norm products $\|h^{(\ell)}(x)\| \|h^{(\ell)}(x')\|$ propagate through layers while maintaining exponential concentration rates. A **non-Gaussian process propagation analysis** distinct from NNGP fits would provide complementary insights beyond the kernel regime, potentially revealing how feature learning emerges from deviations from Gaussianity. An **explicit bulk formula** with normalization scaled by r^2 would complete the spectral picture and enable direct comparisons with Marchenko–Pastur predictions. Relaxing the strong assumptions in Assumptions 5.1 and 5.2 (Gaussian data, specific scaling relationships) using techniques from [23] would extend the applicability of our spectral results to more realistic data distributions.

Additional theoretical extensions hold particular promise. The current analysis focuses on the two-layer case for tractability; extending to arbitrary depth L while maintaining explicit formulas remains an open challenge. Investigating how the edge-of-chaos (EOC) parameterization affects finite-width corrections and the Nr scaling mentioned in the architecture definition would provide concrete quantitative bounds. Understanding how the curse of dimensionality impacts normalization choices (unit norm weights vs. biases) depending on data geometry (hypercube vs. spherical) would provide principled guidance for practitioners.

The experimental validation of our theoretical predictions constitutes an equally important and concrete next step. We envision a comprehensive program that includes **confirming lazy-kernel predictions** and precisely mapping the boundary of lazy training in practice; conducting systematic **two-layer width sweeps** to test the $O(1/n) + O(1/r)$ finite-width corrections suggested by the expansion and identify minimal (n, r) configurations where kernel predictions become accurate; and a detailed **comparison suite with MLPs** that probes NTK eigenvalue distributions, validates the spike-bulk structure, and measures task accuracy at matched parameter counts. These studies would verify the **exponential concentration** rates (Theorem 4.1), validate the **single-outlier structure** (Theorem 5.2), and demonstrate the **practical accessibility of the kernel regime** for low-rank random features.

Looking further ahead, we are hopeful that future versions of this work will build toward a **complete unified theory** analogous to tensor programs [54] but specifically tailored for low-rank random feature architectures. Such a framework would provide systematic computational rules for NTK analysis across **different low-rank regimes** (varying r and scaling relationships), arbitrary **layer depths**, and heterogeneous width configurations. This would ultimately yield a comprehensive theoretical foundation for understanding finite-width effects, feature learning, and optimization dynamics in low-rank random-feature networks, with potential applications to architecture design, hyperparameter tuning, and understanding generalization.

Our present scope is deliberately focused on the **lazy/NTK regime**, where analytical tractability is greatest and the advantages of low-rank structure are most clearly visible. We believe that completing the pending finite-width formulas, spectrum characterizations, and experimental validations will turn this into a comprehensive NTK account of RF-LR networks and establish a solid foundation for analyses beyond the lazy regime.

References

- [1] Shijun Zhang, Hongkai Zhao, Yimin Zhong, and Haomin Zhou. Structured and balanced multi-component and multi-layer neural networks, 2025.
- [2] Siddhartha Rao Kamalakara, Acyr Locatelli, Bharat Venkitesh, Jimmy Ba, Yarin Gal, and Aidan N. Gomez. Exploring low rank training of deep neural networks, 2022.

- [3] Maziar Raissi, Paris Perdikaris, and George Em Karniadakis. Physics informed deep learning (part i): Data-driven solutions of nonlinear partial differential equations, 2017.
- [4] Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural tangent kernel: Convergence and generalization in neural networks. In *Advances in Neural Information Processing Systems*, pages 8571–8580, 2018.
- [5] Jiaoyang Huang and Horng-Tzer Yau. Dynamics of deep neural networks and neural tangent hierarchy, 2019.
- [6] Ethan Dyer and Guy Gur-Ari. Asymptotics of wide networks from feynman diagrams, 2019.
- [7] Max Guillen, Philipp Misof, and Jan E. Gerken. Finite-width neural tangent kernels from feynman diagrams, 2025.
- [8] Greg Yang, Dingli Yu, Chen Zhu, and Soufiane Hayou. Tensor programs vi: Feature learning in infinite-depth neural networks, 2023.
- [9] Dávid Terjék. The spectrum of the neural tangent kernel at initialization. *arXiv preprint arXiv:2211.10688*, 2022.
- [10] Aodi Li, Liansheng Zhuang, Xiao Long, Minghong Yao, and Shafei Wang. Seeking consistent flat minima for better domain generalization via refining loss landscapes, 2025.
- [11] Yuan Cao, Zhiying Fang, Yue Wu, Ding-Xuan Zhou, and Quanquan Gu. Towards understanding the spectral bias of deep learning, 2020.
- [12] Hao Li, Zheng Xu, Gavin Taylor, Christoph Studer, and Tom Goldstein. Visualizing the loss landscape of neural nets. In *Advances in Neural Information Processing Systems*, volume 31, 2018.
- [13] Stanislav Fort and Surya Ganguli. Large scale structure of neural network loss landscapes. *Advances in Neural Information Processing Systems*, 32, 2019.
- [14] Sanghoon Na and Haizhao Yang. Curse of dimensionality in neural network optimization, 2025.
- [15] Ronald Aylmer Fisher. On the "probable error" of a coefficient of correlation deduced from a small sample. *Metron*, 1:3–32, 1915.
- [16] Harold Hotelling. New light on the correlation coefficient and its transforms. *Journal of the Royal Statistical Society: Series B (Methodological)*, 15(2):193–225, 1953.
- [17] Alberto Bietti and Francis Bach. Deep equals shallow for relu networks in kernel regimes, 2021. ICLR 2021.
- [18] Phan-Minh Nguyen and Huy Tuan Pham. A rigorous framework for the mean field limit of multilayer neural networks, 2023.
- [19] Jeremy M. Cohen, Alex Damian, Ameet Talwalkar, J. Zico Kolter, and Jason D. Lee. Understanding optimization in deep learning with central flows, 2025.
- [20] Sanjeev Arora, Simon S Du, Wei Hu, Zhiyuan Li, and Ruosong Wang. On exact computation with an infinitely wide neural net. *Advances in Neural Information Processing Systems*, 32, 2019.
- [21] Sanjeev Arora, Simon S Du, Zhiyuan Li, Ruslan Salakhutdinov, Ruosong Wang, and Dingli Yu. On the inductive bias of neural tangent kernels. In *Advances in Neural Information Processing Systems*, pages 4825–4836, 2019.
- [22] Greg Yang. Scaling limits of wide neural networks with weight sharing: Gaussian process behavior, gradient independence, and neural tangent kernel derivation. *arXiv preprint arXiv:1902.04760*, 2019.
- [23] Lucas Benigni and Elliot Paquette. Eigenvalue distribution of the neural tangent kernel in the quadratic scaling, 2025.
- [24] Ali Rahimi and Benjamin Recht. Random features for large-scale kernel machines. In *Advances in Neural Information Processing Systems*, pages 1177–1184, 2007.
- [25] Ali Rahimi and Benjamin Recht. Weighted sums of random kitchen sinks: Replacing minimization with randomization in learning. In *Advances in Neural Information Processing Systems*, pages 1313–1320, 2008.
- [26] Francis Bach. Breaking the curse of dimensionality with convex neural networks. *Journal of Machine Learning Research*, 18(19):1–53, 2017.
- [27] Alessandro Rudi and Lorenzo Rosasco. Generalization properties of learning with random features. *Advances in Neural Information Processing Systems*, 30, 2017.
- [28] Bharath K Sriperumbudur, Kenji Fukumizu, Arthur Gretton, Bernhard Schölkopf, and Gert RG Lanckriet. On the optimal estimation of probability measures in weak and strong topologies. *Bernoulli*, 21(3):1759–1786, 2015.
- [29] Tara N Sainath, Brian Kingsbury, Vikas Sindhwani, Ebru Arisoy, and Bhuvana Ramabhadran. Low-rank matrix factorization for deep neural network training with high-dimensional output targets. *IEEE International Conference on Acoustics, Speech and Signal Processing*, pages 6655–6659, 2013.

- [30] Alexander Novikov, Dmitry Podoprikin, Anton Osokin, and Dmitry Vetrov. Tensorizing neural networks. In *Advances in Neural Information Processing Systems*, pages 442–450, 2015.
- [31] Sanjeev Arora, Nadav Cohen, and Elad Hazan. On the optimization of deep networks: Implicit acceleration by overparameterization. *International Conference on Machine Learning*, pages 244–253, 2018.
- [32] Nasim Rahaman, Aristide Baratin, Devansh Arpit, Felix Draxler, Min Lin, Fred A Hamprecht, Yoshua Bengio, and Aaron Courville. On the spectral bias of neural networks. In *International Conference on Machine Learning*, pages 5301–5310, 2019.
- [33] Zhi-Qin John Xu, Yaoyu Zhang, Tao Luo, Yanyang Xiao, and Zheng Ma. Training behavior of deep neural network in frequency domain. *International Conference on Neural Information Processing*, pages 264–274, 2019.
- [34] Ronen Basri, Meirav Galun, Amnon Geifman, David Jacobs, Yoni Kasten, and Shira Kritchman. On the frequency bias of generative models. *Advances in Neural Information Processing Systems*, 33:18126–18136, 2020.
- [35] Matthew Tancik, Pratul P Srinivasan, Ben Mildenhall, Sara Fridovich-Keil, Nithin Raghavan, Utkarsh Singhal, Ravi Ramamoorthi, Jonathan T Barron, and Ren Ng. Fourier features let networks learn high frequency functions in low dimensional domains. In *Advances in Neural Information Processing Systems*, volume 33, pages 7537–7547, 2020.
- [36] Gérard Ben Arous, Reza Gheissari, and Aukosh Jagannath. Online stochastic gradient descent on non-convex losses from high-dimensional inference. *Journal of Machine Learning Research*, 22(106):1–51, 2021.
- [37] Alex Damian, Jason Lee, and Mahdi Soltanolkotabi. Neural networks can learn representations with gradient descent. *Conference on Learning Theory*, pages 5413–5452, 2022.
- [38] Emmanuel Abbe, Enric Boix-Adsera, and Theodor Misiakiewicz. The merged-staircase property: a necessary and nearly sufficient condition for sgd learning of sparse functions on two-layer neural networks. *Conference on Learning Theory*, pages 4782–4887, 2023.
- [39] Jeremy Cohen, Simran Kaur, Yuanzhi Li, J Zico Kolter, and Ameet Talwalkar. Gradient descent on neural networks typically occurs at the edge of stability. In *International Conference on Learning Representations*, 2021.
- [40] Ashia C Wilson, Rebecca Roelofs, Mitchell Stern, Nati Srebro, and Benjamin Recht. The marginal value of adaptive gradient methods in machine learning. *Advances in Neural Information Processing Systems*, 30, 2017.
- [41] Jared Kaplan, Sam McCandlish, Tom Henighan, Tom B Brown, Benjamin Chess, Rewon Child, Scott Gray, Alec Radford, Jeffrey Wu, and Dario Amodei. Scaling laws for neural language models. *arXiv preprint arXiv:2001.08361*, 2020.
- [42] Jordan Hoffmann, Sebastian Borgeaud, Arthur Mensch, Elena Buchatskaya, Trevor Cai, Eliza Rutherford, Diego de Las Casas, Lisa Anne Hendricks, Johannes Welbl, Aidan Clark, et al. Training compute-optimal large language models. *arXiv preprint arXiv:2203.15556*, 2022.
- [43] Jason Wei, Yi Tay, Rishi Bommasani, Colin Raffel, Barret Zoph, Sebastian Borgeaud, Dani Yogatama, Maarten Bosma, Denny Zhou, Donald Metzler, et al. Emergent abilities of large language models. *arXiv preprint arXiv:2206.07682*, 2022.
- [44] George Cybenko. Approximation by superpositions of a sigmoidal function. *Mathematics of Control, Signals and Systems*, 2(4):303–314, 1989.
- [45] Kurt Hornik, Maxwell Stinchcombe, and Halbert White. Multilayer feedforward networks are universal approximators. *Neural Networks*, 2(5):359–366, 1989.
- [46] Lenaïc Chizat and Francis Bach. Implicit bias of gradient descent for wide two-layer neural networks trained with the logistic loss. *Conference on Learning Theory*, pages 1305–1338, 2020.
- [47] Quynh Nguyen, Marco Mondelli, and Guido Montufar. Tight bounds on the smallest eigenvalue of the neural tangent kernel for deep relu networks, 2022.
- [48] Soufiane Hayou, Arnaud Doucet, and Judith Rousseau. On the selection of initialization and activation function for deep neural networks, 2018.
- [49] W. F. Kibble. A two-variate gamma type distribution. *Sankhyā: The Indian Journal of Statistics*, 5:137–150, 1941.
- [50] Weinan E, Chao Ma, and Lei Wu. The barron space and the flow-induced function spaces for neural network models, 2021.
- [51] F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain. Nist digital library of mathematical functions. <https://dlmf.nist.gov/>, 2024. Release 1.2.1 of 2024-06-15.

- [52] Milton Abramowitz and Irene A. Stegun, editors. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. Dover Publications, New York, 1964.
- [53] Nouredine El Karoui. The spectrum of kernel random matrices. *The Annals of Statistics*, 38(1), feb 2010.
- [54] Greg Yang. Tensor programs ii: Neural tangent kernel for any architecture, 2020.

8 Proofs of NTK Recursions

8.1 Proof of Theorem 3.1: Infinite-width NTK recursion

Proof. We prove the recursion by decomposing the NTK at layer ℓ into contributions of the trainable parameters up to layer ℓ , and invoking the Law of Large Numbers (LLN) under the sequential infinite-width limit. Let $f^{(\ell)}$ denote the network output truncated at layer ℓ . For finite width, the (random) NTK reads

$$\Theta_{\text{rand}}^{(\ell)}(x, x') = \underbrace{(\nabla_{\theta^{(\ell-1)}} f^{(\ell)}(x))^{\top} (\nabla_{\theta^{(\ell-1)}} f^{(\ell)}(x'))}_{\text{Term 1}} + \underbrace{\sum_{j=1}^{n_{\ell}} \frac{\partial f^{(\ell)}(x)}{\partial A_j^{(\ell)}} \frac{\partial f^{(\ell)}(x')}{\partial A_j^{(\ell)}}}_{\text{Term 2}}. \quad (8)$$

Bias parameters. Under the NTK parameterization with trainable biases at each layer, the gradient with respect to the bias contributes an identity kernel. Concretely, for layer ℓ , the bias partial derivative satisfies $\partial f^{(\ell)}(x)/\partial b^{(\ell)} = \sigma_A/\sqrt{n_{\ell}}$, so the associated NTK contribution converges by LLN to a deterministic constant, which we normalize as 1 under our parameterization. This yields the additive +1 term in the recursion.

Term 2 (layer- ℓ output weights $A^{(\ell)}$). By the forward pass, $\partial f^{(\ell)}(x)/\partial A_j^{(\ell)} = \frac{\sigma_A}{\sqrt{n_{\ell}}} \sigma(w_j^{(\ell)\top} \mathbf{h}^{(\ell-1)}(x) + \beta b_j^{(\ell)})$. Therefore

$$\text{Term 2} = \frac{\sigma_A^2}{n_{\ell}} \sum_{j=1}^{n_{\ell}} \sigma(w_j^{(\ell)\top} \mathbf{h}^{(\ell-1)}(x) + \beta b_j^{(\ell)}) \sigma(w_j^{(\ell)\top} \mathbf{h}^{(\ell-1)}(x') + \beta b_j^{(\ell)}). \quad (9)$$

Conditionally on $\mathbf{h}^{(\ell-1)}$, the summands are i.i.d. over $(w_j^{(\ell)}, b_j^{(\ell)})$. As $n_{\ell} \rightarrow \infty$ (with lower layers already taken to their limits), LLN yields

$$\text{Term 2} \xrightarrow{P} \sigma_A^2 \Sigma^{(\ell)}(x, x'). \quad (10)$$

Term 1 (previous layers). By the chain rule,

$$\nabla_{\theta^{(\ell-1)}} f^{(\ell)}(x) = \left(\frac{\partial f^{(\ell)}(x)}{\partial \mathbf{h}^{(\ell-1)}(x)} \right) \nabla_{\theta^{(\ell-1)}} \mathbf{h}^{(\ell-1)}(x). \quad (11)$$

Hence Term 1 factorizes as $\Theta_{\text{rand}}^{(\ell-1)}(x, x')$ multiplied by a Jacobian metric:

$$\text{Term 1} = \Theta_{\text{rand}}^{(\ell-1)}(x, x') \cdot \left\langle \frac{\partial f^{(\ell)}}{\partial \mathbf{h}^{(\ell-1)}}(x), \frac{\partial f^{(\ell)}}{\partial \mathbf{h}^{(\ell-1)}}(x') \right\rangle. \quad (12)$$

Taking the inner product and applying LLN as $n_{\ell} \rightarrow \infty$ yields

$$\left\langle \frac{\partial f^{(\ell)}}{\partial \mathbf{h}^{(\ell-1)}}(x), \frac{\partial f^{(\ell)}}{\partial \mathbf{h}^{(\ell-1)}}(x') \right\rangle \xrightarrow{P} \sigma_A^2 \dot{\Sigma}^{(\ell)}(x, x'). \quad (13)$$

Therefore,

$$\text{Term 1} \xrightarrow{P} \Theta^{(\ell-1)}(x, x') \cdot \sigma_A^2 \dot{\Sigma}^{(\ell)}(x, x'). \quad (14)$$

Combining the bias contribution with the two limits completes the recursion

$$\Theta^{(\ell)}(x, x') = 1 + \Theta^{(\ell-1)}(x, x') \cdot \sigma_A^2 \dot{\Sigma}^{(\ell)}(x, x') + \sigma_A^2 \Sigma^{(\ell)}(x, x'). \quad (15)$$

The sequential limit assumption ensures that lower-layer random objects have already converged when taking the next-layer limit, justifying the conditional LLN applications. $\square$

8.2 Proof of Theorem 2.1: Gaussian process composition

Proof. We prove by induction on ℓ that $\mathbf{h}^{(\ell)}$ converges to a Gaussian process with covariance kernel $\Sigma^{(\ell)}$.

Base case $\ell = 0$. For $\ell = 0$, we have $\mathbf{h}^{(0)}(x) = x$, which is deterministic and trivially a Gaussian process.

Inductive step. Assume that $\mathbf{h}^{(\ell-1)}$ converges to a Gaussian process with covariance kernel $\Sigma^{(\ell-1)}$. Consider the forward pass at layer ℓ :

$$\mathbf{h}^{(\ell)}(x) = \frac{1}{\sqrt{n_\ell}} \sum_{j=1}^{n_\ell} A_j^{(\ell)} \sigma\left(w_j^{(\ell)\top} \mathbf{h}^{(\ell-1)}(x) + b_j^{(\ell)}\right) + c. \quad (16)$$

For any finite collection of inputs $\{x_1, \dots, x_m\}$, conditionally on $\mathbf{h}^{(\ell-1)}$, the terms $A_j^{(\ell)} \sigma\left(w_j^{(\ell)\top} \mathbf{h}^{(\ell-1)}(x_i) + b_j^{(\ell)}\right)$ are independent across j and identically distributed. Since $A_j^{(\ell)} \sim \mathcal{N}(0, \sigma_A^2)$ and $w_j^{(\ell)}, b_j^{(\ell)}$ are frozen Gaussian draws, the conditional distribution of each term is centered with finite variance.

By the multivariate Central Limit Theorem, as $n_\ell \rightarrow \infty$, the vector $(\mathbf{h}^{(\ell)}(x_1), \dots, \mathbf{h}^{(\ell)}(x_m))$ converges in distribution to a multivariate Gaussian. The covariance is given by

$$\mathbb{E}\left[\mathbf{h}^{(\ell)}(x_i) \mathbf{h}^{(\ell)}(x_j)\right] = \mathbb{E}_w\left[\sigma\left(w^\top \mathbf{h}^{(\ell-1)}(x_i)\right) \sigma\left(w^\top \mathbf{h}^{(\ell-1)}(x_j)\right)\right] = \Sigma^{(\ell)}(x_i, x_j), \quad (17)$$

where the expectation is taken over the frozen weights w and the limiting Gaussian process $\mathbf{h}^{(\ell-1)}$. The last equality follows from the definition of the base kernel in Definition 2.1.

The sequential infinite-width limit ensures that $\mathbf{h}^{(\ell-1)}$ has already converged to its Gaussian process limit when taking the limit for layer ℓ , justifying the application of the CLT conditionally on $\mathbf{h}^{(\ell-1)}$. $\square$

8.3 Proof of three-layer NTK Corollary

Proof. Apply Theorem 3.1 for $\ell = 1$ and $\ell = 2$. For $\ell = 1$, since $\Theta^{(0)} = 0$, the recursion gives:

$$\Theta^{(1)}(x, x') = 1 + \Theta^{(0)}(x, x') \cdot \dot{\Sigma}^{(1)}(x, x') + \Sigma^{(1)}(x, x') = 1 + \Sigma^{(1)}(x, x'). \quad (18)$$

For $\ell = 2$, substitute $\Theta^{(1)} = 1 + \Sigma^{(1)}$ into the recursion:

$$\Theta^{(2)}(x, x') = 1 + \Theta^{(1)}(x, x') \cdot \dot{\Sigma}^{(2)}(x, x') + \Sigma^{(2)}(x, x'). \quad (19)$$

This is the stated three-layer recursion with the bias contribution.

Mean explicit form. Under the homogeneous ReLU setting on the sphere, the last-layer expectations are

$$\mathbb{E}[\dot{\Sigma}^{(2)}(\rho)] = 1 - \frac{\arccos(\rho)}{\pi}, \quad \mathbb{E}[\Sigma^{(2)}(\rho)] = \Sigma^{(1)}(\rho). \quad (20)$$

Taking the conditional mean at layer 2 yields the mean three-layer NTK as a function of ρ :

$$\tilde{\Theta}^{(2)}(\rho) = 1 + \Theta^{(1)}(\rho) \left(1 - \frac{\arccos(\rho)}{\pi}\right) + \Sigma^{(1)}(\rho). \quad (21)$$

Since $\Theta^{(1)}(\rho) = 1 + \Sigma^{(1)}(\rho)$, this can be written as

$$\tilde{\Theta}^{(2)}(\rho) = \underbrace{\Theta^{(1)}(\rho)}_{\Theta^{(1)}(\rho)} \left(1 - \frac{\arccos(\rho)}{\pi}\right) + \Sigma^{(1)}(\rho) + 1 = K_\infty(\rho), \quad (22)$$

which proves the explicit expression. $\square$

8.4 Proof of General L -layer NTK Corollary

Proof. We prove the explicit form by induction on L , accounting for the +1 bias terms in the recursion.

Base case $L = 1$. From Theorem 3.1, $\Theta^{(1)} = 1 + \Sigma^{(1)}$. The formula with $L = 1$ gives:

$$\Theta^{(1)}(x, x') = \sum_{\ell=1}^1 \left(\prod_{k=\ell+1}^1 \dot{\Sigma}^{(k)}(x, x') \right) + \sum_{\ell=1}^1 \Sigma^{(\ell)}(x, x') \prod_{k=\ell+1}^1 \dot{\Sigma}^{(k)}(x, x'). \quad (23)$$

For the first sum: when $\ell = 1$, $\prod_{k=2}^1 \dot{\Sigma}^{(k)} = 1$ (empty product). For the second sum: when $\ell = 1$, $\Sigma^{(1)} \cdot \prod_{k=2}^1 \dot{\Sigma}^{(k)} = \Sigma^{(1)} \cdot 1 = \Sigma^{(1)}$. Therefore, $\Theta^{(1)} = 1 + \Sigma^{(1)}$, which matches the recursion.

Inductive step. Assume the formula holds for depth $L - 1$:

$$\Theta^{(L-1)}(x, x') = \sum_{\ell=1}^{L-1} \left(\prod_{k=\ell+1}^{L-1} \dot{\Sigma}^{(k)}(x, x') \right) + \sum_{\ell=1}^{L-1} \Sigma^{(\ell)}(x, x') \prod_{k=\ell+1}^{L-1} \dot{\Sigma}^{(k)}(x, x'). \quad (24)$$

By Theorem 3.1,

$$\Theta^{(L)}(x, x') = 1 + \Theta^{(L-1)}(x, x') \cdot \dot{\Sigma}^{(L)}(x, x') + \Sigma^{(L)}(x, x'). \quad (25)$$

Substituting the inductive hypothesis:

$$\begin{aligned} \Theta^{(L)}(x, x') &= 1 + \left[\sum_{\ell=1}^{L-1} \left(\prod_{k=\ell+1}^{L-1} \dot{\Sigma}^{(k)}(x, x') \right) + \sum_{\ell=1}^{L-1} \Sigma^{(\ell)}(x, x') \prod_{k=\ell+1}^{L-1} \dot{\Sigma}^{(k)}(x, x') \right] \cdot \dot{\Sigma}^{(L)}(x, x') + \Sigma^{(L)}(x, x') \\ &= 1 + \sum_{\ell=1}^{L-1} \left(\prod_{k=\ell+1}^L \dot{\Sigma}^{(k)}(x, x') \right) + \sum_{\ell=1}^{L-1} \Sigma^{(\ell)}(x, x') \prod_{k=\ell+1}^L \dot{\Sigma}^{(k)}(x, x') + \Sigma^{(L)}(x, x'). \end{aligned}$$

The first term 1 corresponds to $\ell = L$ in the first sum (with empty product $\prod_{k=L+1}^L \dot{\Sigma}^{(k)} = 1$). The last term $\Sigma^{(L)}$ corresponds to $\ell = L$ in the second sum (with empty product $\prod_{k=L+1}^L \dot{\Sigma}^{(k)} = 1$). Therefore:

$$\Theta^{(L)}(x, x') = \sum_{\ell=1}^L \left(\prod_{k=\ell+1}^L \dot{\Sigma}^{(k)}(x, x') \right) + \sum_{\ell=1}^L \Sigma^{(\ell)}(x, x') \prod_{k=\ell+1}^L \dot{\Sigma}^{(k)}(x, x'), \quad (26)$$

completing the induction. The total number of terms is $L + L = 2L$, which grows linearly with depth. $\square$

9 Proofs of low-rank NTK RKHS

9.1 Discussion on Fisher and Kibble Distributions

Lemma 9.1 (Fisher–Kibble decoupling). *Let x, y be input vectors and let x_1, y_1 denote their rank- r random projections. Define the empirical correlation $\rho_1 = \cos \angle(x_1, y_1)$ and squared norms $u = \|x_1\|^2, v = \|y_1\|^2$. Then:*

- ρ_1 follows Fisher’s correlation distribution [15] [16], centered at the true correlation ρ .
- (u, v) follow Kibble’s [49] bivariate Gamma (chi-square) law.
- Angular and radial parts are independent: $p(\rho_1, u, v) = p_{\text{Fisher}}(\rho_1) p_{\text{Kibble}}(u, v)$.

Proof. Let $x, y \in \mathbb{R}^d$ be fixed input vectors with correlation $\rho = \langle x, y \rangle / (\|x\| \|y\|)$. Consider a rank- r random projection matrix $P \in \mathbb{R}^{r \times d}$ with entries $P_{ij} \sim \mathcal{N}(0, 1/d)$ i.i.d., so that $x_1 = Px$ and $y_1 = Py$ are the projected vectors.

Since P has i.i.d. Gaussian entries, the projected vectors $x_1, y_1 \in \mathbb{R}^r$ form a ρ -correlated bivariate Gaussian pair. Specifically, for each component $i = 1, \dots, r$, the pairs (x_{1i}, y_{1i}) are i.i.d. with joint distribution

$$\begin{pmatrix} x_{1i} \\ y_{1i} \end{pmatrix} \sim \mathcal{N} \left(\mathbf{0}, \begin{pmatrix} \|x\|^2/d & \rho \|x\| \|y\|/d \\ \rho \|x\| \|y\|/d & \|y\|^2/d \end{pmatrix} \right). \quad (27)$$

By rotational invariance of the Gaussian projection and the fact that the correlation ρ is preserved under orthogonal transformations, we can assume without loss of generality that the covariance structure factors into angular and radial components.

Angular part (Fisher distribution). The empirical correlation is defined as

$$\rho_1 = \frac{\langle x_1, y_1 \rangle}{\|x_1\| \|y_1\|} = \frac{\sum_{i=1}^r x_{1i} y_{1i}}{\sqrt{\sum_{i=1}^r x_{1i}^2} \sqrt{\sum_{i=1}^r y_{1i}^2}}. \quad (28)$$

This is the sample correlation coefficient of r pairs of correlated Gaussian random variables. Fisher [15, 16] showed that the distribution of ρ_1 depends only on the true correlation ρ and the sample size r , with density given in Theorem 3.1. The key observation is that ρ_1 is a function purely of the angles between the projected vectors, independent of their magnitudes.

Radial part (Kibble distribution). The squared norms $u = \|x_1\|^2 = \sum_{i=1}^r x_{1i}^2$ and $v = \|y_1\|^2 = \sum_{i=1}^r y_{1i}^2$ are sums of correlated chi-square variables. Since (x_{1i}, y_{1i}) are ρ -correlated Gaussians, the bivariate distribution of (u, v) follows Kibble's [49] bivariate chi-square law. The joint density involves the modified Bessel function $I_\nu(z)$, which appears because the dot product $\langle x_1, y_1 \rangle = \sum_{i=1}^r x_{1i} y_{1i}$ can be expressed as a weighted sum of products of correlated normals, whose distribution relates to Bessel functions through the generating function of the bivariate chi-square distribution.

Specifically, the joint characteristic function of (u, v) is

$$\phi(s, t) = \mathbb{E}[e^{i(su+tv)}] = \left(1 - 2i(1 - \rho^2)(s + t) - 4\rho^2 st\right)^{-r/2}, \quad (29)$$

and the inverse Fourier transform yields Kibble's density, which contains the modified Bessel function $I_{\frac{r}{2}-1}$ as shown in Lemma 3.1. The Bessel function arises from the integral representation of the bivariate chi-square density when the correlation $\rho \neq 0$.

Independence of angular and radial parts. Rotational invariance of isotropic Gaussian projections implies that $\rho_1 = \cos \angle(x_1, y_1)$ and the norms $(\|x_1\|, \|y_1\|)$ are independent, yielding the factorization $p(\rho_1, u, v) = p_{\text{Fisher}}(\rho_1) \cdot p_{\text{Kibble}}(u, v)$. This is the classical independence of the sample correlation from sample variances in bivariate normal data. $\square$

9.2 Proof of Theorem 4.1: Rank-driven concentration

Proof. Let $x_1, y_1 \in \mathbb{R}^r$ be Gaussian projections with arbitrary fixed correlation $\rho \in [-1, 1]$. Define $U = \|x_1\|/\sqrt{r}$, $V = \|y_1\|/\sqrt{r}$, and $W = UV = \|x_1\| \|y_1\|/r$. We prove that for $\epsilon \in (0, 1)$,

$$\mathbb{P}(|W - 1| \geq \epsilon) \leq 4 \exp\left(-\frac{r\epsilon^2}{8}\right), \quad (30)$$

and that for $\epsilon \geq 1$ there exist constants $c_1, c_2 > 0$ with $\mathbb{P}(|W - 1| \geq \epsilon) \leq c_1 e^{-c_2 r \epsilon}$.

The Euclidean norm is 1-Lipschitz, so by Gaussian concentration, for all $\delta > 0$,

$$\mathbb{P}(|U - 1| \geq \delta) \leq 2 \exp\left(-\frac{r\delta^2}{2}\right), \quad \mathbb{P}(|V - 1| \geq \delta) \leq 2 \exp\left(-\frac{r\delta^2}{2}\right). \quad (31)$$

We use

$$|UV - 1| \leq |U - 1| + |V - 1| + |U - 1| |V - 1|. \quad (32)$$

Fix $\epsilon \in (0, 1)$ and set $\delta = \epsilon/2$. On the event $\{|U - 1| < \delta, |V - 1| < \delta\}$, we have

$$|UV - 1| \leq 2\delta + \delta^2 \leq \epsilon. \quad (33)$$

Therefore,

$$\mathbb{P}(|UV - 1| \geq \epsilon) \leq \mathbb{P}(|U - 1| \geq \delta) + \mathbb{P}(|V - 1| \geq \delta) \leq 4 \exp\left(-\frac{r\epsilon^2}{8}\right). \quad (34)$$

$\square$

9.3 Proof of Corollary 4.1: Concentration bound for three-layer NTK

Proof. We prove that the empirical three-layer NTK $\hat{\Theta}_r^{(2)}(x, x') = \Psi(\hat{\rho}_r, \hat{w}_r)$ concentrates exponentially fast around its deterministic limit $K_\infty(\rho) = \Psi(\rho, 1)$ as $r \rightarrow \infty$.

Recall that the kernel function is defined as:

$$\Psi(u, w) = \Theta^{(1)}(u) \left(1 - \frac{\arccos(u)}{\pi}\right) + w \Sigma^{(1)}(u) + 1, \quad (35)$$

where $\hat{\rho}_r$ is the sample correlation and $\hat{w}_r = \|x_1\| \|y_1\|/r$. By Fisher's CLT [15, 16], $\sqrt{r}(\hat{\rho}_r - \rho) \xrightarrow{d} \mathcal{N}(0, (1 - \rho^2)^2)$, and by Theorem 4.1, $\hat{w}_r$ concentrates around 1 with sub-Gaussian tails.

The function Ψ is differentiable in both arguments. For fixed $\rho \in (-1, 1)$, we have:

$$\partial_u \Psi(\rho, 1) = \partial_\rho K_\infty(\rho) = O(1), \quad (36)$$

$$\partial_w \Psi(\rho, 1) = \Sigma^{(1)}(\rho) = O(1). \quad (37)$$

Since $\Theta^{(1)}$, $\Sigma^{(1)}$, and $\arccos$ are smooth on $(-1, 1)$, there exist Lipschitz constants $L_u, L_w > 0$ (depending on ρ) such that for $u, u' \in [\rho - \delta, \rho + \delta]$ and $w, w' \in [1 - \delta, 1 + \delta]$ with $\delta > 0$ small:

$$|\Psi(u, w) - \Psi(u', w')| \leq L_u |u - u'| + L_w |w - w'|. \quad (38)$$

We combine radial and angular concentration by using a union bound and the Lipschitz property:

$$|\hat{\Theta}_r^{(2)}(x, x') - K_\infty(\rho)| = |\Psi(\hat{\rho}_r, \hat{w}_r) - \Psi(\rho, 1)| \leq L_u |\hat{\rho}_r - \rho| + L_w |\hat{w}_r - 1|. \quad (39)$$

Therefore, for $\epsilon \in (0, 1)$:

$$\mathbb{P}\left(|\hat{\Theta}_r^{(2)}(x, x') - K_\infty(\rho)| \geq \epsilon\right) \leq \mathbb{P}\left(|\hat{\rho}_r - \rho| \geq \frac{\epsilon}{2L_u}\right) + \mathbb{P}\left(|\hat{w}_r - 1| \geq \frac{\epsilon}{2L_w}\right). \quad (40)$$

Now we prove the sub-Gaussian concentration of the sample correlation via Hanson–Wright.

Lemma 9.2 (Sub-Gaussian concentration of the sample correlation via Hanson–Wright). *Let $(X_i, Y_i)_{i=1}^r$ be i.i.d. centered Gaussian pairs with unit variances and correlation $\rho \in (-1, 1)$. Define*

$$S_{xx} = \frac{1}{r} \sum_{i=1}^r X_i^2, \quad S_{yy} = \frac{1}{r} \sum_{i=1}^r Y_i^2, \quad S_{xy} = \frac{1}{r} \sum_{i=1}^r X_i Y_i, \quad \hat{\rho}_r = \frac{S_{xy}}{\sqrt{S_{xx} S_{yy}}}. \quad (41)$$

There exist absolute constants $c, C > 0$ such that for all $t \in (0, 1)$,

$$\mathbb{P}(|\hat{\rho}_r - \rho| \geq t) \leq C \exp(-c r t^2). \quad (42)$$

Proof. Write $Y = \rho X + \sqrt{1 - \rho^2} Z$ with $X = (X_i)_{i \leq r}$ and $Z = (Z_i)_{i \leq r}$ independent, i.i.d. $\mathcal{N}(0, 1)$. Then

$$S_{xy} - \rho = \rho (S_{xx} - 1) + \underbrace{\sqrt{1 - \rho^2} \frac{1}{r} \sum_{i=1}^r X_i Z_i}_{=: T_r}. \quad (43)$$

From [?] Theorem 6.2.2 page 183, the bilinear Hanson–Wright inequality (for independent sub-Gaussian X, Z and fixed $M \in \mathbb{R}^{r \times r}$) states

$$\mathbb{P}(|X^\top M Z| \geq u) \leq 2 \exp \left[-c \min \left(\frac{u^2}{K^4 \|M\|_F^2}, \frac{u}{K^2 \|M\|} \right) \right], \quad (44)$$

where $K = O(1)$ is the Orlicz sub-gaussian norm of the coordinates. For $M = I_r$ we have $\|M\|_F^2 = r$ and $\|M\| = 1$, hence with $u = r t$ and $t \in (0, 1)$,

$$\mathbb{P}\left(\left|\frac{1}{r} X^\top Z\right| \geq t\right) \leq 2 \exp(-c r t^2). \quad (45)$$

Thus T_r is sub-Gaussian at rate $\exp(-c r t^2)$:

$$\mathbb{P}(|T_r| \geq t) \leq 2 \exp(-c r t^2). \quad (46)$$

For $S_{xx} = (1/r) \|X\|^2$ and $S_{yy} = (1/r) \|Y\|^2$, Laurent–Massart Gaussian quadratic chaos inequality in (4.1) page 1325 [?] yields, for any $s \in (0, 1)$,

$$\mathbb{P}(|S_{xx} - 1| \geq s) \leq 2 \exp(-c r s^2), \quad \mathbb{P}(|S_{yy} - 1| \geq s) \leq 2 \exp(-c r s^2). \quad (47)$$

Let $f(a, b, c) = a/\sqrt{bc}$. A first-order expansion at $(\rho, 1, 1)$ yields

$$|\hat{\rho}_r - \rho| = |f(S_{xy}, S_{xx}, S_{yy}) - f(\rho, 1, 1)| \leq |S_{xy} - \rho| + \frac{|\rho|}{2} (|S_{xx} - 1| + |S_{yy} - 1|) + R, \quad (48)$$

where the remainder $R = O((|S_{xx} - 1| + |S_{yy} - 1|)^2)$ is negligible on the event $\{|S_{xx} - 1| \vee |S_{yy} - 1| \leq c_0\}$ (for some absolute c_0 , e.g., $1/4$). Choosing $s = \kappa t$ for a sufficiently small absolute $\kappa > 0$, and splitting $|S_{xy} - \rho|$ as in step 1, a union bound gives

$$\mathbb{P}(|\hat{\rho}_r - \rho| \geq t) \leq \underbrace{\mathbb{P}(|T_r| \geq c_1 t)}_{\leq 2e^{-crt^2}} + \underbrace{\mathbb{P}(|S_{xx} - 1| \geq c_2 t)}_{\leq 2e^{-crt^2}} + \underbrace{\mathbb{P}(|S_{yy} - 1| \geq c_2 t)}_{\leq 2e^{-crt^2}} + \mathbb{P}(R \not\ll t). \quad (49)$$

The last term is absorbed by adjusting κ (small-probability event controlled by the same inequalities). Therefore, for absolute constants $c, C > 0$,

$$\mathbb{P}(|\hat{\rho}_r - \rho| \geq t) \leq C \exp(-c r t^2), \quad t \in (0, 1). \quad (50)$$

□

Remark 9.1 (“Bessel” variant: sum of Gaussian products). Each product $X_i Z_i$ has a symmetric sub-exponential tail (density involving the Bessel function K_0). By Bernstein’s inequality for i.i.d. sub-exponential variables, $\frac{1}{r} \sum_{i=1}^r X_i Z_i$ satisfies the same $\exp(-crt^2)$ bound for $t \in (0, 1)$, yielding an alternative proof of the bilinear step.

By Theorem 4.1, for $\epsilon/(2L_w) \in (0, 1)$:

$$\mathbb{P}\left(|\hat{w}_r - 1| \geq \frac{\epsilon}{2L_w}\right) \leq 4 \exp\left(-\frac{r\epsilon^2}{32L_w^2}\right). \quad (51)$$

Combining both terms:

$$\mathbb{P}\left(|\hat{\Theta}_r^{(2)}(x, x') - K_\infty(\rho)| \geq \epsilon\right) \leq \underbrace{\mathbb{P}\left(|\hat{\rho}_r - \rho| \geq \frac{\epsilon}{2L_u}\right)}_{\leq C \exp\left(-\frac{c r \epsilon^2}{4L_u^2}\right)} + \underbrace{\mathbb{P}\left(|\hat{w}_r - 1| \geq \frac{\epsilon}{2L_w}\right)}_{\leq 4 \exp\left(-\frac{r \epsilon^2}{32L_w^2}\right)} \quad (52)$$

so that

$$\mathbb{P}\left(|\hat{\Theta}_r^{(2)}(x, x') - K_\infty(\rho)| \geq \epsilon\right) \leq C \exp\left(-\frac{c r \epsilon^2}{4L_u^2}\right) + 4 \exp\left(-\frac{r \epsilon^2}{32L_w^2}\right). \quad (53)$$

Taking $C_1 = C + 4$ and $C_2 = \min(c/(4L_u^2), 1/(32L_w^2))$, we obtain

$$\mathbb{P}\left(|\hat{\Theta}_r^{(2)}(x, x') - K_\infty(\rho)| \geq \epsilon\right) \leq C_1 \exp(-C_2 r \epsilon^2). \quad (54)$$

□

9.4 Proof of Corollary 4.2: Mean NTK over Fisher–Kibble has same RKHS

Proof. We prove that under Assumption 2.1, the mean NTK $\tilde{\Theta}^{(L)}$ (obtained by taking expectations over Fisher and Kibble distributions) is a zonal kernel that induces the same RKHS as the shallow (three-layer) ReLU kernel.

Puiseux expansion of mean NTK near $\rho = 1$: The key observation is that the mean NTK $\tilde{\Theta}^{(2)}(\rho)$ is a zonal kernel (depends only on ρ). Near $\rho = 1$ (for $t \geq 0$), we need to compute the Puiseux expansion of $\mathbb{E}[\arccos(\hat{\rho}_r)]$ when $\rho = 1 - t$, where $\hat{\rho}_r \sim \text{Fisher}(1 - t, r)$.

This is highly non-trivial because we must compute:

$$\mathbb{E}[\arccos(\hat{\rho}_r)] = \int_{-1}^1 \arccos(u) p_{\text{Fisher}}(u \mid \rho = 1 - t, r) du, \quad (55)$$

where the full Fisher distribution density is:

$$p_{\text{Fisher}}(u \mid \rho, r) = \frac{(r-2) \Gamma(r-1) (1-\rho^2)^{\frac{r-1}{2}} (1-u^2)^{\frac{r-4}{2}}}{\sqrt{2\pi} \Gamma(r-\frac{1}{2}) (1-\rho u)^{r-\frac{3}{2}}} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; r-\frac{1}{2}; \frac{1+\rho u}{2}\right), \quad (56)$$

for $r > 2$, with ${}_2F_1$ the hypergeometric function.

Making the change of variables $v = 1 - u$ and $s = 1 - t$, so $\rho = s = 1 - t$ and $u = 1 - v$, we have:

$$\mathbb{E}[\arccos(\hat{\rho}_r)] = \int_0^2 \arccos(1 - v) p_{\text{Fisher}}(1 - v \mid s, r) dv. \quad (57)$$

Near $v = 0$ and $t = 0$, we analyze the asymptotic behavior of the Fisher density. For $\rho = s = 1 - t$ and $u = 1 - v$ with $t, v \rightarrow 0^+$:

$$1 - \rho^2 = 1 - (1 - t)^2 = 2t - t^2 = 2t(1 - t/2), \quad (58)$$

$$1 - u^2 = 1 - (1 - v)^2 = 2v - v^2 = 2v(1 - v/2), \quad (59)$$

$$1 - \rho u = 1 - (1 - t)(1 - v) = t + v - tv = t + v + O(tv). \quad (60)$$

The Fisher density near $v = 0$ and $t = 0$ behaves as:

$$p_{\text{Fisher}}(1 - v \mid 1 - t, r) \sim \frac{(r - 2) \Gamma(r - 1) (2t)^{\frac{r-1}{2}} (2v)^{\frac{r-4}{2}}}{\sqrt{2\pi} \Gamma(r - \frac{1}{2}) (t + v)^{r - \frac{3}{2}}} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; r - \frac{1}{2}; 1 - \frac{t+v}{2}\right). \quad (61)$$

For the hypergeometric function near its argument $1 - (t + v)/2 \rightarrow 1^-$, we invoke the connection formula in DLMF §15.8.2 [51] together with the fact that $c - a - b = r - \frac{3}{2} > 0$ for $r \geq 2$ and is a half-integer (hence no logarithmic term appears). Specifically, for $a = b = \frac{1}{2}$, $c = r - \frac{1}{2}$, and $z \rightarrow 1^-$, the connection formula yields

$${}_2F_1(a, b; c; z) = \frac{\Gamma(c) \Gamma(c - a - b)}{\Gamma(c - a) \Gamma(c - b)} + O((1 - z)^{c - a - b}). \quad (62)$$

Setting $z = 1 - \frac{t+v}{2}$ we obtain, uniformly for $t, v \rightarrow 0^+$,

$${}_2F_1\left(\frac{1}{2}, \frac{1}{2}; r - \frac{1}{2}; 1 - \frac{t+v}{2}\right) = C_1(r) + O((t + v)^{r - \frac{3}{2}}), \quad (63)$$

where the leading constant $C_1(r)$ depends only on r and not on t or v (its closed form is given below). Consequently, at leading order the hypergeometric factor may be replaced by $C_1(r)$ in the neighborhood of $t = v = 0$.

The dominant contribution to the integral comes from the region where $v \approx t$ (the density concentrates around the mean). Near this region, $\arccos(1 - v) = \sqrt{2v} + O(v^{3/2})$, so:

$$\mathbb{E}[\arccos(\hat{\rho}_r)] = \int_0^\infty \sqrt{2v} \cdot \frac{(r - 2) \Gamma(r - 1) (2t)^{\frac{r-1}{2}} (2v)^{\frac{r-4}{2}}}{\sqrt{2\pi} \Gamma(r - \frac{1}{2}) (t + v)^{r - \frac{3}{2}}} C_1(r) dv + O(t^{3/2}). \quad (64)$$

Making the substitution $w = v/t$, we have $v = tw$, $dv = t dw$, and $t + v = t(1 + w)$:

$$\mathbb{E}[\arccos(\hat{\rho}_r)] = \int_0^\infty \sqrt{2tw} \cdot \frac{(r - 2) \Gamma(r - 1) (2t)^{\frac{r-1}{2}} (2tw)^{\frac{r-4}{2}}}{\sqrt{2\pi} \Gamma(r - \frac{1}{2}) (t(1 + w))^{r - \frac{3}{2}}} C_1(r) \cdot t dw + O(t^{3/2}). \quad (65)$$

The extension of the upper limit to ∞ is justified: for large w , the integrand behaves like $w^{\frac{r-3}{2}} / (1 + w)^{r - \frac{3}{2}} = O(w^{-r/2})$, which is integrable for all $r \geq 2$. Hence the tail beyond any fixed cutoff W contributes $o(t^{1/2})$ uniformly as $t \rightarrow 0^+$, and is absorbed into the $O(t^{3/2})$ remainder.

Simplifying the powers of t :

$$= \sqrt{2t} \cdot t^{\frac{r-1}{2} + \frac{r-4}{2} + 1 - (r - \frac{3}{2})} \cdot \frac{(r - 2) \Gamma(r - 1) (2)^{\frac{r-1}{2}} (2w)^{\frac{r-4}{2}}}{\sqrt{2\pi} \Gamma(r - \frac{1}{2}) (1 + w)^{r - \frac{3}{2}}} C_1(r) \cdot \sqrt{w} dw + O(t^{3/2}), \quad (66)$$

where the exponent is $\frac{r-1}{2} + \frac{r-4}{2} + 1 - (r - \frac{3}{2}) = 0$, so:

$$\mathbb{E}[\arccos(\hat{\rho}_r)] = \sqrt{2t} \cdot I(r) + O(t^{3/2}), \quad (67)$$

where

$$I(r) = \int_0^\infty \sqrt{w} \cdot \frac{(r - 2) \Gamma(r - 1) (2)^{\frac{r-1}{2}} (2w)^{\frac{r-4}{2}}}{\sqrt{2\pi} \Gamma(r - \frac{1}{2}) (1 + w)^{r - \frac{3}{2}}} C_1(r) dw. \quad (68)$$

Evaluating the integral yields the exact constant

$$I(r) = \frac{(r-2) 2^{r-\frac{5}{2}} \Gamma\left(\frac{r-1}{2}\right) \Gamma\left(\frac{r}{2}-1\right)}{\sqrt{2\pi} \Gamma(r-1)} C_1(r), \quad C_1(r) = \frac{\Gamma\left(r-\frac{1}{2}\right) \Gamma\left(r-\frac{3}{2}\right)}{\Gamma(r-1)^2}. \quad (69)$$

Therefore,

$$\mathbb{E}[\arccos(\hat{\rho}_r)] = \sqrt{2t} I(r) + O(t^{3/2}). \quad (70)$$

Therefore, the mean NTK expansion is:

$$\tilde{\Theta}^{(2)}(1-t) = K_\infty(1) - \left[\frac{2}{\pi} \frac{\sqrt{2}}{\pi} \Theta^{(1)}(1) \frac{I(r)}{2} \right] t^{1/2} + O(t^{3/2}), \quad (71)$$

so, for ReLU with $\Theta^{(1)}(1) = \frac{1}{2}$,

$$\tilde{\Theta}^{(2)}(1-t) = K_\infty(1) - \left[\frac{2}{\pi} + \frac{\sqrt{2}}{2\pi} I(r) \right] t^{1/2} + O(t^{3/2}). \quad (72)$$

This exhibits the complete r -dependent constant multiplying $t^{1/2}$.

Remark 9.2 (Closed form via Beta function and asymptotics). Using the change of variables $w = v/t$ and Euler's Beta integral, the inner integral appearing in $I(r)$ can be evaluated explicitly:

$$\int_0^\infty \frac{w^{\frac{r-3}{2}}}{(1+w)^{r-\frac{3}{2}}} dw = B\left(\frac{r-1}{2}, \frac{r}{2}-1\right) = \frac{\Gamma\left(\frac{r-1}{2}\right) \Gamma\left(\frac{r}{2}-1\right)}{\Gamma\left(r-\frac{3}{2}\right)}. \quad (73)$$

Substituting this identity into the pre-factors gives the stated closed form for $I(r)$ in (74) below, which matches the expression above:

$$I(r) = \frac{(r-2) 2^{r-\frac{5}{2}}}{\sqrt{2\pi}} \frac{\Gamma\left(\frac{r-1}{2}\right) \Gamma\left(\frac{r}{2}-1\right)}{\Gamma(r-1)} C_1(r), \quad C_1(r) = \frac{\Gamma\left(r-\frac{1}{2}\right) \Gamma\left(r-\frac{3}{2}\right)}{\Gamma(r-1)^2}. \quad (74)$$

Moreover, by Stirling's formula and standard Gamma-ratio bounds (e.g., Gautschi's inequality), one obtains for fixed $r \geq 3$:

$$0 < \inf_{r \geq 3} I(r) \leq I(r) \leq \sup_{r \geq 3} I(r) < \infty, \quad (75)$$

i.e., $I(r) = \Theta(1)$ as $r \rightarrow \infty$. In particular, the leading Puiseux coefficient

$$1 - \left[\frac{2}{\pi} + \frac{\sqrt{2}}{2\pi} I(r) \right] t^{1/2} + O(t^{3/2}) \quad (76)$$

is strictly positive and depends smoothly on r . This confirms that the endpoint exponent $\frac{1}{2}$ (hence the RKHS spectral decay rate) is unchanged by the Fisher–Kibble expectation, while the multiplicative constant is an $O(1)$ perturbation of the shallow ReLU constant $2/\pi$.

RKHS equivalence via endpoint behavior. The mean NTK $\tilde{\Theta}^{(2)}(\rho)$ is a zonal kernel with a Puiseux expansion that differs from the deterministic kernel $K_\infty(\rho)$ due to the Fisher–Kibble expectations. However, the leading-order behavior near $\rho = 1$ is preserved: both kernels have the same $t^{1/2}$ leading term in their Puiseux expansions.

By Theorem 1 of [17], the RKHS spectral decay is determined by the leading-order Puiseux expansion coefficient (the $t^{1/2}$ term). Since $\tilde{\Theta}^{(2)}(\rho)$ and the shallow ReLU kernel share the same leading-order expansion $1 - \frac{2}{\pi} t^{1/2} + \dots$ (with different higher-order corrections), they induce the same RKHS. The higher-order corrections from Fisher–Kibble expectations affect the $O(t^{3/2})$ and higher terms, but do not change the spectral decay rate, ensuring RKHS equivalence.

Extension to general L . For general depth L , the mean NTK involves nested expectations over Fisher and Kibble distributions at each layer. The key is that homogeneity ensures the endpoint Puiseux expansions are preserved (up to $O(r^{-1/2})$ corrections per layer), so the RKHS structure remains the same. Proofs is deferred to a future version. $\square$

10 Spectrum of 3-layers Gram NTK Matrix

10.1 Asymptotic Signal-Noise Decomposition

Proof of Theorem 5.1. We establish the asymptotic distribution of the empirical kernel $\hat{\Theta}_r^{(2)} = \Psi(\hat{\rho}_r, \hat{w}_r)$ via a joint CLT and the multivariate Delta method. Let $\mathbf{Z}_r = (\hat{\rho}_r, \hat{w}_r)^\top$. For isotropic Gaussian projections in rank r , Fisher's theory [15, 16] gives

$$\sqrt{r}(\hat{\rho}_r - \rho) \xrightarrow{d} \mathcal{N}(0, (1 - \rho^2)^2). \quad (77)$$

Writing $u = \|x\|^2$, $v = \|y\|^2$ and $\hat{w}_r = \sqrt{uv}/r$, Kibble's bivariate chi-square yields

$$\sqrt{r}(\hat{w}_r - 1) \xrightarrow{d} \mathcal{N}(0, 1 + \rho^2). \quad (78)$$

Angular and radial components are independent for isotropic Gaussians, hence

$$\sqrt{r}(\mathbf{Z}_r - (\rho, 1)^\top) \xrightarrow{d} \mathcal{N}(\mathbf{0}, \Lambda(\rho)), \quad \Lambda(\rho) = \begin{pmatrix} (1 - \rho^2)^2 & 0 \\ 0 & 1 + \rho^2 \end{pmatrix}. \quad (79)$$

The map $\Psi(u, w) = \Theta^{(1)}(u)(1 - \arccos(u)/\pi) + w \Sigma^{(1)}(u) + 1$ is C^1 near $(\rho, 1)$. Therefore,

$$\sqrt{r}(\Psi(\mathbf{Z}_r) - \Psi(\rho, 1)) \xrightarrow{d} \mathcal{N}(0, \nabla \Psi(\rho, 1)^\top \Lambda(\rho) \nabla \Psi(\rho, 1)). \quad (80)$$

We have $\partial_u \Psi(\rho, 1) = \partial_\rho K_\infty(\rho)$ and $\partial_w \Psi(\rho, 1) = \Sigma^{(1)}(\rho)$. Using the diagonal covariance, we obtain

$$\sigma_{\text{tot}}^2(\rho) = (\partial_\rho K_\infty(\rho))^2 (1 - \rho^2)^2 + (\Sigma^{(1)}(\rho))^2 (1 + \rho^2). \quad (81)$$

Hence

$$\hat{\Theta}_r^{(2)} = K_\infty(\rho) + \frac{1}{\sqrt{r}} \mathcal{G}(\rho) + o_p(r^{-1/2}), \quad \mathcal{G}(\rho) \sim \mathcal{N}(0, \sigma_{\text{tot}}^2(\rho)), \quad (82)$$

which proves the claim. $\square$

10.2 Proof of Theorem 5.2: Deformed Marchenko–Pastur with Spike

Proof. We prove the spectral limit using Theorem 2.1 of [53] on the spectrum of inner product kernel random matrices. This connects the microscopic kernel structure (ReLU/Kibble calculations) with the macroscopic matrix behavior (spectrum).

Setting. Let $\mathbf{M} = [\Theta_{ij}^{(2)}]_{i,j=1}^n$ be the $n \times n$ NTK matrix with entries $M_{ij} = \Theta^{(2)}(x_i, x_j) = K_\infty(\rho_{ij}) + O_p(r^{-1/2})$, where $\rho_{ij} = \langle x_i, x_j \rangle / (\|x_i\| \|x_j\|)$ is the cosine similarity and $K_\infty(\rho) = \Theta^{(1)}(\rho)(1 - \arccos(\rho)/\pi) + \Sigma^{(1)}(\rho) + 1$ is the deterministic three-layer kernel limit.

Under Assumption 5.1, we have $n \sim r$ (so n/p remains bounded where p is the ambient dimension) and the data points x_i are normalized. The kernel function $f(\rho) = K_\infty(\rho)$ is C^1 in a neighborhood of $l = \lim_{p \rightarrow \infty} \text{trace}(\Sigma_p)/p = 1$ (for normalized data) and C^3 in a neighborhood of 0, satisfying the regularity conditions of [53].

Step 1: Equivalent Matrix Structure. By Theorem 2.1 of [53], in probability as $n, p \rightarrow \infty$, the kernel matrix $\mathbf{M}$ is equivalent in operator norm to the matrix $\mathbf{K}$:

$$\mathbf{K} = \underbrace{\alpha \mathbf{1}\mathbf{1}^T}_{\text{Spike}} + \underbrace{\beta \frac{XX^T}{p}}_{\text{Bulk}} + \underbrace{\gamma I_n}_{\text{Shift}}, \quad (83)$$

where $X = [x_1, \dots, x_n] \in \mathbb{R}^{p \times n}$ is the data matrix. The three spectral coefficients α, β, γ are determined by the kernel function $f(\rho) = K_\infty(\rho)$ and its derivatives evaluated at $\rho = 0$ and $\rho = 1$:

$$\alpha = f(0) = K_\infty(0), \quad (84)$$

$$\beta = f'(0) = K'_\infty(0), \quad (85)$$

$$\gamma = f(1) - f(0) - f'(0) = K_\infty(1) - K_\infty(0) - K'_\infty(0). \quad (86)$$

Spectral Coefficients. The Spike (α) The spike coefficient is the kernel value at orthogonality ($\rho = 0$):

$$\alpha = K_\infty(0) = \Theta^{(1)}(0) \left(1 - \frac{\arccos(0)}{\pi}\right) + \Sigma^{(1)}(0) + 1. \quad (87)$$

Since $\arccos(0) = \pi/2$, we obtain:

$$\alpha = \Theta^{(1)}(0) \left(1 - \frac{\pi}{2\pi}\right) + \Sigma^{(1)}(0) + 1 = \Theta^{(1)}(0) \left(1 - \frac{1}{2}\right) + \Sigma^{(1)}(0) + 1 = \frac{\Theta^{(1)}(0)}{2} + \Sigma^{(1)}(0) + 1. \quad (88)$$

For ReLU activations with normalized inputs, $\Theta^{(1)}(0) = \Sigma^{(1)}(0) = 0$, hence $\alpha = 1$. The rank-one mean term $\alpha \mathbf{1}\mathbf{1}^T$ induces an outlier eigenvalue:

$$\lambda_{\text{spike}} = n \cdot \alpha = n \cdot K_\infty(0), \quad (89)$$

of order $O(n)$, isolated from the bulk spectrum.

B. The Bulk (β): The Linear Scale. The bulk coefficient is the derivative of the kernel at $\rho = 0$, capturing the linear sensitivity:

$$\beta = K'_\infty(0) = \frac{d}{d\rho} \left[\Theta^{(1)}(\rho) \left(1 - \frac{\arccos(\rho)}{\pi}\right) + \Sigma^{(1)}(\rho) + 1 \right]_{\rho=0}. \quad (90)$$

Differentiating term by term:

$$\begin{aligned} \frac{d}{d\rho} \left[\Theta^{(1)}(\rho) \left(1 - \frac{\arccos(\rho)}{\pi}\right) \right] &= \Theta^{(1)'}(\rho) \left(1 - \frac{\arccos(\rho)}{\pi}\right) + \Theta^{(1)}(\rho) \cdot \frac{1}{\pi\sqrt{1-\rho^2}}, \\ \frac{d}{d\rho} [\Sigma^{(1)}(\rho)] &= \Sigma^{(1)'}(\rho). \end{aligned}$$

Evaluating at $\rho = 0$ (where $\arccos(0) = \pi/2$ and $\sqrt{1-0} = 1$):

$$\beta = \Theta^{(1)'}(0) \left(1 - \frac{1}{2}\right) + \Theta^{(1)}(0) \cdot \frac{1}{\pi} + \Sigma^{(1)'}(0) = \frac{\Theta^{(1)'}(0)}{2} + \frac{\Theta^{(1)}(0)}{\pi} + \Sigma^{(1)'}(0). \quad (91)$$

For ReLU kernels with normalized inputs, a common normalization yields $\Theta^{(1)}(0) = 0$, $\Theta^{(1)'}(0) = 1/\pi$, and $\Sigma^{(1)'}(0) = 1/(2\pi)$, hence $\beta = 1/(2\pi) + 1/(2\pi) = 1/\pi$.

The term $\beta XX^T/p$ has the same spectral distribution as a Wishart matrix scaled by β . When $n \asymp p$ with aspect ratio $\gamma_{\text{ratio}} = n/r$, the bulk eigenvalues follow a deformed Marchenko–Pastur distribution scaled by β , supported on:

$$\left[\beta(1 - \sqrt{\gamma_{\text{ratio}}})^2, \beta(1 + \sqrt{\gamma_{\text{ratio}}})^2 \right]. \quad (92)$$

The Shift (γ) (Diagonal Residual) The shift coefficient accounts for the difference between the diagonal value and the linear approximation:

$$\gamma = K_\infty(1) - K_\infty(0) - K'_\infty(0) = \alpha + \beta \cdot 1 + \text{non-linear terms} - \alpha - \beta = \text{non-linear residual}. \quad (93)$$

Computing $K_\infty(1)$ (on-diagonal, where $\arccos(1) = 0$):

$$K_\infty(1) = \Theta^{(1)}(1) \cdot 1 + \Sigma^{(1)}(1) + 1 = \Theta^{(1)}(1) + \Sigma^{(1)}(1) + 1. \quad (94)$$

Therefore:

$$\gamma = [\Theta^{(1)}(1) + \Sigma^{(1)}(1) + 1] - \alpha - \beta. \quad (95)$$

For normalized ReLU kernels, $\Theta^{(1)}(1) = \Sigma^{(1)}(1) = 1/2$, so $K_\infty(1) = 1 + 1 + 1 = 3$, and $\gamma = 3 - 1 - 3/(2\pi) = 2 - 3/(2\pi) > 0$.

The diagonal shift γI_n adds γ to all eigenvalues, acting as natural Tikhonov regularization. This ensures the kernel matrix is well-conditioned (bounded away from zero) and stabilizes inversion.

Final Spectral Distribution. Combining the three components, the spectrum of $\mathbf{M}$ converges (in probability) to:

- **Spike (rank-one outlier):** A single eigenvalue $\lambda_{\text{spike}} = n \cdot \alpha = n \cdot K_{\infty}(0)$ of order $O(n)$, isolated from the bulk.
- **Bulk (deformed Marchenko–Pastur):** The remaining $n - 1$ eigenvalues follow a deformed MP distribution, supported on:

$$\left[\gamma + \beta(1 - \sqrt{\gamma_{\text{ratio}}})^2, \gamma + \beta(1 + \sqrt{\gamma_{\text{ratio}}})^2 \right], \quad (96)$$

where $\gamma_{\text{ratio}} = \lim_{n \rightarrow \infty} n/r \in (0, \infty)$. The bulk width is determined by $\beta = K'_{\infty}(0)$, and the bulk location is shifted by $\gamma = K_{\infty}(1) - K_{\infty}(0) - K'_{\infty}(0)$.

The theorem is proved. □