

Analysis II

Janis Hutz
<https://janishutz.com>

September 30, 2025

TITLE PAGE COMING SOON

“Some funny quote from the lecture still needed”

- Özlem Imamoglu, 2025

HS2025, ETHZ

Cheat-Sheet based on Lecture notes and Script

<https://metaphor.ethz.ch/x/2025/hs/401-0213-16L/sc/script-analysis-II.pdf>

Contents

1	Introduction	3
2	Differential Equations	4
2.1	Introduction	4

1 Introduction

This Cheat-Sheet does not serve as a replacement for solving exercises and getting familiar with the content. There is no guarantee that the content is 100% accurate, so use at your own risk. If you discover any errors, please open an issue or fix the issue yourself and then open a Pull Request here:

<https://github.com/janishutz/eth-summaries>

This Cheat-Sheet was designed with the HS2025 page limit of 10 A4 pages in mind. Thus, the whole Cheat-Sheet can be printed full-sized, if you exclude the title page, contents and this page. You could also print it as two A5 pages per A4 page and also print the [Analysis I summary](#) in the same manner, allowing you to bring both to the exam

2 Differential Equations

2.1 Introduction

Ex 2.1.1: $f'(x) = f(x)$ has only solution $f(x) = ae^x$ for any $a \in \mathbb{R}$; $f' - a = 0$ has only solution $f(x) = \int_{x_0}^x a(t) \, dt$

T 2.1.6: Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a differential function of two variables. Let $x_0 \in \mathbb{R}$ and $y_0 \in \mathbb{R}^2$. The Ordinary Differential Equation (ODE) $y' = F(x, y)$ has a unique solution f defined on a “largest” interval I that contains x_0 such that $y_0 = f(x_0)$