

STA 517 3.0 Programming and Statistical Computing with R

Tutorial 1

1 Creating vectors

1. Create the following vectors:

(a) 1, 2, 3, ..., 100

(b) 2, 4, 6, 8, ..., 100

(c)

[1]	1.000000	1.497487	1.994975	2.492462	2.989950	3.487437
[7]	3.984925	4.482412	4.979899	5.477387	5.974874	6.472362
[13]	6.969849	7.467337	7.964824	8.462312	8.959799	9.457286
[19]	9.954774	10.452261	10.949749	11.447236	11.944724	12.442211
[25]	12.939698	13.437186	13.934673	14.432161	14.929648	15.427136
[31]	15.924623	16.422111	16.919598	17.417085	17.914573	18.412060
[37]	18.909548	19.407035	19.904523	20.402010	20.899497	21.396985
[43]	21.894472	22.391960	22.889447	23.386935	23.884422	24.381910
[49]	24.879397	25.376884	25.874372	26.371859	26.869347	27.366834
[55]	27.864322	28.361809	28.859296	29.356784	29.854271	30.351759
[61]	30.849246	31.346734	31.844221	32.341709	32.839196	33.336683
[67]	33.834171	34.331658	34.829146	35.326633	35.824121	36.321608
[73]	36.819095	37.316583	37.814070	38.311558	38.809045	39.306533
[79]	39.804020	40.301508	40.798995	41.296482	41.793970	42.291457
[85]	42.788945	43.286432	43.783920	44.281407	44.778894	45.276382
[91]	45.773869	46.271357	46.768844	47.266332	47.763819	48.261307
[97]	48.758794	49.256281	49.753769	50.251256	50.748744	51.246231
[103]	51.743719	52.241206	52.738693	53.236181	53.733668	54.231156
[109]	54.728643	55.226131	55.723618	56.221106	56.718593	57.216080
[115]	57.713568	58.211055	58.708543	59.206030	59.703518	60.201005
[121]	60.698492	61.195980	61.693467	62.190955	62.688442	63.185930
[127]	63.683417	64.180905	64.678392	65.175879	65.673367	66.170854
[133]	66.668342	67.165829	67.663317	68.160804	68.658291	69.155779
[139]	69.653266	70.150754	70.648241	71.145729	71.643216	72.140704
[145]	72.638191	73.135678	73.633166	74.130653	74.628141	75.125628
[151]	75.623116	76.120603	76.618090	77.115578	77.613065	78.110553
[157]	78.608040	79.105528	79.603015	80.100503	80.597990	81.095477
[163]	81.592965	82.090452	82.587940	83.085427	83.582915	84.080402
[169]	84.577889	85.075377	85.572864	86.070352	86.567839	87.065327
[175]	87.562814	88.060302	88.557789	89.055276	89.552764	90.050251
[181]	90.547739	91.045226	91.542714	92.040201	92.537688	93.035176
[187]	93.532663	94.030151	94.527638	95.025126	95.522613	96.020101
[193]	96.517588	97.015075	97.512563	98.010050	98.507538	99.005025
[199]	99.502513	100.000000				

2. Generate a sequence using the code `seq(from=1, to=10, by=1)`. What other ways can you generate the same sequence?
3. Using the function `rep()` , create the below sequence 1, 2, 3, 4, 1, 2, 3, 4, 1, 2, 3, 4
4. Create a vector that shows the square root the integers from 1 to 100.
5. Observe the differences in running the following codes.

```
vec1 <- 1.8:20.8
vec1
```

```
[1] 1.8 2.8 3.8 4.8 5.8 6.8 7.8 8.8 9.8 10.8 11.8 12.8 13.8 14.8 15.8
[16] 16.8 17.8 18.8 19.8 20.8
```

```
vec2 <- 1.8:30
vec2
```

```
[1] 1.8 2.8 3.8 4.8 5.8 6.8 7.8 8.8 9.8 10.8 11.8 12.8 13.8 14.8 15.8
[16] 16.8 17.8 18.8 19.8 20.8 21.8 22.8 23.8 24.8 25.8 26.8 27.8 28.8 29.8
```

2 Object classes and type of objects

6. Use `typeof` to identify the storage mode of the following objects and `class` to identify object classes.

```
a <- c("MON", "TUES", "WED", "THUR", "FRI")
```

```
b <- c(1, 2, 3, 4, 5)
```

```
c <- c(1L, 2L, 3L, 4L, 5L)
```

```
d <- c(TRUE, FALSE, TRUE, TRUE)
```

```
e <- c(2+3i, 1+2i, 5+3i)
```

```
f <- c("MON", TRUE, 1, 1L)
```

7. Explore comment on the output of following vector functions.

```
a1 <- vector("numeric", 8)
a2 <- vector("complex", 8)
a3 <- vector("logical", 8)
a4 <- vector("character", 8)
```

```
b1 <- numeric(8)
b2 <- complex(8)
b3 <- logical(8)
b4 <- character(8)
```

8. Consider the vector

```
set.seed(32020)
st_normal <- rnorm(100)
st_normal
```

```
[1] 0.18183635 -0.92262020 2.06110995 -1.50040396 -1.69529463 2.45410426
[7] 0.16552699 -2.20702891 -0.21274657 -0.69387976 -0.67516314 1.03136276
[13] 0.77649171 0.60913641 -1.06664784 0.34027083 -0.47879695 -0.40281847
[19] -1.12500580 -0.79235873 -0.89371755 -2.72593829 0.99052081 -0.53966792
[25] 2.44848942 1.82337921 -0.52409631 -2.52099047 -0.01338390 -0.67771367
[31] -0.26224412 -1.96067034 0.03172268 -0.83045197 1.60051305 0.04106971
[37] 0.93303006 -1.31390340 -0.25427286 -0.61430209 -0.09897693 0.33713741
[43] 0.45989743 -0.79752346 -0.77387974 -0.57871649 -1.24023942 -1.74035257
[49] -0.02742062 -2.21931034 0.23715755 -0.47101092 -0.22116294 -1.45243410
[55] 0.27650330 -1.76656058 0.01328862 -1.30263545 1.20788668 1.47504605
[61] -2.19540879 0.44796633 0.39314554 -3.15206211 -0.32687439 -0.54550496
[67] 1.39978830 -2.19770996 1.46683852 -1.19686302 0.87487978 -0.83723410
[73] 1.37510059 -0.80996752 0.56198382 0.40264681 0.13343941 -0.05576293
[79] 1.66654211 -0.78997663 0.29758171 0.36613867 0.80338650 -1.43640458
[85] -0.56015981 -0.12409835 -0.75476839 0.32283051 1.46941104 -0.30940270
[91] -1.14718708 -0.93229533 0.06524165 -0.20590515 -0.69251943 0.93134043
[97] 0.28856808 1.04544874 0.24806814 0.22931507
```

Drop the elements corresponds to the positions multiply of 10 (10, 20, 30, ...)

```
st_normal[-seq(1, 100, by=10)]
```

```
[1] -0.92262020 2.06110995 -1.50040396 -1.69529463 2.45410426 0.16552699
[7] -2.20702891 -0.21274657 -0.69387976 1.03136276 0.77649171 0.60913641
[13] -1.06664784 0.34027083 -0.47879695 -0.40281847 -1.12500580 -0.79235873
[19] -2.72593829 0.99052081 -0.53966792 2.44848942 1.82337921 -0.52409631
[25] -2.52099047 -0.01338390 -0.67771367 -1.96067034 0.03172268 -0.83045197
[31] 1.60051305 0.04106971 0.93303006 -1.31390340 -0.25427286 -0.61430209
[37] 0.33713741 0.45989743 -0.79752346 -0.77387974 -0.57871649 -1.24023942
[43] -1.74035257 -0.02742062 -2.21931034 -0.47101092 -0.22116294 -1.45243410
[49] 0.27650330 -1.76656058 0.01328862 -1.30263545 1.20788668 1.47504605
[55] 0.44796633 0.39314554 -3.15206211 -0.32687439 -0.54550496 1.39978830
[61] -2.19770996 1.46683852 -1.19686302 -0.83723410 1.37510059 -0.80996752
[67] 0.56198382 0.40264681 0.13343941 -0.05576293 1.66654211 -0.78997663
[73] 0.36613867 0.80338650 -1.43640458 -0.56015981 -0.12409835 -0.75476839
[79] 0.32283051 1.46941104 -0.30940270 -0.93229533 0.06524165 -0.20590515
[85] -0.69251943 0.93134043 0.28856808 1.04544874 0.24806814 0.22931507
```

9. Create a vector with elements from 1 to 100 incrementing by 0.4

```
seq(1, 100, by=0.4)
```

```
[1] 1.0 1.4 1.8 2.2 2.6 3.0 3.4 3.8 4.2 4.6 5.0 5.4 5.8 6.2 6.6
[16] 7.0 7.4 7.8 8.2 8.6 9.0 9.4 9.8 10.2 10.6 11.0 11.4 11.8 12.2 12.6
[31] 13.0 13.4 13.8 14.2 14.6 15.0 15.4 15.8 16.2 16.6 17.0 17.4 17.8 18.2 18.6
[46] 19.0 19.4 19.8 20.2 20.6 21.0 21.4 21.8 22.2 22.6 23.0 23.4 23.8 24.2 24.6
[61] 25.0 25.4 25.8 26.2 26.6 27.0 27.4 27.8 28.2 28.6 29.0 29.4 29.8 30.2 30.6
```

```
[76] 31.0 31.4 31.8 32.2 32.6 33.0 33.4 33.8 34.2 34.6 35.0 35.4 35.8 36.2 36.6
[91] 37.0 37.4 37.8 38.2 38.6 39.0 39.4 39.8 40.2 40.6 41.0 41.4 41.8 42.2 42.6
[106] 43.0 43.4 43.8 44.2 44.6 45.0 45.4 45.8 46.2 46.6 47.0 47.4 47.8 48.2 48.6
[121] 49.0 49.4 49.8 50.2 50.6 51.0 51.4 51.8 52.2 52.6 53.0 53.4 53.8 54.2 54.6
[136] 55.0 55.4 55.8 56.2 56.6 57.0 57.4 57.8 58.2 58.6 59.0 59.4 59.8 60.2 60.6
[151] 61.0 61.4 61.8 62.2 62.6 63.0 63.4 63.8 64.2 64.6 65.0 65.4 65.8 66.2 66.6
[166] 67.0 67.4 67.8 68.2 68.6 69.0 69.4 69.8 70.2 70.6 71.0 71.4 71.8 72.2 72.6
[181] 73.0 73.4 73.8 74.2 74.6 75.0 75.4 75.8 76.2 76.6 77.0 77.4 77.8 78.2 78.6
[196] 79.0 79.4 79.8 80.2 80.6 81.0 81.4 81.8 82.2 82.6 83.0 83.4 83.8 84.2 84.6
[211] 85.0 85.4 85.8 86.2 86.6 87.0 87.4 87.8 88.2 88.6 89.0 89.4 89.8 90.2 90.6
[226] 91.0 91.4 91.8 92.2 92.6 93.0 93.4 93.8 94.2 94.6 95.0 95.4 95.8 96.2 96.6
[241] 97.0 97.4 97.8 98.2 98.6 99.0 99.4 99.8
```

10. Consider the vector `x`.

```
x <- 1:10
```

What does each of the following codes do?

```
x[3]
x[c(2, 4)]
x[-1]
x[c(2, -4)]
x[c(2.4, 3.54)]
```

3 Filtering vectors based on conditions

11. Consider the vector

```
x <- c(80, 39, NA, 51, 51, 11, NA, NA, NA, 100, 80, 70)
```

Write an R code to extract non-missing values in `x`

Write an R code to extract missing values and odd-numbers in `x`

Write an R code to extract odd numbers on `x`

Which values of `x` are NOT in the set 1:50

4 Modify a vector

12. Consider the following vector `age` which includes the age of 10 individuals

```
age <- c(20, 30, 40, 41, 32, 32, 25, NA, NA, -4, -6, 9999, 10000)
```

i. Convert all negative values to ``NA``.

ii. Convert all values of ``age`` that are NOT from 10 to 100 and calculate the mean of valid responses.

15. Consider the following vector of 100 random numbers generated from the standard normal distribution.
- i. Change the first five values in the vector to 1.
 - ii. Change the last five values in the vector to 0.
 - iii. Assign all values greater than 0.5 to 1 and all values less than 0.5 to 0.
 - iv. Recode the 0 values to “MALE” and others to “FEMALE”