

# Rényi pseudo-distances

Bc. Jan Kučera

FNSPE, CTU

Work supervisor: Ing. Václav Kůs Ph.D.

26.6.2012

# Table of contents

- 1 Decomposable pseudo-distance
- 2 Rényi pseudo-distance
- 3 Applications for distributions
  - Normal distribution
  - Laplace distribution
  - Exponential distribution
  - Weibull distribution

# Decomposable pseudo-distance

We say, that  $\mathfrak{D} : \mathcal{P} \rightarrow \mathbb{R}$  is *pseudo-distance* if  $\forall P, Q \in \mathcal{P}$

$$\mathfrak{D}(P, Q) \geq 0 \quad \text{and} \quad \mathfrak{D}(P, Q) = 0 \Leftrightarrow P = Q$$

and this pseudo-distance is *decomposable* if there exist functionals so that  $\mathfrak{D}^0 : \mathcal{P} \rightarrow \mathbb{R}$ ,  $\mathfrak{D}^1 : \tilde{\mathcal{P}} \rightarrow \mathbb{R}$  and measurable  $\rho_\theta : \mathcal{X} \rightarrow \mathbb{R}$ ,  $\theta \in \Theta$ , so that  $\forall \theta \in \Theta$  and  $\forall Q \in \tilde{\mathcal{P}}$  there exists finite  $\int \rho_\theta dQ$  and

$$\mathfrak{D}(P_\theta, Q) = \mathfrak{D}^0(P_\theta) + \mathfrak{D}^1(Q) + \int \rho_\theta dQ.$$

- we don't presume triangle inequality or symmetry

## Minimal distance estimator

Functional  $T_{\mathfrak{D}} : \tilde{\mathcal{P}} \rightarrow \Theta$  defines *minimal distance* estimator if  $\mathfrak{D}(P_{\theta}, Q)$  is decomposable and  $T_{\mathfrak{D}}(Q) \in \Theta$  minimizes

$$T_{\mathfrak{D}}(Q) = \arg \min_{\theta \in \Theta} \left[ \mathfrak{D}^0(P_{\theta}) + \int \rho_{\theta} dQ \right], \quad \forall Q \in \tilde{\mathcal{P}}$$

## Rényi pseudo-distance

Let for some  $\beta > 0$

$$p^\beta, q^\beta, \ln p \in L_1(Q), \quad \forall P \in \mathcal{P}, Q \in \tilde{\mathcal{P}}.$$

holds. Then for some  $\alpha$ ,  $0 < \alpha \leq \beta$ ,  $P \in \mathcal{P}$ ,  $Q \in \tilde{\mathcal{P}}$  is decomposable, that is

$$\mathcal{R}_\alpha(P, Q) = \mathcal{R}_\alpha^0(P) + \mathcal{R}_\alpha^1(Q) - \frac{1}{\alpha} \ln \left( \int p^\alpha dQ \right),$$

where

$$\mathcal{R}_\alpha^0(P) = \frac{1}{1+\alpha} \ln \left( \int p^\alpha dP \right), \quad \mathcal{R}_\alpha^1(Q) = \frac{1}{\alpha(1+\alpha)} \ln \left( \int q^\alpha dQ \right).$$

Moreover for  $\alpha \searrow 0$

$$\mathcal{R}_0(P, Q) = \int (\ln q - \ln p) dQ$$

If we replace  $Q$  by the empirical distribution  $P_n$ , we get

$$\theta_{\mathfrak{R}_{\alpha,n}} = \begin{cases} \arg \max_{\theta \in \Theta} \left( \int p_{\theta}^{1+\alpha}(x) dx \right)^{-\frac{\alpha}{1+\alpha}} \frac{1}{n} \sum_{i=1}^n p_{\theta}^{\alpha}(x_i), \\ \quad \text{for } 0 < \alpha \leq \beta, \\ \arg \max_{\theta \in \Theta} \frac{1}{n} \sum_{i=1}^n \ln p_{\theta}(x_i), \\ \quad \text{for } \alpha = 0. \end{cases}$$


So for  $\alpha = 0$  the  $\theta_{\alpha,n} = \theta_{MLE}$

# Normal distribution

$$p_{\theta} = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left[ -\frac{(x - \mu)^2}{2\sigma} \right]$$


$$\theta_{\mathfrak{R}_{\alpha}, n} = \arg \max_{\theta \in \Theta} \frac{1}{n \sigma^{\frac{\alpha}{1+\alpha}}} \sum_{i=1}^n \exp \left[ -\alpha \frac{(x_i - \mu)^2}{2\sigma^2} \right]$$

distances in the parametric space:






$$\text{IF}(x; T_{\mathfrak{R}_\alpha}, \mu) = (1 + \alpha)^{\frac{3}{2}}(x - \mu)e^{-\frac{\alpha}{2}(x - \mu)^2}$$



$$\text{IF}(x; T_{\mathfrak{R}_\alpha, \sigma}) = \frac{(1 + \alpha)^{\frac{5}{2}} \sigma}{2} \left( \left( \frac{x}{\sigma} \right)^2 - \frac{1}{1 + \alpha} \right) e^{-\frac{\alpha x^2}{2\sigma^2}}$$



$$\lim_{x \rightarrow \pm\infty} \text{IF}(x; T_{\mathfrak{R}_\alpha}, \cdot) = 0$$

→ estimator is robust against outliers

$$\sup_x |\text{IF}(x; T_{\mathfrak{R}_\alpha}, \cdot)| \text{ is finite}$$

→ estimator is robust against point errors

$\alpha \backslash n$	500		
	$m(\mu)$	$s(\mu)$	$eref(\mu)$
	$m(\sigma)$	$s(\sigma)$	$eref(\sigma)$
0.0	0.000	0.037	1.000
	1.672	0.051	1.000
0.01	-0.000	0.037	1.032
	1.652	0.050	1.031
0.05	0.001	0.033	1.248
	1.572	0.046	1.244
0.1	-0.000	0.031	1.426
	1.478	0.041	1.560
0.2	-0.000	0.029	1.650
	1.333	0.035	2.097
0.5	-0.000	0.028	1.733
	1.156	0.028	3.295

Table: Rényi:  $p = N(0, 1)$ , data:  $(1 - \varepsilon)N(0, 1) + \varepsilon N(0, 10)$ ,  $\varepsilon = 0.2$ ,  
 $K = 10000$

$\alpha \backslash n$	500		
	$m(\mu)$ $m(\sigma)$	$s(\mu)$ $s(\sigma)$	$eref(\mu)$ $eref(\sigma)$
0.0	-0.000 0.949	0.021 0.016	1.000 1.000
0.01	-0.000 0.949	0.021 0.016	1.002 0.969
0.05	0.001 0.945	0.021 0.016	0.991 0.967
0.1	0.000 0.942	0.021 0.017	0.990 0.889
0.2	-0.001 0.931	0.021 0.017	1.032 0.865
0.5	-0.000 0.905	0.021 0.020	0.998 0.641

Table: Rényi:  $p = N(0, 1)$ , data:  $0.9N(0, 1) + 0.1N_{0.1x}(0, 1)$ ,  $K = 1000$

$\alpha \backslash n$	500		
	$m(\mu)$ $m(\sigma)$	$s(\mu)$ $s(\sigma)$	$eref(\mu)$ $eref(\sigma)$
0.0	-0.002	0.075	1.000
	3.294	0.231	1.000
0.01	-0.000	0.065	1.327
	3.104	0.220	1.103
0.05	0.001	0.040	3.489
	2.118	0.189	1.493
0.1	0.001	0.026	8.356
	1.277	0.053	18.986
0.2	0.000	0.025	8.796
	1.087	0.024	91.365
0.5	-0.000	0.026	8.390
	1.029	0.021	123.291


Table: Rényi:  $p = N(0, 1)$ , data:  $0.9N(0, 1) + 0.1N_{10.0x}(0, 1)$ ,  $K = 1000$

# Laplace distribution

$$p_{\theta} = \frac{1}{2\lambda} e^{-\frac{|x-\mu|}{\lambda}}$$


$$\theta_{\mathfrak{R}_{\alpha},n} = \arg \max_{\theta \in \Theta} (2\lambda)^{-\frac{\alpha}{1+\alpha}} \frac{1}{n} \sum_{i=1}^n \exp \left[ -\alpha \frac{|x_i - \mu|}{\lambda} \right]$$

$$\text{IF}(x; T_{\mathfrak{R}_\alpha}, \mu) = (1 + \alpha)^{\frac{3}{2}} (x - \mu) e^{-\frac{\alpha}{2}(x - \mu)^2}$$





$$\text{IF}(x; T_{\mathfrak{R}_\alpha}, \lambda) = (1 + \alpha)^2 (-\lambda + (1 + \alpha)|x|) e^{-\frac{\alpha|x|}{\lambda}}$$



$\alpha \backslash n$	500		
	$m(\mu)$ $m(\lambda)$	$s(\mu)$ $s(\lambda)$	$eref(\mu)$ $eref(\lambda)$
0.0	0.000	0.065	1.000
	3.723	0.355	1.000
0.01	0.003	0.063	1.090
	3.593	0.338	1.102
0.05	0.002	0.063	1.063
	3.239	0.306	1.348
0.1	-0.002	0.062	1.118
	2.796	0.285	1.556
0.2	0.000	0.062	1.105
	2.059	0.205	3.005
0.3	-0.003	0.060	1.198
	1.626	0.154	5.326
0.5	0.000	0.058	1.267
	1.300	0.108	10.756
1.0	-0.001	0.068	0.932
	1.126	0.110	10.386


Table: Rényi:  $p = L(0, 1)$ , data:  $(1 - \varepsilon)L(0, 1) + \varepsilon L(0, 10)$ ,  $\varepsilon = 0.3$ ,  
 $K = 1000$

# Exponential distribution


$$p_{\theta} = \frac{1}{\lambda} e^{-\frac{x-\mu}{\lambda}}$$

$$\theta_{\mathfrak{R}_{\alpha}, n} = \arg \max_{\theta \in \Theta} \lambda^{-\frac{\alpha}{1+\alpha}} \frac{1}{n} \sum_{i=1}^n \exp \left[ -\alpha \frac{x_i - \mu}{\lambda} \right]$$

$$\text{IF}(x; T_{\mathfrak{R}_\alpha}, \mu) = e^{-\frac{\alpha}{2}(x-\mu)^2} (1 + \alpha)^{\frac{3}{2}} (x - \mu)$$



$$\text{IF}(x; T_{\mathfrak{R}_\alpha}, \lambda) = (1 + \alpha)^2 (-\lambda + (1 + \alpha)x) e^{-\frac{\alpha x}{\lambda}}$$



$\alpha \backslash n$	500		
	$m(\lambda)$	$s(\lambda)$	$eref(\lambda)$
0.0	3.901	0.432	1.000
0.01	3.614	0.356	1.475
0.05	3.241	0.317	1.862
0.1	2.796	0.274	2.495
0.15	2.389	0.239	3.258
0.2	2.049	0.209	4.281
0.3	1.622	0.153	7.999
0.5	1.310	0.110	15.330
1.0	1.139	0.109	15.806

Table: Rényi:  $p = E(0, 1)$ , data:  $(1 - \varepsilon)E(0, 1) + \varepsilon E(0, 10)$ ,  $\varepsilon = 0.3$ ,  
 $K = 1000$

# Weibull distribution

$$p_{\theta} = \frac{k}{\lambda} \left( \frac{x - \mu}{\lambda} \right)^{k-1} \exp \left[ - \left( \frac{x - \mu}{\lambda} \right)^k \right]$$

$$\begin{aligned} \theta_{\mathfrak{R}_{\alpha,n}} &= \arg \max_{\theta \in \Theta} \left( \frac{k}{\lambda} \right)^{\frac{\alpha}{1+\alpha}} (1 + \alpha)^{\frac{\alpha}{1+\alpha} \frac{1+\alpha+k}{k}} \Gamma \left( \frac{1 + \alpha + k}{k} \right)^{-\frac{\alpha}{1+\alpha}} \\ &\quad \frac{1}{n} \sum_{i=1}^n \left( \frac{x_i - \mu}{\lambda} \right)^{\alpha(k-1)} \exp \left[ -\alpha \left( \frac{x_i - \mu}{\lambda} \right)^k \right] \end{aligned}$$

$\alpha \backslash n$	500		
	$m(\lambda)$	$s(\lambda)$	$eref(\lambda)$
0.0	4.713	0.387	1.000
0.01	4.568	0.364	1.130
0.05	4.149	0.373	1.077
0.1	3.511	0.383	1.020
0.15	2.574	0.428	0.815
0.2	1.664	0.261	2.201
0.3	1.226	0.076	25.766
0.5	1.096	0.057	46.725
1.0	1.043	0.055	49.962

Table: Rényi:  $p = W(1.5, 1)$ , data:  $(1 - \varepsilon)W(1.5, 1) + \varepsilon W(1.5, 10)$ ,  
 $\varepsilon = 0.3$ ,  $K = 1000$



Thank you for your attention.