

Computer Assignment # 5

Due: May 26, 2024, 23:59:59

Consider the following unconstrained problem

$$\min_x f(x) = - \sum_{i=1}^m \log(1 - \mathbf{a}_i^T \mathbf{x}) - \sum_{j=1}^n \log\left(1 - \frac{1}{2}x_j^2\right),$$

where variable $\mathbf{x} \in \mathbb{R}^n$, and **dom** $f = \{\mathbf{x} : \mathbf{a}_i^T \mathbf{x} < 1, i = 1, 2, \dots, m, |x_j| \leq 1, j = 1, 2, \dots, n\}$. This is the problem of computing the analytic center of the set of linear inequalities

$$\mathbf{a}_i^T \mathbf{x} < 1, i = 1, 2, \dots, m, ; |x_j| < 1, j = 1, 2, \dots, n.$$

Note that you can choose $x^{(0)} = 0$ as your initial point. You can generate instances of this problem by choosing \mathbf{a}_i from some distribution on \mathbb{R}^n .

- Use the gradient method to solve the problem, using reasonable choices for the backtracking parameters, and a stopping criterion of a form $\|\nabla f(\mathbf{x})\|_2 \leq \eta$. Plot the objective function and step length versus iteration number. Experiment with the backtracking parameters α and β to see their effect on the total number of iterations required. Carry these experiments out for several instances of the problem, of different sizes.
- Repeat using Newton's method, with stopping criterion based on the Newton decrement λ^2 . Use the chain rule to find expressions for $\nabla f(\mathbf{x})$ and $\nabla^2 f(\mathbf{x})$

Submission Policy leftmargin=*

- Place all your scripts inside the folder called "codes".
- Place convergence figures with verbal explanations inside report (word or pdf).
- Put "codes" and report inside zip file called `ca05_XXXXXX.zip`, where XXXXXX is your student ID and submit it to e3.