

# Weighted Finite Automata and Noncommutative Rational Series

Jan Pantner ([jan.pantner@gmail.com](mailto:jan.pantner@gmail.com))

October 1, 2024

# 1 WFA, Linear Representations, Rational Series

## 1.1 Weighted Finite Automata

**Definition 1.1.1.** Let  $K$  be a semiring and  $A$  an alphabet.

- (1) A **weighted (finite) automaton** (WFA) with weights in  $K$  is a tuple  $(Q, I, E, T)$  consisting of a finite set  $Q$  of **states**, and maps  $I: Q \rightarrow K$  (**initial weights**),  $E: Q \times A \times Q \rightarrow K$  (**transition function**),  $T: Q \rightarrow K$  (**terminal weights**).
- (2) A triple  $(p, a, q)$  with  $E(p, a, q) \neq 0$  is an **edge/transition** with **label**  $a$ , **starting state**  $p$ , **ending state**  $q$  and **weight**  $E(p, a, q) \in K$ .

- (3) A **path/run** is a sequence of edges  $c = (q, a_1, q_1)(q_1, a_2, p_2) \dots (q_{n-1}, a_n, q_n)$ . Its **weight** is

$$E(q, a_1, q_1) \cdot E(q_1, a_2, p_2) \cdots E(q_{n-1}, a_n, q_n)$$

and its **label** is  $w = a_1 a_2 \dots a_n \in A^*$ .

- (4) The **behaviour** of  $\mathcal{A}$  is the series  $[[\mathcal{A}]] \in K \langle\langle A \rangle\rangle$  defined by

$$([\mathcal{A}], w) = \sum_{q_0, \dots, q_n \in Q} I(q_0) E(q, a_1, q_1) E(q_1, a_2, p_2) \cdots E(q_{n-1}, a_n, q_n) T(q_n),$$

where  $w = a_1 \cdots a_n$ ,  $a_i \in A$ .

**Definition 1.1.2.** Terminology:

1. A state  $q$  is **initial** if  $I(q) \neq 0$  and **terminal** if  $T(q) \neq 0$ .
2. A **successful run/accepting run** is a run from an initial state to a terminal state.

## 1.2 Linear representation, recognizable series

**Definition 1.2.1.** Let  $K$  be a semiring and  $A$  an alphabet.

- (1) A series  $S \in K \langle\langle A \rangle\rangle$  is **(K-)recognizable** if there exist  $n \geq 0$ ,  $\lambda \in K^{1 \times b}$ ,  $\gamma \in K^{n \times 1}$ , and a monoid morphism  $\mu: A^* \rightarrow K^{d \times d}$  such that for every  $w \in A^*$  we have

$$(S, w) = \lambda \mu(w) \gamma.$$

- (2) The triple  $(\lambda, \mu, \gamma)$  is a **linear representation** of  $S$  with **dimension**  $n$ .

**Proposition 1.2.2.** A series  $S \in K \langle\langle A \rangle\rangle$  is recognizable if and only if there exists a WFA  $\mathcal{A}$  such that  $S = [[\mathcal{A}]]$ .

*Proof.*

□

### 1.3 Model-theoretic Characterization

**Lemma 1.3.1.** Let  $K$  be a semiring and  $A$  an alphabet.

- (1) For  $x \in A^*$ , the map  $S \mapsto x^{-1}S$  is a  $K$ -module morphism.
- (2) For every  $x, y \in A^*$  and for every  $S \in K \langle\langle A \rangle\rangle$  we have  $(xy)^{-1}S = y^{-1}(x^{-1}S)$ .

*Proof.* Left as an exercise. □

**Definition 1.3.2.** A submodule  $M \subseteq K \langle\langle A \rangle\rangle$  is **stable** if for every  $S \in M$  and every  $x \in A^*$  we have  $x^{-1}S \subseteq M$  (equivalently for every  $a \in A$  we have  $a^{-1}M \subseteq M$ ).

**Theorem 1.3.3.** A series  $S \in K \langle\langle A \rangle\rangle$  is recognizable if and only if there exists a stable finitely generated left submodule  $M \subseteq K \langle\langle A \rangle\rangle$  such that  $S \in M$ .

*Proof.* □

October 8, 2024

## Index

linear representation, [2](#)

recognizable series, [2](#)

weighted finite automaton, [2](#)