

PART III Applications

Classical moment problems

December 10, 2025

Hamburger's moment problem:

Given a sequence of numbers $\{h_k\}_{k=0}^{\infty}$, decide if there exists a measure μ on \mathbb{R} s.t. $h_k = \int_{\mathbb{R}} x^k d\mu$, $k \geq 0$.

(\Leftrightarrow which sequences of reals are moment sequences of measures?)

Trigonometric moment problem:

Given a sequence $\{t_k\}_{k=-\infty}^{\infty} \subset \mathbb{C}$, is there a $\mu \geq 0$ s.t. $t_k = \int_{\mathbb{T}} z^k d\mu$?

Obvious restrictions:

• Hamburger case: $\sum_{k,j=0}^N h_{k+j} a_k \bar{a}_j \stackrel{(*)}{\geq} 0 \quad \forall \{a_k\}_{k=0}^N \subset \mathbb{C}, N \geq 0$

$$0 \leq \int_{\mathbb{R}} \left| \sum_{k=0}^N a_k x^k \right|^2 d\mu = \sum_{k,j=0}^N a_k \bar{a}_j \int_{\mathbb{R}} x^{k+j} d\mu = \sum_{k,j=0}^N a_k \bar{a}_j h_{k+j}$$

• Trigonometric case: $\sum_{k,j=-N}^N t_{k-j} a_k \bar{a}_j \stackrel{(**)}{\geq} 0 \quad \forall \{a_k\}_{k=-N}^N \subset \mathbb{C}, N \geq 0$

$$0 \leq \int_{\mathbb{T}} \left| \sum_{k=-N}^N a_k z^k \right|^2 d\mu = \sum_{k,j=-N}^N a_k \bar{a}_j \int_{\mathbb{T}} z^k \bar{z}^j d\mu = \sum_{k,j=-N}^N a_k \bar{a}_j t_{k-j}$$

Theorem [Hamburger]: The assumption $(*)$ is sufficient for the solvability of the Hamburger case.

Theorem: The assumption $(**)$ is sufficient for the solvability of the Trigonometric moment problem. Moreover, we have $\sum_{k,j=-N}^N t_{k-j} a_k \bar{a}_j > 0 \quad \forall \{a_k\}_{k=-N}^N \Leftrightarrow \{t_k\}$ is the moment sequence of a measure μ such that $\# \text{supp } \mu = +\infty$. [$(***)$]

Our goal is to prove $(***)$.

Proof: $H_0 = \left(\text{span}\{z^k\}_{k \in \mathbb{Z}}, \left\langle \sum_{-N}^N a_k z^k, \sum_{-N}^N \tilde{a}_k z^k \right\rangle := \sum_{k,j} t_{k-j} a_k \bar{\tilde{a}}_j \right)$

\hookrightarrow pre Hilbert space, because it is linear, and $\langle \cdot, \cdot \rangle$ is the inner product on H_0 , but H_0 is not complete w.r.t. $\|\sum_{-N}^N a_k z^k\| = \sqrt{\langle \sum_{-N}^N a_k z^k, \sum_{-N}^N a_k z^k \rangle}$

General functional analysis implies that $\exists H$ -Hilbert space such that $H_0 \subset H$ as a dense linear subset.

$T: \sum_{k=0}^N a_k z^k \longmapsto \sum_{k=0}^N a_k z^{k+1}$ - densely defined operator on H :

$$\begin{aligned} \|T(\sum_{-N}^N a_k z^k)\|^2 &= \|\sum_{-N}^N a_k z^{k+1}\|^2 = \sum_{-N+1}^{N+1} t_{k-j} a_{k-1} \overline{a_{j-1}} = \sum_{-N+1}^{N+1} t_{(k-1)-(j-1)} a_{k-1} \overline{a_{j-1}} = \\ &= \sum_{-N+1}^{N+1} a_{k-1} \overline{z^k} = \sum_{-N}^N t_{k-j} a_k \bar{a}_j = \|\sum_{-N}^N a_k z^k\|^2 \end{aligned}$$

$\Rightarrow T$ is an isometry initially defined on H_0 .

Let's extend it to the whole space H . $\Rightarrow T$ is isometry on H ,

$T(H) \supset H_0$ - dense in H , since $T(H)$ is closed, we have $T(H) = H$.

$\Rightarrow T$ is unitary. Moreover, there is $h=1$ s.t. $\text{span}\{T^k T^{*j} h\}$ is dense in H .
(H_0)

By the spectral theorem, there is a measure μ s.t. $\text{supp } \mu = \mathbb{T} \subset \mathbb{T}$:

$T \cong M_z$ on $L^2(\mu)$.

$$\langle T^k h, h \rangle_H = \langle M_z^k 1, 1 \rangle_{L^2(\mu)} \quad \forall k \geq 0 \text{ for } h=1 \text{ in } H.$$

||

$$\langle T^k 1, 1 \rangle_H = \langle z^k, 1 \rangle_H = \sum_0^k t_{i-j} \delta_k(i) \delta_0(j) = t_k$$

$$\langle M_z^k 1, 1 \rangle_{L^2(\mu)} = \langle z^k, 1 \rangle_{L^2(\mu)} = \int_{\mathbb{T}} z^k d\mu$$

$$\Rightarrow t_k = \int_{\mathbb{T}} z^k d\mu, \quad k \geq 0$$

$$\underline{t_{-k}} = \overline{t_k} = \overline{\int_{\mathbb{T}} z^k d\mu} = \int_{\mathbb{T}} \overline{z^k} d\mu = \int_{\mathbb{T}} z^{-k} d\mu, \quad k \geq 0 \quad \left. \vphantom{\int_{\mathbb{T}} z^k d\mu} \right\} \Rightarrow t_k \text{ is the moment sequence}$$

$$\left\{ \begin{aligned} t_k &= \langle z^k, 1 \rangle \\ t_{-k} &= \langle z^{-k}, 1 \rangle = \langle T^k z^{-k}, T^k 1 \rangle = \langle 1, z^k \rangle = \overline{\langle z^k, 1 \rangle} = \overline{t_k} \Rightarrow \underline{t_{-k}} = t_k \end{aligned} \right.$$

It remains to show that the measure μ is such that $\# \text{supp } \mu = \infty$.

$$\Leftrightarrow \int_{\mathbb{T}} |\sum_{-N}^N a_k z^k|^2 d\mu > 0 \quad (\text{true by assumption}).$$



December 11, 2025