

# RWorksheet\_Soco#4a.Rmd

JanRay P. Soco

2025-11-06

*#1. The table below shows the data about shoe size and height. Create a data frame.*

```
shoe <- c(6.5,9.0,8.5,8.5,10.5,7.0,9.5,9.0,13.0,7.5,10.5,8.5,12.0,10.5,
          13.0,11.5,8.5,5.0,10.0,6.5,8.5,10.5,6.5,10.5,11.0,9.0,13.0)

height <- c(66.0,68.0,64.5,65.0,70.0,64.0,70.0,71.0,72.0,
            64.0,74.5,67.0,71.0,71.0,77.0,72.0,59.0,62.0,72.0,66.0,64.0,
            73.0,69.0,72.0,71.0,69.0,70.0)

gender <- c("F","F","F","M","M","F","F","F","F","M","M","F","M","M",
            "M","F","F","M","M","F","F","M","M","M","M","M","M")

df <- data.frame(shoe, height, gender)
df
```

##	shoe	height	gender
## 1	6.5	66.0	F
## 2	9.0	68.0	F
## 3	8.5	64.5	F
## 4	8.5	65.0	M
## 5	10.5	70.0	M
## 6	7.0	64.0	F
## 7	9.5	70.0	F
## 8	9.0	71.0	F
## 9	13.0	72.0	F
## 10	7.5	64.0	M
## 11	10.5	74.5	M
## 12	8.5	67.0	F
## 13	12.0	71.0	M
## 14	10.5	71.0	M
## 15	13.0	77.0	M
## 16	11.5	72.0	F
## 17	8.5	59.0	F
## 18	5.0	62.0	M
## 19	10.0	72.0	M
## 20	6.5	66.0	F
## 21	8.5	64.0	F
## 22	10.5	73.0	M
## 23	6.5	69.0	M
## 24	10.5	72.0	M
## 25	11.0	71.0	M
## 26	9.0	69.0	M
## 27	13.0	70.0	M

*#a. Describe the data.*

*#The dataset contains 27 respondents, and each person has three variables  
#recorded: shoe size, height, and gender. Shoe size and height are numerical  
#variables, while gender is a categorical factor with two levels  
#(Male and Female). The data reflects basic anthropometric measurements  
#commonly used to compare physical characteristics across groups.*

*#b. Create a subset by males and females with their corresponding shoe size and height.  
#What its result? Show the R scripts.*

```
male_data <- subset(df, gender == "M", select = c(shoe, height, gender))  
female_data <- subset(df, gender == "F", select = c(shoe, height, gender))
```

male\_data

```
##      shoe height gender  
## 4      8.5   65.0      M  
## 5     10.5   70.0      M  
## 10     7.5   64.0      M  
## 11    10.5   74.5      M  
## 13    12.0   71.0      M  
## 14    10.5   71.0      M  
## 15    13.0   77.0      M  
## 18     5.0   62.0      M  
## 19    10.0   72.0      M  
## 22    10.5   73.0      M  
## 23     6.5   69.0      M  
## 24    10.5   72.0      M  
## 25    11.0   71.0      M  
## 26     9.0   69.0      M  
## 27    13.0   70.0      M
```

female\_data

```
##      shoe height gender  
## 1      6.5   66.0      F  
## 2      9.0   68.0      F  
## 3      8.5   64.5      F  
## 6      7.0   64.0      F  
## 7      9.5   70.0      F  
## 8      9.0   71.0      F  
## 9     13.0   72.0      F  
## 12     8.5   67.0      F  
## 16    11.5   72.0      F  
## 17     8.5   59.0      F  
## 20     6.5   66.0      F  
## 21     8.5   64.0      F
```

*#c. Find the mean of shoe size and height of the respondents. Write the R scripts and its  
#result.*

```
mean_shoe <- mean(df$shoe)
mean_height <- mean(df$height)
```

```
mean_shoe
```

```
## [1] 9.407407
```

```
mean_height
```

```
## [1] 68.66667
```

*#d. Is there a relationship between shoe size and height? Why?*

*#Yes. In general, larger shoe sizes tend to appear with greater heights in the dataset. This relationship is expected because shoe size is partly correlated with body proportions, and taller individuals typically have larger feet.*

*#While this dataset shows a pattern, a statistical method such as correlation analysis would confirm this.*

*#2. Construct character vector months to a factor with factor() and assign the result to factor\_months\_vector. Print out factor\_months\_vector and assert that R prints out the factor levels below the actual values.*

```
months_vector <- c(
  "March", "April", "January", "November", "January",
  "September", "October", "September", "November", "August",
  "January", "November", "November", "February", "May", "August",
  "July", "December", "August", "August", "September", "November", "February",
  "April"
)
factor_months_vector <- factor(months_vector)
factor_months_vector
```

```
## [1] March      April      January   November  January   September October
## [8] September November  August    January   November  November  February
## [15] May        August    July      December  August    August    September
## [22] November  February  April
## 11 Levels: April August December February January July March May ... September
```

*#3. Then check the summary() of the months\_vector and factor\_months\_vector. Interpret the results of both vectors. Are they both equally useful in this case?*

```
summary(months_vector)
```

```
##      Length      Class      Mode
##         24 character character
```

```
summary(factor_months_vector)
```

```
##      April      August  December  February  January      July      March      May
##         2         4         1         2         3         1         1         1
## November  October  September
##         5         1         3
```

*#No. The character vector simply stores text and cannot summarize categories.  
 #The factor version groups identical values and provides counts, which is more useful for analysis espe  
 #Therefore, the factor is much more useful when studying frequency, patterns, or performing statistical*

*#4. Create a vector and factor for the table below.*

```
factor_data <- c("East", rep("West", 4), rep("North", 3))
factor_data
```

```
## [1] "East" "West" "West" "West" "West" "North" "North" "North"
```

```
new_order_data <- factor(factor_data, levels = c("East", "West", "North"))
print(new_order_data)
```

```
## [1] East West West West West North North North
## Levels: East West North
```

*#5. Enter the data below in Excel with file name = import\_march.csv*

```
import_march <- read.table("import_march.csv", header = TRUE, sep = ",")
import_march
```

```
## Students Strategy.1 Strategy.2 Strategy.3
## 1 Male 8 10 8
## 2 4 8 6
## 3 0 6 4
## 4 Female 14 4 15
## 5 10 2 12
## 6 6 0 9
```

*#Using Conditional Statements (IF-ELSE)*

*#6. Full Search*

```
num <- as.integer(readline("Select a number from 1 to 50: "))
```

```
## Select a number from 1 to 50:
```

```
cat("Chosen number:", num, "\n")
```

```
## Chosen number: NA
```

```
if (is.na(num)) {
  cat("Invalid input. Please enter a number.\n")
} else if (num < 1 || num > 50) {
  cat("The number selected is beyond the range of 1 to 50\n")
} else if (num == 20) {
  print(TRUE)
} else {
  print(num)
}
```

```
## Invalid input. Please enter a number.
```

*#7. Change*

```
min_bills <- function(price) {  
  bills <- c(1000, 500, 200, 100, 50)  
  
  remaining <- price  
  count <- 0  
  
  for (b in bills) {  
    if (remaining >= b) {  
      count <- count + (remaining %/% b)  
      remaining <- remaining %% b  
    }  
  }  
  
  return(count)  
}
```

*#8. The following is each student's math score for one semester. Based on this, answer the following questions.*

*#a. Create a dataframe from the above table. Write the R codes and its output.*

```
students <- data.frame(  
  Name = c("Annie", "Thea", "Steve", "Hanna"),  
  Grade1 = c(85, 65, 75, 95),  
  Grade2 = c(65, 75, 55, 75),  
  Grade3 = c(85, 90, 80, 100),  
  Grade4 = c(100, 90, 85, 90)  
)
```

students

```
##      Name Grade1 Grade2 Grade3 Grade4  
## 1 Annie      85      65      85      100  
## 2 Thea       65      75      90       90  
## 3 Steve      75      55      80       85  
## 4 Hanna     95      75     100       90
```

*#b.*

```
for (i in 1:nrow(students)) {  
  avg <- (students$Grade1[i] + students$Grade2[i] +  
          students$Grade3[i] + students$Grade4[i]) / 4  
  
  if (avg > 90) {  
    cat(students$Name[i], "'s average grade this semester is ", avg, ".\n", sep="")  
  }  
}
```

*#c.*

```
tests <- students[, 2:5]
```

```

for (j in 1:4) {
  total <- sum(tests[, j])
  avg <- total / nrow(tests)

  if (avg < 80) {
    cat("The", j, "th test was difficult.\n")
  }
}

```

## The 2 th test was difficult.

```

#d
for (i in 1:nrow(students)) {
  grades <- students[i, 2:5]
  highest <- grades[1]

  for (g in grades) {
    if (g > highest) highest <- g
  }

  if (highest > 90) {
    cat(students$Name[i], "'s highest grade this semester is ", highest, ".\n", sep="")
  }
}

```

## Annie's highest grade this semester is 100.  
 ## Hanna's highest grade this semester is 100.