

RWorksheet_SOCO-3_B

*#1. Create a data frame using the table below.
#a. Write the codes.
#b. Describe the data. Get the structure or the summary of the data*

```
respondents <- c(1,2,3,4,5,6,7,8,9,10,
                 11,12,13,14,15,16,17,18,19,20)
sex <- c(2,2,1,2,2,2,2,2,2,2,
         1,2,2,2,2,2,2,2,1,2)
fathers_occupation <- c(1,3,3,3,1,2,3,1,1,1,
                       3,2,1,3,3,1,3,1,2,1)
persons_at_home <- c(5,7,3,8,5,9,6,7,8,4,
                     7,5,4,7,8,8,3,11,7,6)
siblings_at_school <- c(6,4,4,1,2,1,5,3,1,2,
                        3,2,5,5,2,1,2,5,3,2)
types_of_houses <- c(1,2,3,1,1,3,3,1,2,3,
                     2,3,2,2,3,3,3,3,3,2)

data <- data.frame(respondents, sex, fathers_occupation,
                   persons_at_home, siblings_at_school, types_of_houses)

data
```

	respondents	sex	fathers_occupation	persons_at_home	siblings_at_school
## 1	1	2	1	5	6
## 2	2	2	3	7	4
## 3	3	1	3	3	4
## 4	4	2	3	8	1
## 5	5	2	1	5	2
## 6	6	2	2	9	1
## 7	7	2	3	6	5
## 8	8	2	1	7	3
## 9	9	2	1	8	1
## 10	10	2	1	4	2
## 11	11	1	3	7	3
## 12	12	2	2	5	2
## 13	13	2	1	4	5
## 14	14	2	3	7	5
## 15	15	2	3	8	2
## 16	16	2	1	8	1
## 17	17	2	3	3	2
## 18	18	2	1	11	5
## 19	19	1	2	7	3
## 20	20	2	1	6	2
##	types_of_houses				
## 1	1				
## 2	2				
## 3	3				

```
## 4      1
## 5      1
## 6      3
## 7      3
## 8      1
## 9      2
## 10     3
## 11     2
## 12     3
## 13     2
## 14     2
## 15     3
## 16     3
## 17     3
## 18     3
## 19     3
## 20     2
```

```
#Getting the Structure and summary of the data.
str(data)
```

```
## 'data.frame':  20 obs. of  6 variables:
## $ respondents      : num  1 2 3 4 5 6 7 8 9 10 ...
## $ sex              : num  2 2 1 2 2 2 2 2 2 2 ...
## $ fathers_occupation: num  1 3 3 3 1 2 3 1 1 1 ...
## $ persons_at_home   : num  5 7 3 8 5 9 6 7 8 4 ...
## $ siblings_at_school: num  6 4 4 1 2 1 5 3 1 2 ...
## $ types_of_houses   : num  1 2 3 1 1 3 3 1 2 3 ...
```

```
summary(data)
```

```
##  respondents      sex      fathers_occupation persons_at_home
##  Min.   : 1.00    Min.   :1.00    Min.   :1.00      Min.   : 3.0
## 1st Qu.: 5.75    1st Qu.:2.00    1st Qu.:1.00      1st Qu.: 5.0
## Median :10.50    Median :2.00    Median :2.00      Median : 7.0
## Mean   :10.50    Mean   :1.85    Mean   :1.95      Mean   : 6.4
## 3rd Qu.:15.25    3rd Qu.:2.00    3rd Qu.:3.00      3rd Qu.: 8.0
## Max.   :20.00    Max.   :2.00    Max.   :3.00      Max.   :11.0
## siblings_at_school types_of_houses
##  Min.   :1.00      Min.   :1.0
## 1st Qu.:2.00      1st Qu.:2.0
## Median :2.50      Median :2.5
## Mean   :2.95      Mean   :2.3
## 3rd Qu.:4.25      3rd Qu.:3.0
## Max.   :6.00      Max.   :3.0
```

```
#c. Is the mean number of siblings attending is 5?
#No, since the mean based on the data is 2.95 as the display shown.
mean_siblings <- mean(data$siblings_at_school)
mean_siblings == 5
```

```
## [1] FALSE
```

```
mean_siblings
```

```
## [1] 2.95
```

```
#d. Extract the 1st two rows and all columns
```

```
data[1:2, ]
```

```
## respondents sex fathers_occupation persons_at_home siblings_at_school
## 1          1  2                   1             5             6
## 2          2  2                   3             7             4
## types_of_houses
## 1          1
## 2          2
```

```
#e. Extract 3rd and 5th rows with 2nd and 4th columns
```

```
data [c(3, 5), c(2, 4)]
```

```
## sex persons_at_home
## 3  1             3
## 5  2             5
```

```
#f. Select the variable types of houses then store the vector that results as types_houses. Write the c
types_houses <- data$types_of_houses
types_houses
```

```
## [1] 1 2 3 1 1 3 3 1 2 3 2 3 2 2 3 3 3 3 3 2
```

```
#g. Select only all Males respondent that their father occupation was farmer. Write the codes and its o
```

```
male_farmer <- subset(data, sex == 1 & fathers_occupation == 1)
male_farmer
```

```
## [1] respondents          sex          fathers_occupation persons_at_home
## [5] siblings_at_school types_of_houses
## <0 rows> (or 0-length row.names)
```

```
#h. Select only all females respondent that have greater than or equal to 5 number of siblings attendin
```

```
female_5plus <- subset(data, sex == 2 & siblings_at_school >= 5)
female_5plus
```

```
## respondents sex fathers_occupation persons_at_home siblings_at_school
## 1          1  2                   1             5             6
## 7          7  2                   3             6             5
## 13         13  2                   1             4             5
## 14         14  2                   3             7             5
## 18         18  2                   1            11             5
## types_of_houses
## 1          1
## 7          3
## 13         2
## 14         2
## 18         3
```