

Symbolic-Numeric Computing

1 Interval Arithmetic

- deciding the size of the working precision
- symbolic-numeric factorization

2 Constrained Optimization

- optimize subject to constraints
- Lagrange multipliers

MCS 320 Lecture 25
Introduction to Symbolic Computation
Jan Verschelde, 17 July 2026

Symbolic-Numeric Computing

1 Interval Arithmetic

- deciding the size of the working precision
- symbolic-numeric factorization

2 Constrained Optimization

- optimize subject to constraints
- Lagrange multipliers

How Much Precision is Needed for Accuracy?

Problem: Evaluate $f(x, y) =$

$$(333.75 - x^2)y^6 + x^2(11x^2y^2 - 121y^4 - 2) + 5.5y^8 + x/(2y)$$

at (77617, 33096).

An example of Stefano Taschini: [Interval Arithmetic: Python Implementation and Applications](#). In the *Proceedings of the 7th Python in Science Conference (SciPy 2008)*.

What are the problems?

- What must the precision be for accurate results?
- Increasing the precision often gives false confidence.
- How to compute bounds on the accuracy of an approximation?

Solution: compute with intervals $[a, b]$.

Symbolic-Numeric Computing

1 Interval Arithmetic

- deciding the size of the working precision
- symbolic-numeric factorization

2 Constrained Optimization

- optimize subject to constraints
- Lagrange multipliers

A Symbolic-Numeric Factorization

- A numeric factorization is over the complex numbers in $\mathbb{C}$.
- A symbolic factorization extends $\mathbb{Q}$ with algebraic numbers.

A *symbolic-numeric factorization* of a polynomial p uses complex interval arithmetic to represent the roots of p .

For example, in 10 bits of precision, $p = x^3 + x + 1$ factors as

$$(x - 0.341? - 1.17?I)(x - 0.341? + 1.17?I)(x + 0.682?),$$

where $I = \sqrt{-1}$ and $0.682?$ is short for $[0.68164, 0.68262]$.

Symbolic-Numeric Computing

1 Interval Arithmetic


- deciding the size of the working precision
- symbolic-numeric factorization

2 Constrained Optimization

- **optimize subject to constraints**
- Lagrange multipliers

Constrained Optimization

Find the point (x, y, z) on a sphere where $f(x, y, z)$ is optimal.



The constraint is the blue sphere, the target f is drawn in red.

Symbolic-Numeric Computing

1 Interval Arithmetic

- deciding the size of the working precision
- symbolic-numeric factorization

2 Constrained Optimization

- optimize subject to constraints
- Lagrange multipliers

Lagrange Multipliers

Let $f(x, y, z)$ be the target, what we want to optimize, and $g(x, y, z) = 0$ is the constraint.

At an optimal solution:

$$\nabla f \parallel \nabla g,$$

the gradients of f and g are parallel to each other.

The $\nabla f \parallel \nabla g$ is expressed via the multiplier λ .

Solve the polynomial system

$$\begin{cases} \nabla f(x, y, z) - \lambda \nabla g(x, y, z) = 0 \\ g(x, y, z) = 0. \end{cases}$$