

Analyysi I: Limits

Joona Piirainen

Autumn 2026

Abstract

Notes for the University of Helsinki course MAT11003, taught in Finnish from 01.08.2026–17.10.2026.

Contents

1	Course information	1
2	Lecture notes	2
2.1	01.09.2026	2
2.1.1	The absolute value function	2
2.2	03.09.2026	3
2.2.1	More on absolute values	3
2.2.2	Field and ordering properties of $\mathbb{R}$	4
2.3	08.09.2026	5
2.3.1	Sequences	5
2.3.2	Limit of a sequence	6
2.4	10.09.2026	7
2.4.1	Uniqueness of limits	7
2.4.2	Algebra of limits	8
2.4.3	Limits and order	10
2.4.4	Subsequences	11
2.5	15.09.2026	12
2.5.1	Upper and lower bounds	12
2.5.2	The completeness axiom	12
2.5.3	Maximum and properties of the supremum	13
2.5.4	Monotone sequences	13

1 Course information

The official course information is available on the University of Helsinki course page.

2 Lecture notes

2.1 01.09.2026

2.1.1 The absolute value function

Definition 2.1. For any real number $x \in \mathbb{R}$, the *absolute value* (or *modulus*) of x , denoted by $|x|$, is defined by

$$|x| = \begin{cases} x, & \text{if } x \geq 0, \\ -x, & \text{if } x < 0. \end{cases}$$

Geometrically, $|x - y|$ is the distance between the points x and y on the real line.

Example 2.2. Fix $y \in \mathbb{R}$ and let $h > 0$. The inequality $|x - y| < h$ can be written in several equivalent ways:

$$\begin{aligned} |x - y| < h &\iff -h < x - y < h \\ &\iff y - h < x < y + h \\ &\iff x \in (y - h, y + h). \end{aligned}$$

Thus, $|x - y| < h$ means that x lies in the open interval of radius h centred at y .

Example 2.3. As a concrete instance of the preceding equivalences,

$$\begin{aligned} |2x - 1| < 3 &\iff -3 < 2x - 1 < 3 \\ &\iff -2 < 2x < 4 \\ &\iff -1 < x < 2 \\ &\iff x \in (-1, 2). \end{aligned}$$

Theorem 2.4 (Triangle and reverse triangle inequalities). *For all $x, y \in \mathbb{R}$,*

$$\begin{aligned} |x + y| &\leq |x| + |y|, \\ ||x| - |y|| &\leq |x - y|. \end{aligned}$$

Proof. By definition 2.1,

$$\begin{aligned} -|x| &\leq x \leq |x|, \\ -|y| &\leq y \leq |y|. \end{aligned}$$

Adding the inequalities gives

$$-(|x| + |y|) \leq x + y \leq |x| + |y|,$$

which is equivalent to $|x + y| \leq |x| + |y|$.

Applying this triangle inequality to $x = (x - y) + y$ gives

$$|x| - |y| \leq |x - y|.$$

Interchanging x and y similarly gives $|y| - |x| \leq |x - y|$. Therefore

$$-|x - y| \leq |x| - |y| \leq |x - y|,$$

which is equivalent to the reverse triangle inequality. □

2.2 03.09.2026

2.2.1 More on absolute values

Theorem 2.5. *We have*

$$|xy| = |x||y|$$

for all $x, y \in \mathbb{R}$.

Proof. We prove by splitting into four cases. Firstly, if $x, y \geq 0$, we have $xy \geq 0$, so $|xy| = xy$. But in this case also both $|x| = x$ and $|y| = y$, so $|xy| = xy = |x||y|$. Secondly, if $x, y < 0$, we have $xy > 0$, $|xy| = xy = (-x)(-y) = |x||y|$. Now suppose $x < 0$ and $y \geq 0$. Then $xy \leq 0$, and hence $|xy| = -xy = (-x)y = |x||y|$. The case $x \geq 0, y < 0$ follows from commutativity of multiplication of real numbers. $\square$

Example 2.6. Let $x \in (1, 3)$ and $y \in (4, 8)$. Find an upper bound for

$$|x + y - 8|.$$

Kolmioepäyhtälön nojalla saadaan

$$\begin{aligned} |x + y - 8| &= |(x - 2) + (y - 6)| \\ &\leq |x - 2| + |y - 6| \\ &< 1 + 2 \\ &= 3. \end{aligned}$$

Thus we have found that 3 is an upper bound for $|x + y - 8|$.

Example 2.7. Let $x \in (2, 6)$ and $y \in (2, 8)$. Find an upper bound for

$$|xy - 20|.$$

We have $|x - 4| < 2$ and $|y - 5| < 3$. We have

$$\begin{aligned} |xy - 20| &= |xy - 4y + 4y - 20| \\ &\leq |xy - 4y| + |4y - 20| \\ &= |y||x - 4| + 4|y - 5| \\ &\leq 8 \cdot 2 + 4 \cdot 3 \\ &= 28. \end{aligned}$$

It follows that 28 is an upper bound for $|xy - 20|$.

Example 2.8. Find a positive real number m such that

$$|x^2 - 16| \leq m|x - 4|,$$

where $x \in (0, 10)$.

When $x = 4$, we get $0 \leq 0$, which holds with every $m \in \mathbb{R}$. Now suppose $x \neq 4$. We simplify the inequality and obtain

$$|x + 4| \leq m.$$

Since $x \in (0, 10)$, $|x + 4| \in (4, 14)$. We have thus obtained

$$|x + 4| \leq 14,$$

so we can choose $m = 14$.

2.2.2 Field and ordering properties of $\mathbb{R}$

Definition 2.9. $(F, +, \cdot)$ is a field if and only if

1. The algebraic structure $(F, +)$ is an *abelian group*.
2. The algebraic structure $(F^*, \cdot)$ is an *abelian group*, where $F^* = F \setminus \{0_F\}$.
3. The operation $\cdot$ distributes over $+$.

Theorem 2.10. $\mathbb{R}$ is a field.

Proof. Consider the usual addition and multiplication on $\mathbb{R}$.

First, $(\mathbb{R}, +)$ is an abelian group. Addition is closed, associative, and commutative; 0 is an additive identity; and every $x \in \mathbb{R}$ has the additive inverse $-x$.

Next, $(\mathbb{R} \setminus \{0\}, \cdot)$ is an abelian group. Multiplication is associative and commutative, and 1 is its identity element. If $x \neq 0$, then $x^{-1} = 1/x$ is a nonzero real number satisfying

$$x \cdot x^{-1} = 1.$$

The product of two nonzero real numbers is nonzero, so multiplication is closed on $\mathbb{R} \setminus \{0\}$.

Finally, multiplication distributes over addition:

$$x(y + z) = xy + xz$$

for all $x, y, z \in \mathbb{R}$. Therefore $(\mathbb{R}, +, \cdot)$ is a field. □

Definition 2.11 (Ordered field). An *ordered field* is a field $(F, +, \cdot)$ equipped with a relation $<$ such that, for all $x, y, z \in F$,

1. exactly one of $x < y$, $x = y$, or $x > y$ holds;
2. if $x < y$ and $y < z$, then $x < z$;
3. if $x < y$, then $x + z < y + z$; and
4. if $0_F < x$ and $0_F < y$, then $0_F < xy$.

Here $x > y$ means $y < x$. The first two conditions state that $<$ is a strict total order, while the last two state that the order is compatible with the field operations.

Theorem 2.12. The field $(\mathbb{R}, +, \cdot)$, equipped with the usual order $<$, is an ordered field.

Proof. We have already shown that $(\mathbb{R}, +, \cdot)$ is a field. The usual strict order on $\mathbb{R}$ satisfies trichotomy and transitivity, so it satisfies the first two ordering axioms.

Suppose that $x < y$. Then $0 < y - x$, and for every $z \in \mathbb{R}$,

$$(y + z) - (x + z) = y - x > 0.$$

Hence $x + z < y + z$. Furthermore, if $0 < x$ and $0 < y$, then the usual sign rule for multiplication gives $0 < xy$. Thus the order is compatible with addition and multiplication, and $\mathbb{R}$ is an ordered field. □

Theorem 2.13 (Uniqueness of additive identity). *The identity element 0_F of the abelian group $(F, +)$ is unique.*

Proof. Suppose that 0_F and $0'_F$ are both additive identity elements of $(F, +)$. Since 0_F is an additive identity,

$$0_F + 0'_F = 0'_F.$$

Since $0'_F$ is also an additive identity,

$$0_F + 0'_F = 0_F.$$

Therefore $0_F = 0'_F$. □

Theorem 2.14 (Cancellation law). *Let $x, y, z \in \mathbb{R}$. If $x + y = x + z$, then $y = z$.*

Proof. Using the additive identity, additive inverse, associativity, and the assumption $x + y = x + z$, we obtain

$$\begin{aligned} y &= 0 + y \\ &= ((-x) + x) + y \\ &= (-x) + (x + y) \\ &= (-x) + (x + z) \\ &= ((-x) + x) + z \\ &= 0 + z \\ &= z. \end{aligned}$$

□

2.3 08.09.2026

2.3.1 Sequences

Definition 2.15. Let X and Y be sets. A function $f: X \rightarrow Y$ is a relation $f \subseteq X \times Y$ such that, for every $x \in X$, there exists exactly one $y \in Y$ with $(x, y) \in f$. This unique y is denoted by $f(x)$.

The set X is the *domain* of f , and Y is its *codomain*. The set

$$f(X) = \{f(x) : x \in X\}$$

is its *range* or *image*.

Definition 2.16. A sequence of real numbers is a function $x: \mathbb{N} \rightarrow \mathbb{R}$. For $n \in \mathbb{N}$, the value $x(n)$, usually denoted by x_n , is the n -th term of the sequence. We denote the sequence by $(x_n)_{n \in \mathbb{N}}$, or simply by (x_n) .

2.3.2 Limit of a sequence

Definition 2.17. Let (x_n) be a sequence of real numbers. We say that (x_n) *converges* to $a \in \mathbb{R}$ if, for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that, for every $n \in \mathbb{N}$ with $n > N$,

$$|x_n - a| < \epsilon.$$

In this case, we write $x_n \rightarrow a$ as $n \rightarrow \infty$, or $\lim_{n \rightarrow \infty} x_n = a$.

Let us look at some examples.

Example 2.18. Let (x_n) be defined by

$$x_n = \frac{2n}{n+1}.$$

We will show that

$$\lim_{n \rightarrow \infty} x_n = 2.$$

Let $\epsilon > 0$. Choose $N \in \mathbb{N}$ such that $N > \frac{2}{\epsilon}$. Now for every $n > N$, we have

$$\begin{aligned} |x_n - 2| &= \left| \frac{2n}{n+1} - 2 \right| \\ &= \frac{2}{n+1} < \frac{2}{n} < \frac{2}{\frac{2}{\epsilon}} = \epsilon. \end{aligned}$$

Thus $x_n \rightarrow 2$ when $n \rightarrow \infty$.

Example 2.19. Let (x_n) be defined by

$$\begin{aligned} x_0 &= \cdots = x_7 = 42, \\ x_n &= 7 + \frac{1}{n}, \quad n \geq 8. \end{aligned}$$

We will show that

$$\lim_{n \rightarrow \infty} x_n = 7.$$

Let $\epsilon > 0$. Choose $N \in \mathbb{N}$ such that $N > \max \left\{ 8, \frac{1}{\epsilon} \right\}$. Now for every $n > N$, we have

$$|x_n - 7| = \frac{1}{n} < \epsilon.$$

Thus $x_n \rightarrow 7$ as $n \rightarrow \infty$.

Let us now tackle a simple divergence proof.

Example 2.20. Let us show that the sequence x_n defined by $x_n = n$ diverges. Assume that the sequence converged to some $a \in \mathbb{R}$. Taking $\epsilon = 1$, there exists $N \in \mathbb{N}$ such that, for every $n > N$,

$$|x_n - a| < \epsilon = 1.$$

Choose $n \in \mathbb{N}$ such that

$$n > \max \{N, |a| + 1\}.$$

By the triangle inequality,

$$n = |n| = |(n - a) + a| \leq |n - a| + |a|.$$

Subtracting $|a|$ from both sides and using $n > |a| + 1$, we obtain

$$|x_n - a| = |n - a| \geq n - |a| > 1,$$

contradicting the assumption $|x_n - a| < 1$. Therefore (x_n) diverges.

Example 2.21. Let (x_n) be defined by

$$\begin{aligned} x_0 &= \cdots = x_{42} = 1, \\ x_n &= \frac{n}{n - 42}, \quad n \geq 43. \end{aligned}$$

We will show that

$$\lim_{n \rightarrow \infty} x_n = 1.$$

Let $\epsilon > 0$. Choose $N \in \mathbb{N}$ such that

$$N > \max \left\{ 84, \frac{84}{\epsilon} \right\}.$$

Now, for every $n > N$, we have $n > 84$, and hence $\frac{n}{2} - 42 > 0$. Therefore

$$\begin{aligned} |x_n - 1| &= \left| \frac{n}{n - 42} - 1 \right| \\ &= \frac{42}{n - 42} \\ &= \frac{42}{\frac{n}{2} + \frac{n}{2} - 42} \\ &< \frac{84}{n} < \epsilon. \end{aligned}$$

Thus $x_n \rightarrow 1$ as $n \rightarrow \infty$.

2.4 10.09.2026

2.4.1 Uniqueness of limits

Theorem 2.22 (Uniqueness of limits). *A convergent sequence of real numbers has a unique limit. That is, if $x_n \rightarrow a$ and $x_n \rightarrow b$ as $n \rightarrow \infty$, then $a = b$.*

Proof. Suppose for contradiction that $a \neq b$, and set

$$\epsilon = \frac{|a - b|}{2} > 0.$$

Since $x_n \rightarrow a$ and $x_n \rightarrow b$, there exist $N_a, N_b \in \mathbb{N}$ such that

$$n > N_a \implies |x_n - a| < \epsilon$$

and

$$n > N_b \implies |x_n - b| < \epsilon.$$

Choose $n > \max \{N_a, N_b\}$. By the triangle inequality,

$$|a - b| \leq |a - x_n| + |x_n - b| < 2\epsilon = |a - b|,$$

which is impossible. Therefore $a = b$. $\square$

Theorem 2.23 (Convergent sequences are bounded). *Every convergent sequence of real numbers is bounded.*

Proof. Suppose that $x_n \rightarrow a$. Taking $\epsilon = 1$, there exists $N \in \mathbb{N}$ such that

$$n > N \implies |x_n - a| < 1.$$

By the triangle inequality, for every $n > N$,

$$|x_n| \leq |x_n - a| + |a| < 1 + |a|.$$

Let

$$M = \max \{|x_0|, |x_1|, \dots, |x_N|, 1 + |a|\}.$$

Then $|x_n| \leq M$ for every $n \in \mathbb{N}$, so (x_n) is bounded. $\square$

2.4.2 Algebra of limits

Theorem 2.24 (Algebraic limit laws). *Suppose that $x_n \rightarrow a$ and $y_n \rightarrow b$, and let $c \in \mathbb{R}$. Then*

1. $x_n \pm y_n \rightarrow a \pm b$.
2. $cx_n \rightarrow ca$.
3. $x_n y_n \rightarrow ab$.
4. If $b \neq 0$, then $y_n \neq 0$ for all sufficiently large n , and

$$\frac{x_n}{y_n} \rightarrow \frac{a}{b}.$$

Proof. 1. Let $\epsilon > 0$. Since $x_n \rightarrow a$ and $y_n \rightarrow b$, there exist $N_x, N_y \in \mathbb{N}$ such that

$$n > N_x \implies |x_n - a| < \frac{\epsilon}{2}$$

and

$$n > N_y \implies |y_n - b| < \frac{\epsilon}{2}.$$

Thus, for $n > \max \{N_x, N_y\}$,

$$|(x_n + y_n) - (a + b)| \leq |x_n - a| + |y_n - b| < \epsilon.$$

Hence $x_n + y_n \rightarrow a + b$. Replacing y_n and b by $-y_n$ and $-b$ gives $x_n - y_n \rightarrow a - b$.

2. If $c = 0$, then $cx_n = ca = 0$ for every n . If $c \neq 0$, choose $N \in \mathbb{N}$ such that

$$n > N \implies |x_n - a| < \frac{\epsilon}{|c|}.$$

Then, for every $n > N$,

$$|cx_n - ca| = |c||x_n - a| < \epsilon.$$

Therefore $cx_n \rightarrow ca$.

3. Since (y_n) converges, it is bounded. Choose $M > 0$ such that $|y_n| \leq M$ for every n . There exist $N_1, N_2 \in \mathbb{N}$ such that

$$n > N_1 \implies |x_n - a| < \frac{\epsilon}{2M}$$

and

$$n > N_2 \implies |y_n - b| < \frac{\epsilon}{2(|a| + 1)}.$$

For $n > \max\{N_1, N_2\}$,

$$\begin{aligned} |x_n y_n - ab| &\leq |x_n y_n - a y_n| + |a y_n - ab| \\ &= |y_n| |x_n - a| + |a| |y_n - b| \\ &< M \frac{\epsilon}{2M} + |a| \frac{\epsilon}{2(|a| + 1)} \\ &< \epsilon. \end{aligned}$$

Therefore $x_n y_n \rightarrow ab$.

4. Suppose that $b \neq 0$. Since $y_n \rightarrow b$, there exists $N_0 \in \mathbb{N}$ such that

$$n > N_0 \implies |y_n - b| < \frac{|b|}{2}.$$

The reverse triangle inequality then gives

$$|y_n| \geq |b| - |y_n - b| > \frac{|b|}{2} > 0$$

for every $n > N_0$. In particular, $y_n \neq 0$ eventually. Choose $N_1 \in \mathbb{N}$ such that

$$n > N_1 \implies |y_n - b| < \frac{\epsilon |b|^2}{2}.$$

For $n > \max\{N_0, N_1\}$,

$$\left| \frac{1}{y_n} - \frac{1}{b} \right| = \frac{|y_n - b|}{|y_n| |b|} < \frac{2}{|b|^2} |y_n - b| < \epsilon.$$

Hence $1/y_n \rightarrow 1/b$; the finitely many terms with $n \leq N_0$ may be defined arbitrarily because they do not affect the limit. Applying the product law to (x_n) and $(1/y_n)$ yields

$$\frac{x_n}{y_n} \rightarrow \frac{a}{b}.$$

□

2.4.3 Limits and order

Theorem 2.25. *Suppose that $x_n \rightarrow x$. Then*

1. *if $x_n \leq M$ for every $n \in \mathbb{N}$, then $x \leq M$;*
2. *if $x_n \geq m$ for every $n \in \mathbb{N}$, then $x \geq m$.*

Proof. 1. Suppose for contradiction that $x > M$, and set

$$\epsilon = \frac{x - M}{2} > 0.$$

Since $x_n \rightarrow x$, there exists $N \in \mathbb{N}$ such that, for every $n > N$,

$$|x_n - x| < \epsilon.$$

It follows that

$$x_n > x - \epsilon = x - \frac{x - M}{2} = \frac{x + M}{2} > M,$$

contradicting $x_n \leq M$. Therefore $x \leq M$.

2. Suppose for contradiction that $x < m$, and set

$$\epsilon = \frac{m - x}{2} > 0.$$

Since $x_n \rightarrow x$, there exists $N \in \mathbb{N}$ such that, for every $n > N$,

$$|x_n - x| < \epsilon.$$

It follows that

$$x_n < x + \epsilon = x + \frac{m - x}{2} = \frac{x + m}{2} < m,$$

contradicting $x_n \geq m$. Therefore $x \geq m$.

□

Theorem 2.26 (Squeeze theorem). *Suppose that*

$$x_n \leq y_n \leq z_n$$

for every $n \in \mathbb{N}$. If $x_n \rightarrow a$ and $z_n \rightarrow a$, then $y_n \rightarrow a$.

Proof. Let $\epsilon > 0$. Since $x_n \rightarrow a$, there exists $N_x \in \mathbb{N}$ such that

$$n > N_x \implies a - \epsilon < x_n < a + \epsilon.$$

Similarly, since $z_n \rightarrow a$, there exists $N_z \in \mathbb{N}$ such that

$$n > N_z \implies a - \epsilon < z_n < a + \epsilon.$$

Therefore, for every $n > \max\{N_x, N_z\}$,

$$a - \epsilon < x_n \leq y_n \leq z_n < a + \epsilon.$$

Hence $|y_n - a| < \epsilon$, so $y_n \rightarrow a$.

□

The inequalities in the squeeze theorem need only hold for all sufficiently large n , since changing finitely many terms does not affect convergence.

Example 2.27. Consider the oscillating sequence

$$x_n = \frac{(-1)^n}{n}, \quad n \geq 1.$$

Its sign alternates, but

$$-\frac{1}{n} \leq \frac{(-1)^n}{n} \leq \frac{1}{n}.$$

Both bounding sequences converge to 0. Indeed, given $\epsilon > 0$, choosing $N > 1/\epsilon$ gives $1/n < \epsilon$ whenever $n > N$. The squeeze theorem therefore gives

$$\lim_{n \rightarrow \infty} \frac{(-1)^n}{n} = 0.$$

Notice that the squeeze theorem lets us prove convergence without trying to handle even and odd indices separately.

Example 2.28. Consider

$$x_n = \sqrt{n^2 + n} - n, \quad n \geq 1.$$

Rationalising the expression gives

$$x_n = \frac{n}{\sqrt{n^2 + n} + n}.$$

Since

$$n < \sqrt{n^2 + n} < n + 1,$$

we obtain

$$\frac{n}{2n + 1} < x_n < \frac{1}{2}.$$

By the algebraic limit laws,

$$\lim_{n \rightarrow \infty} \frac{n}{2n + 1} = \lim_{n \rightarrow \infty} \frac{1}{2 + 1/n} = \frac{1}{2}.$$

The upper bound is the constant sequence $1/2$, so both bounds converge to $1/2$. Hence the squeeze theorem yields

$$\lim_{n \rightarrow \infty} (\sqrt{n^2 + n} - n) = \frac{1}{2}.$$

2.4.4 Subsequences

Definition 2.29. Let (x_n) be a sequence, and let (n_k) be a strictly increasing sequence of natural numbers:

$$n_0 < n_1 < n_2 < \cdots.$$

The sequence (x_{n_k}) , obtained by selecting the terms of (x_n) whose indices are $n_0, n_1, n_2, \dots$, is called a *subsequence* of (x_n) .

Theorem 2.30. *If $x_n \rightarrow x$, then every subsequence of (x_n) converges to x .*

Proof. Let (x_{n_k}) be a subsequence of (x_n) , and let $\epsilon > 0$. Since $x_n \rightarrow x$, there exists $N \in \mathbb{N}$ such that

$$n > N \implies |x_n - x| < \epsilon.$$

Because (n_k) is strictly increasing, $n_k \geq k$ for every $k \in \mathbb{N}$. Thus, whenever $k > N$, we also have $n_k > N$, and hence

$$|x_{n_k} - x| < \epsilon.$$

Therefore $x_{n_k} \rightarrow x$. Since the subsequence was arbitrary, every subsequence of (x_n) converges to the same limit x . $\square$

2.5 15.09.2026

2.5.1 Upper and lower bounds

Definition 2.31. Let $\emptyset \neq E \subseteq \mathbb{R}$. A number $s \in \mathbb{R}$ is an *upper bound* of E if

$$x \leq s \quad \text{for every } x \in E.$$

If such a number exists, E is *bounded above*.

Definition 2.32. Let $\emptyset \neq E \subseteq \mathbb{R}$. A number $p \in \mathbb{R}$ is a *lower bound* of E if

$$p \leq x \quad \text{for every } x \in E.$$

If such a number exists, E is *bounded below*.

Definition 2.33. Let $\emptyset \neq E \subseteq \mathbb{R}$ be bounded above. An upper bound s of E is the *least upper bound*, or *supremum*, of E if $s \leq u$ for every upper bound u of E . We write

$$s = \sup E.$$

Definition 2.34. Let $\emptyset \neq E \subseteq \mathbb{R}$ be bounded below. A lower bound p of E is the *greatest lower bound*, or *infimum*, of E if $p \geq \ell$ for every lower bound ℓ of E . We write

$$p = \inf E.$$

If E is not bounded above, we write $\sup E = +\infty$. Similarly, if E is not bounded below, we write $\inf E = -\infty$.

2.5.2 The completeness axiom

Completeness axiom. Every nonempty subset of $\mathbb{R}$ that is bounded above has a supremum in $\mathbb{R}$.

Corollary 2.35. *Every nonempty subset $E \subseteq \mathbb{R}$ that is bounded below has an infimum in $\mathbb{R}$.*

Proof. Let c be a lower bound of E , and define

$$F = \{-x : x \in E\}.$$

Since $c \leq x$ for every $x \in E$, we have $-x \leq -c$. Thus F is bounded above. It is also nonempty, so the completeness axiom gives $\sup F \in \mathbb{R}$. For every $x \in E$,

$$-x \leq \sup F \leq -c, \quad \text{hence} \quad x \geq -\sup F \geq c.$$

Therefore $-\sup F$ is a lower bound of E . Since the argument applies to every lower bound c of E , it is the greatest lower bound. Consequently,

$$\inf E = -\sup F \in \mathbb{R}.$$

□

2.5.3 Maximum and properties of the supremum

Definition 2.36. A number s is the *greatest element*, or *maximum*, of a set E if $s \in E$ and $x \leq s$ for every $x \in E$. We write $s = \max E$.

Theorem 2.37. Let $\emptyset \neq E \subseteq \mathbb{R}$ be bounded above. If $s = \max E$, then $s = \sup E$.

Proof. Since $x \leq s$ for every $x \in E$, the number s is an upper bound of E . Hence $\sup E \leq s$. On the other hand, $s \in E$, so $s \leq \sup E$. Thus $s = \sup E$. □

Theorem 2.38. The supremum of a nonempty subset of $\mathbb{R}$ that is bounded above is unique.

Proof. Suppose that s and s' are both least upper bounds of E . Since s' is an upper bound and s is a least upper bound, $s \leq s'$. Similarly, since s is an upper bound and s' is a least upper bound, $s' \leq s$. Therefore $s = s'$. □

Theorem 2.39. Let $\emptyset \neq E \subseteq \mathbb{R}$ be bounded above. For every $\epsilon > 0$, there exists $x \in E$ such that

$$\sup E - \epsilon < x \leq \sup E.$$

Proof. Since $\sup E$ is an upper bound, $x \leq \sup E$ for every $x \in E$. Suppose, for a contradiction, that for some $\epsilon > 0$,

$$x \leq \sup E - \epsilon \quad \text{for every } x \in E.$$

Then $\sup E - \epsilon$ is an upper bound of E , but $\sup E - \epsilon < \sup E$. This contradicts the fact that $\sup E$ is the least upper bound. Hence the claimed element x exists. □

2.5.4 Monotone sequences

Definition 2.40. A real sequence (x_n) is

- *increasing* (nondecreasing) if $x_n \leq x_{n+1}$ for every $n \in \mathbb{N}$;
- *decreasing* (nonincreasing) if $x_{n+1} \leq x_n$ for every $n \in \mathbb{N}$;
- *strictly increasing* if $x_n < x_{n+1}$ for every $n \in \mathbb{N}$;
- *strictly decreasing* if $x_{n+1} < x_n$ for every $n \in \mathbb{N}$.

A sequence is *monotone* if it is increasing or decreasing, and *strictly monotone* if it is strictly increasing or strictly decreasing.