

1 Lecture 1 May 02 2018

1.1 Introduction

1.1.1 Numbers

The following are some of the number sets that we are already familiar with:

$$\begin{aligned}\mathbb{N} &= \{1, 2, 3, \dots\} & \mathbb{Z} &= \{\dots, -2, -1, 0, 1, 2, \dots\} \\ \mathbb{Q} &= \left\{ \frac{a}{b} : a \in \mathbb{Z}, b \in \mathbb{N} \right\} & \mathbb{R} &= \text{set of real numbers} \\ \mathbb{C} &= \{a + bi : a, b \in \mathbb{R}, i = \sqrt{-1}\} = \text{set of complex numbers}\end{aligned}$$

For $n \in \mathbb{Z}$, let \mathbb{Z}_n denote the set of integers modulo n , i.e.

$$\mathbb{Z}_n = \{[0], [1], \dots, [n-1]\}$$

where the $[r]$, $0 \leq r \leq n-1$, are the congruence classes, i.e.

$$[r] = \{z \in \mathbb{Z} : z \equiv r \pmod{n}\}$$

These sets share some common properties, e.g. $+$ and \times . Let's try to break that down to make further observation.

NOTE THAT for $R = \mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$, or \mathbb{Z}_n , R has 2 operations, i.e. addition and multiplication.

Addition If $r_1, r_2, r_3 \in R$, then

- **(closure)** $r_1 + r_2 \in R$
- **(associativity)** $r_1 + (r_2 + r_3) = (r_1 + r_2) + r_3$

Also, if $R \neq \mathbb{N}$, then $\exists 0 \in R$ (the **additive identity**) such that

$$\forall r \in R \quad r + 0 = r = 0 + r.$$

Also, $\forall r \in R$, $\exists(-r) \in R$ such that

$$r + (-r) = 0 = (-r) + r.$$

Multiplication For $r_1, r_2, r_3 \in R$, we have

- (**closure**) $r_1 r_2 \in R$
- (**associativity**) $r_1(r_2 r_3) = (r_1 r_2)r_3$

Also, $\exists 1 \in R$ (a.k.a the **multiplicative identity**), such that

$$\forall r \in R \quad r \cdot 1 = r = 1 \cdot r.$$

Finally, for $R = \mathbb{Q}$, \mathbb{R} , or \mathbb{C} , $\forall r \in R$, $\exists r^{-1} \in R$ such that

$$r \cdot r^{-1} = 1 = r^{-1} \cdot r.$$

Note that for $R = \mathbb{Z}_n$, where $n \in \mathbb{Z}$, not all $[r] \in \mathbb{Z}_n$ have a multiplicative inverse. For example, for $[2] \in \mathbb{Z}_4$, there is no $[x] \in \mathbb{Z}_4$ such that $[2][x] = [1]$.¹

¹ This is best proven using techniques introduced in MATH135/145.

1.1.2 Matrices

For $n \in \mathbb{N} \setminus \{1\}$, an $n \times n$ matrix over \mathbb{R} ² is an $n \times n$ array that can be expressed as follows:

$$A = [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

where for $1 \leq i, j \leq n$, $a_{ij} \in \mathbb{R}$. We denote $M_n(\mathbb{R})$ as the set of all $n \times n$ matrices over \mathbb{R} .

As in Section 1.1.1, we can perform **addition and multiplication** on $M_n(\mathbb{R})$.

² \mathbb{R} can be replaced by \mathbb{Q} or \mathbb{C} .

Define the identity matrix, $I \in M_n(\mathbb{R})$, as follows:

$$I = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix}.$$

Then we have that I is the **multiplicative identity**, since

$$AI = A = IA.$$

However, contrary to matrix addition, $\forall A \in M_n(\mathbb{R})$, it is not always true that $\exists A^{-1} \in M_n(\mathbb{R})$ such that

$$AA^{-1} = I = A^{-1}A.$$

This is especially true if the **determinant** of A is 0.

Also, we can always find some $A, B \in M_n(\mathbb{R})$ such that

$$AB \neq BA,$$

i.e. matrix multiplication is not always commutative.

THE COMMON PROPERTIES of the operations from above: **closure, associativity, and existence of an inverse**, are not unique to just addition and multiplication. We shall see in the next lecture that there are other operations where these properties will continue to hold, e.g. **permutations**.

2 *Index*

additive identity, 10
associativity, 9

closure, 9

mutiplicative identity, 10