

UW W17 PMATH333 - Definitions and Theorems

Johnson Ng

April 15, 2017

Preface

PMATH333 is offered as a course that attempts to bridge the gap for students who have taken the regular math courses instead of the advanced math courses, in particular for MATH147, MATH148 and MATH247 in UW. This set of notes is taken from the W2017 term.

Contents

1	The Real Number System	13
1.1	Ordered Field Axioms	13
1.2	Completeness Axiom	17
2	Sequences	20
2.1	Limits of Sequences	20
2.2	Limit Theorems	22
2.3	Bolzano-Weierstrass Theorem	24
2.4	Cauchy Sequences	25
3	Functions on \mathbb{R}	26
3.1	Two-Sided Limits	26
3.2	Continuity	30
4	Differentiability on \mathbb{R}	33
4.1	The Derivative	33
4.2	Differentiability Theorems	34
4.3	Inverse Function Theorems	35
4.4	Mean Value Theorem	36
4.5	l'Hôpital's Rule	38
5	Integrability on \mathbb{R}	39
5.1	The Riemann Integral	39
5.2	Upper and Lower Riemann Sums	40
5.3	Evaluating Integrals of Continuous Functions	42
5.4	Basic Properties of Integrals	42
5.5	Fundamental Theorem of Calculus	43
6	Series of \mathbb{R}	44
6.1	Introduction	44

<i>CONTENTS</i>	3
6.2 Convergence Tests	45
6.3 Fubini's Theorem for Series	47
7 Sequences and Series of Functions	49
7.1 Pointwise Convergence	49
7.2 Uniform Convergence	49
7.3 Series of Functions	51
7.4 Power Series	52
7.5 Operations on Power Series	53
7.6 Taylor Series	55
8 Topology in Euclidean Space	56
A ZF Set Theory and the Axiom of Choice	57
A.1 Introduction	57
A.2 ZFC Axioms of Set Theory	58
A.3 Relations, Equivalence Relations, Functions and Recursion	64
A.4 Construction of Integers, Rational, Real and Complex Numbers	68
B Functions and Cardinality	70
B.1 Functions	70
B.2 Cardinality	71

List of Definitions

Definition 1.1.1	Removal	13
Definition 1.1.2	Disjoint	13
Definition 1.1.3	Intervals	13
Definition 1.1.4	Ring	14
Definition 1.1.5	Commutative Ring	14
Definition 1.1.6	Field	14
Definition 1.1.7	Order	15
Definition 1.1.8	Ordered Field	16
Definition 1.1.9	Absolute Value	16
Definition 1.2.1	Upper and Lower Bounds	17
Definition 1.2.2	Supremum and Infimum	17
Definition 1.2.3	Floor and Ceiling Functions	18
Definition 2.1.1	Sequence	20
Definition 2.1.2	Subsequence	20
Definition 2.1.3	Extended Ordered Field	20
Definition 2.1.4	Convergence, Divergence and Limits of a Sequence	21
Definition 2.2.1	Bounds	24
Definition 2.3.1	Increasing, Decreasing, and Monotonic Sequences	24
Definition 2.3.2	Rearrangement of a Sequence	25
Definition 2.4.1	Cauchy	25

Definition 3.1.1	Limit Point	26
Definition 3.1.2	Limit Point from Above and Below	26
Definition 3.1.3	Infinity As A Limit Point	27
Definition 3.2.1	Continuity	30
Definition 3.2.2	Limit Point and Continuity	30
Definition 3.2.3	Maximum, Minimum and Extreme Values	31
Definition 3.2.4	Uniform Continuity	32
Definition 4.1.1	Differentiable on a Point	33
Definition 4.1.2	Differentiable in a Domain	33
Definition 4.1.3	Differentiable n-times & n-th Derivative	34
Definition 4.1.4	Linearization & Tangent Line	34
Definition 4.4.1	Local Maximum and Minimum	36
Definition 5.1.1	Partition & Subintervals	39
Definition 5.1.2	The Riemann Sum	39
Definition 5.1.3	(Riemann) Integrable	40
Definition 5.2.1	Upper and Lower Riemann Sums	40
Definition 5.2.2	Upper and Lower Integral	41
Definition 5.4.1	Integral at a Point & Integral from the Right	43
Definition 5.5.1	Antiderivative	43
Definition 6.1.1	Series	44
Definition 6.1.2	Error	45
Definition 6.2.1	Limit Supremum	46
Definition 6.2.2	Alternating Sequence	47
Definition 6.2.3	Converge Absolutely & Converge Conditionally	47
Definition 6.3.1	Multiplication of Series	47
Definition 7.1.1	Pointwise Convergence of Sequences of Functions	49
Definition 7.2.1	Uniform Convergence of Sequences of Functions	49

Definition 7.3.1	Series of Functions	51
Definition 7.4.1	Power Series	52
Definition 7.4.2	Interval and Radius of Convergence	53
Definition 7.6.1	Taylor Series and Polynomial	55
Definition A.2.1	Mathematical Symbols	58
Definition A.2.2	Formula	58
Definition A.2.3	Free or Bounded Variable	59
Definition A.2.4	Is Bound By and Binds	59
Definition A.2.5	Free Variable, Statement, Statement About	59
Definition A.2.6	Unique Existence	59
Definition A.2.7	Empty Set Axiom	60
Definition A.2.8	Extension Axiom	60
Definition A.2.9	\emptyset	60
Definition A.2.10	Subset	61
Definition A.2.11	Separation Axiom	61
Definition A.2.12	Pair Axiom	61
Definition A.2.13	Union Axiom	61
Definition A.2.14	Union	61
Definition A.2.15	Intersection	62
Definition A.2.16	Power Set Axiom	62
Definition A.2.17	Power Set	62
Definition A.2.18	Ordered Pair	62
Definition A.2.19	Successor, Inductive	63
Definition A.2.20	Axiom of Infinity	63
Definition A.2.21	Natural Numbers	63
Definition A.2.22	Replacement Axiom	64
Definition A.2.23	Axiom of Choice	64

Definition A.3.1	Binary Relation	64
Definition A.3.2	Domain, Range, Image, Inverse Image, Inverse, Composition .	64
Definition A.3.3	Equivalence Relation	65
Definition A.3.4	Equivalence Class	65
Definition A.3.5	Partition	65
Definition A.3.6	Set of Representatives	66
Definition A.3.7	Function	66
Definition A.3.8	One-to-one & Onto	67
Definition A.3.9	Left and Right Inverses	67
Definition A.3.10	Invertible	67
Definition A.4.1	Sum and Product	68
Definition A.4.2	Integers	68
Definition A.4.3	Rational Numbers	68
Definition A.4.4	Real Numbers	69
Definition A.4.5	Complex Numbers	69
Definition B.1.1	Range, Image, and Inverse Image	70
Definition B.1.2	Composite Function	70
Definition B.1.3	Bijection	70
Definition B.1.4	Identity Function	71
Definition B.2.1	Equal Cardinality	71
Definition B.2.2	Properties for Cardinality of Sets	71
Definition B.2.3	Finiteness and Countability of Sets	72
Definition B.2.4	Countability and \aleph_0	73

List of Theorems

Theorem 1.1.1	Properties of Sets	13
Theorem 1.1.2	\mathbb{Q} and \mathbb{R} as Fields	14
Theorem 1.1.3	Cancellations & Identities	15
Theorem 1.1.4	Properties of Fields	15
Theorem 1.1.5	$\mathbb{N}, \mathbb{Z}, \mathbb{Q}$ and \mathbb{R} are ordered	16
Theorem 1.1.6	\mathbb{Q} and \mathbb{R} as Ordered Fields	16
Theorem 1.1.7	Properties of Ordered Fields	16
Theorem 1.1.8	Properties of Absolute Values	17
Theorem 1.1.9	Basic Order Properties in \mathbb{Z}	17
Theorem 1.2.1	Approximation Property of Supremum and Infimum	18
Theorem 1.2.2	Completeness Properties of \mathbb{R}	18
Theorem 1.2.3	Well-Ordering Properties of \mathbb{Z} in \mathbb{R}	18
Theorem 1.2.4	Floor and Ceiling Properties of \mathbb{Z} in \mathbb{R}	18
Theorem 1.2.5	Archimedean Properties of \mathbb{Z} in \mathbb{R}	18
Theorem 1.2.6	Density of \mathbb{Q}	19
Theorem 2.1.1	Independence of Limit from Initial Terms	21
Theorem 2.1.2	Uniqueness of Limit	22
Theorem 2.2.1	Basic Limits	22
Theorem 2.2.2	Operations on Limits	22
Theorem 2.2.3	Extended Operations on Limits	22

Theorem 2.2.4	Monotonic Surjective Functions	23
Theorem 2.2.5	Basic Elementary functions Acting on Limits	23
Theorem 2.2.6	Comparison Theorem for Sequences	24
Theorem 2.2.7	Squeeze Theorem for Sequences	24
Theorem 2.3.1	Monotone Convergence Theorem	24
Theorem 2.3.2	Nested Interval Theorem	25
Theorem 2.3.3	Convergence of Subsequences and Rearrangements	25
Theorem 2.3.4	Bolzano-Weierstrass Theorem	25
Theorem 2.4.1	Cauchy Criterion for Convergence	25
Theorem 3.1.1	Two-sided Limits	28
Theorem 3.1.2	Sequential Characterization of Limits of Functions	28
Theorem 3.1.3	Local Determination of Limits	28
Theorem 3.1.4	Uniqueness of Limits	28
Theorem 3.1.5	Extended Operations on Limits	29
Theorem 3.1.6	Basic Elementary Functions Acting on Limits	29
Theorem 3.1.7	Comparison Theorem for Functions	30
Theorem 3.1.8	Squeeze Theorem for Functions	30
Theorem 3.2.1	Sequential Characterization of Continuity	31
Theorem 3.2.2	Operations on Continuous Functions	31
Theorem 3.2.3	Composition of Continuous Functions	31
Corollary 3.2.3.1	Continuity of Elementary Functions	31
Theorem 3.2.4	Functions Acting on Limits	31
Theorem 3.2.5	Intermediate Value Theorem	31
Theorem 3.2.6	Extreme Value Theorem	32
Theorem 3.2.7	Closed Bounded Intervals and Uniform Continuity	32
Theorem 4.1.1	Definition of Differentiability in ϵ - δ	34
Theorem 4.1.2	Differentiability \implies Continuity	34

<i>CONTENTS</i>		10
Theorem 4.2.1	Local Determination of the Derivative	34
Theorem 4.2.2	Operations on Derivatives	35
Theorem 4.2.3	Chain Rule	35
Theorem 4.3.1	Monotonic Functions	35
Theorem 4.3.2	Continuity and Strictly Monotonous Functions	35
Theorem 4.3.3	Inverse Function Theorem	36
Theorem 4.3.4	Derivatives of the Basic Elementary Functions	36
Theorem 4.4.1	Fermat's Theorem	36
Theorem 4.4.2	Rolle's Theorem	37
Theorem 4.4.3	Mean Value Theorem	37
Theorem 4.4.4	Cauchy/Generalized Mean Value Theorem	37
Corollary 4.4.4.1	Trichotomy of The Derivative	37
Corollary 4.4.4.2	The Second Derivative Test	37
Theorem 4.5.1	l'Hôpital's Rule	38
Theorem 5.2.1	Upper and Lower Riemann Sums and the Riemann Sum	41
Theorem 5.2.2	Partition Refinement	41
Theorem 5.2.3	Upper Integral \geq Lower Integral	41
Theorem 5.2.4	Equivalent Definitions of Integrability	41
Theorem 5.3.1	Continuous Functions are Integrable	42
Lemma 5.3.2	Summation Formulas	42
Theorem 5.4.1	Linearity	42
Theorem 5.4.2	Comparison Theorem for Integrals	42
Theorem 5.4.3	Additivity	42
Theorem 5.4.4	Estimation	43
Theorem 5.5.1	The Fundamental Theorem of Calculus	43
Theorem 6.1.1	First Finitely Many Terms Do Not Affect Convergence	44
Theorem 6.1.2	Linearity	45

<i>CONTENTS</i>	11
Theorem 6.1.3	Series of Positive Terms 45
Theorem 6.1.4	Cauchy Criterion 45
Theorem 6.2.1	Divergence Test 45
Theorem 6.2.2	Integral Test 46
Corollary 6.2.2.1	p-Series 46
Theorem 6.2.3	Comparison Test 46
Theorem 6.2.4	Limit Comparison Test 46
Theorem 6.2.5	Ratio Test 46
Theorem 6.2.6	Root Test 46
Theorem 6.2.7	Alternating Series Test 47
Theorem 6.2.8	Absolute Converges \implies Convergence 47
Theorem 6.3.1	Fubini's Theorem for Series 47
Theorem 7.2.1	Cauchy Criterion for Uniform Convergence - Sequence of Functions 50
Theorem 7.2.2	Uniform Convergence, Limits and Continuity - Sequence of Functions 50
Theorem 7.2.3	Uniform Convergence and Integration - Sequence of Functions 50
Theorem 7.2.4	Uniform Convergence and Differentiation - Sequence of Functions 50
Theorem 7.3.1	Cauchy Criterion for Uniform Convergence - Series of Functions 51
Theorem 7.3.2	Uniform Convergence, Limits and Continuity - Series of Functions 51
Theorem 7.3.3	Term-by-Term Integration 51
Theorem 7.3.4	Term-by-Term Differentiation 51
Theorem 7.3.5	Weierstrass M-Test 52
Lemma 7.4.1	Abel's Formula 52
Theorem 7.4.2	The Interval and Radius of Convergence 52
Theorem 7.4.3	Abel's Theorem 52
Theorem 7.5.1	Continuity of Power Series 53
Theorem 7.5.2	Addition and Subtraction of Power Series 53

<i>CONTENTS</i>	12
Theorem 7.5.3	Multiplication of Power Series 53
Theorem 7.5.4	Division of Power Series 53
Theorem 7.5.5	Composition of Power Series 54
Theorem 7.5.6	Integration of Power Series 54
Theorem 7.5.7	Differentiation of Power Series 54
Theorem 7.6.1	Coefficients of the Taylor Series 55
Theorem 7.6.2	Taylor's Theorem 55
Theorem 7.6.3	(?) Accuracy of the Maclaurin Expansion 55
Theorem A.2.1	Uniqueness of the Empty Set 60
Theorem A.2.2	Existence & Uniqueness of an Inductive Set 63
Theorem A.2.3	Principle of Induction 63
Theorem A.3.1	Domain, Range, Image and Inverse Image as Sets 65
Theorem A.3.2	Inverse and Composition as Binary Relations 65
Theorem A.3.3	Correspondence of Equivalence Relations and Partitions 66
Theorem A.3.4	Surjective and Injective VS Inverses 67
Theorem A.3.5	The Recursion Theorem 67
Theorem B.1.1	Bijectiveness and Inverse of the Composite Function 70
Theorem B.1.2	Bijectiveness and Invertability of Functions 71
Corollary B.1.2.1	Relationship between Injection and Surjection 71
Theorem B.2.1 72
Corollary B.2.1.1 72
Theorem B.2.2	$ \mathbb{N} $ as a Threshold for Finiteness and Countability 72
Theorem B.2.3 72
Theorem B.2.4	Set Cartesian Product and Union, and \mathbb{Q} are Countable 73
Theorem B.2.5	\mathbb{R} as an Uncountable Set 73
Theorem B.2.6	Cantor-Schröder-Bernstein Theorem 73

Chapter 1

The Real Number System

1.1 Ordered Field Axioms

Please review [Appendix A](#). We shall use all of the set notations that are introduced in Appendix A. We will also introduce one more notation.

Definition 1.1.1 (Removal)

Let A and B be sets. The set A remove B , denoted as $A \setminus B$, is the set

$$A \setminus B = \{x | x \in A \wedge x \notin B\}$$

Definition 1.1.2 (Disjoint)

Let A and B be sets. We say that A and B are disjoint when $A \cap B = \emptyset$

Theorem 1.1.1 (Properties of Sets)

Let $A, B, C \subseteq X$. Then

1. (Idempotence) $A \cup A = A, A \cap A = A$
2. (Identity) $A \cup \emptyset = A, A \cap \emptyset = \emptyset, A \cup X = X, A \cap X = A$
3. (Associativity) $(A \cup B) \cup C = A \cup (B \cup C)$ and $(A \cap B) \cap C = A \cap (B \cap C)$
4. (Commutativity) $A \cup B = B \cup A$ and $A \cap B = B \cap A$
5. (Distributivity) $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ and $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
6. (De Morgan's Laws) $X \setminus (A \cup B) = (X \setminus A) \cap (X \setminus B)$ and $X \setminus (A \cap B) = (X \setminus A) \cup (X \setminus B)$

Definition 1.1.3 (Intervals)

For $a, b \in \mathbb{R}$ with $a \leq b$ we write

$$\begin{aligned}
(a, b) &= \{x \in \mathbb{R} | a < x < b\}, & [a, b] &= \{x \in \mathbb{R} | a \leq x \leq b\}, \\
(a, b] &= \{x \in \mathbb{R} | a < x \leq b\}, & [a, b) &= \{x \in \mathbb{R} | a \leq x < b\}, \\
(a, \infty) &= \{x \in \mathbb{R} | a < x\}, & [a, \infty) &= \{x \in \mathbb{R} | a \leq x\}, \\
(-\infty, b) &= \{x \in \mathbb{R} | x < b\}, & (-\infty, b] &= \{x \in \mathbb{R} | x \leq b\}, \\
(-\infty, \infty) &= \mathbb{R}
\end{aligned}$$

An interval in \mathbb{R} is any set of one of the above forms. In the case that $a = b$, we have $(a, b) = [a, b) = (a, b] = \emptyset$ and $[a, b] = \{a\}$, and these intervals are called **degenerate** intervals. The intervals $\emptyset, (a, b), (a, \infty), (-\infty, b)$ and (∞, ∞) are called open intervals. The intervals $\emptyset, [a, b], [a, \infty), (-\infty, b]$ and $(-\infty, \infty)$ are called closed intervals.

Remark

Note on how the intervals \emptyset and $(-\infty, \infty)$ are both open and closed intervals.

Definition 1.1.4 (Ring)

A ring is a set F with two distinct elements $0, 1 \in F$ and two binary operations $+$ and \cdot such that

1. (Additive Associativity) For all $x, y, z \in F$ we have $(x + y) + z = x + (y + z)$,
2. (Additive Commutativity) For all $x, y \in F$ we have $x + y = y + x$,
3. (Additive Identity) For all $x \in F$ we have $0 + x = x$.
4. (Additive Inverse) $\forall x \in F \exists !y \in F$ $x + y = 0$
5. (Multiplicative Associativity) $\forall x, y, z \in F$ we have $(x \cdot y) \cdot z = x \cdot (y \cdot z)$,
6. (Multiplicative Identity) $\forall x \in F$ we have $1 \cdot x = x = x \cdot 1$,
7. (Distributivity) $\forall x, y, z \in F$ we have $x \cdot (y + z) = (x \cdot y) + (x \cdot z)$

Definition 1.1.5 (Commutative Ring)

A ring F is a commutative ring if it has the following (additional) property:
(Multiplicative Commutativity) $\forall x, y \in F$ we have $x \cdot y = y \cdot x$.

Definition 1.1.6 (Field)

A commutative ring F is a field if it has the following (additional) property:
(Multiplicative Inverse) $\forall x \neq 0 \in F \exists !y \in F$ such that $x \cdot y = 1$.

Remark

For the sake of simplicity, we will write $x \cdot y = xy$ for any x and y .

Theorem 1.1.2 (\mathbb{Q} and \mathbb{R} as Fields)

\mathbb{Q} and \mathbb{R} are fields.

Remark

Note that \mathbb{Z} and \mathbb{N} are not fields since their elements do not have a multiplicative inverse. They are, however, commutative rings.

Remark (Some shorthand notations)

Let F be a field and let $a, b \in F$. We denote the unique additive inverse of a by $-a$ and we write $a - b = a + (-b)$. When $a \neq 0$, we denote the unique multiplicative inverse of a by a^{-1} and we write $b \div a = \frac{b}{a} = ba^{-1}$.

Theorem 1.1.3 (Cancellations & Identities)

Let F be a field. Then $\forall x, y, z \in F$ we have

1. (Additive Cancellation) $x + y = x + z \implies x = z$
2. (Uniqueness of Additive Identity) $x + y = x \implies y = 0$
3. (Multiplicative Cancellation) $xy = xz \implies (x \neq 0 \implies y = z)$
4. (Uniqueness of Multiplicative Identity) $xy = x \implies y = 1$
5. (No Zero Divisors) $xy = 0 \implies (x = 0 \vee y = 0)$

Theorem 1.1.4 (Properties of Fields)

Let F be a field. Then for all $x, y \in F$ we have

$$\begin{array}{llll} 0 \cdot x = 0 & -(-x) = x & -(x + y) = -x - y & (-1)x = x \\ (-x)y = -(xy) & (-x)(-y) = xy & (a^{-1})^{-1} = a & (ab)^{-1} = a^{-1}b^{-1} \\ & & (-a)^{-1} = -a^{-1} & \end{array}$$

Definition 1.1.7 (Order)

An order on a set X is a binary relation \leq on X such that

1. (Totality) $\forall x, y \in X (x \leq y \vee y \leq x)$
2. (Antisymmetry) $\forall x, y \in X (x \leq y \wedge y \leq x) \implies x = y$
3. (Transitivity) $\forall x, y, z \in X (x \leq y \wedge y \leq z) \implies x \leq z$

Remark (Order defined using the $<$ operator)

Note that we may also make a definition of the above using $<$ instead of \leq . Then the properties that will define an order will be:

1. (Trichotomy Property) $\forall x, y \in X (x < y \vee y < x \vee x = y)$
2. (Transitive Property) $\forall x, y, z \in X (x < y \wedge y < z) \implies x < z$
3. (Additive Property) $\forall x, y, z \in X x < y \implies x + z < y + z$

4. (Multiplicative Property) $\forall a, b, c \in X$ we have

$$(a) \ a < b \wedge c > 0 \implies ac < bc$$

$$(b) \ a < b \wedge c < 0 \implies bc < ac$$

Remark (Non-negative and Non-positive)

Let $a \in \mathbb{R}$. We say that a is non-negative when $0 \leq a$ and that a is non-positive $a \leq 0$.

Remark

Some ways of writing the order symbol. Let $a, b, c \in \mathbb{R}$

- $b \not\leq a$ is equivalent to $b \not\leq a$
- $b \leq a$ is equivalent to $b < a \vee b = a$
- If $a \leq b$ and $b \leq c$, we can write $a \leq b \leq c$.

Theorem 1.1.5 ($\mathbb{N}, \mathbb{Z}, \mathbb{Q}$ and \mathbb{R} are ordered)

Each of $\mathbb{N}, \mathbb{Z}, \mathbb{Q}$ and \mathbb{R} is an ordered set using the standard order \leq . Under the inclusions $\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R}$ the orders coincide (e.g. when $a, b \in \mathbb{N}$ we have $a \leq b$ in \mathbb{N} if and only if $a \leq b$ in \mathbb{R})

Definition 1.1.8 (Ordered Field)

An ordered field is a field F with an order \leq such that for all $x, y, z \in F$

1. $x \leq y \implies x + z \leq y + z$, and
2. $0 \leq x \wedge 0 \leq y \implies 0 \leq xy$.

Theorem 1.1.6 (\mathbb{Q} and \mathbb{R} as Ordered Fields)

\mathbb{Q} and \mathbb{R} are ordered fields.

Theorem 1.1.7 (Properties of Ordered Fields)

Let F be an ordered field. Then $\forall x, y, z \in F$ we have

1. $x > 0 \implies -x < 0$ and $x < 0 \implies -x > 0$
2. $x \neq 0 \implies x^2 > 0$ and in particular $1 \neq 0$
3. $0 < x < y \implies 0 < \frac{1}{y} < \frac{1}{x}$

Definition 1.1.9 (Absolute Value)

Let F be an ordered field. For $a \in F$ we define the absolute value of a to be

$$|a| = \begin{cases} a & \text{if } a \geq 0, \\ -a & \text{if } a \leq 0. \end{cases}$$

Theorem 1.1.8 (Properties of Absolute Values)

Let F be an ordered field. Then for all $x, y, z \in F$ we have

1. (Positive Definiteness) $|x| \geq 0$ and $|x| = 0 \iff x = 0$
2. (Symmetry) $|x - y| = |y - x|$
3. (Multiplicativeness) $|xy| = |x||y|$
4. (Triangle Inequality) $|x + y| \leq |x| + |y|$
5. (Approximation) $|x - y| \leq z \implies y - z \leq x \leq y + z$

Theorem 1.1.9 (Basic Order Properties in \mathbb{Z})

1. $\forall n \in \mathbb{Z} (n \in \mathbb{N} \iff n \geq 0)$
2. $\forall k, n \in \mathbb{Z} (k \leq n \iff k < n + 1)$

1.2 Completeness Axiom

Definition 1.2.1 (Upper and Lower Bounds)

Let X be an ordered set and let $A \subseteq X$

1. We say that A is bounded above (in X) when $\exists b \in X \forall a \in A \ a \leq b$, in which case we call b the upper bound of A .
2. We say that A is bounded below (in X) when $\exists c \in X \forall a \in A \ c \leq a$, in which case we call c the lower bound of A .

We say that A is bounded when it is bounded above and below.

Definition 1.2.2 (Supremum and Infimum)

Let X be an ordered set and let $A \subseteq X$.

1. We say that A has a supremum (or the least upper bound) when

$$\exists b \in X (\forall a \in A \ a \leq b) \quad \forall c \in X (\forall a \in A \ a \leq c) \quad b < c.$$

We write $b = \sup A$.

Now if $b = \sup A$ and $b \in A$, we call b the maximum of A , and denote it as $b = \max A$.

2. We say that A has an infimum (or the greatest lower bound) when

$$\exists d \in X (\forall a \in A \, d \leq a) \quad \forall c \in X (\forall a \in A \, c \leq a) \quad c < d.$$

We write $d = \inf A$.

Now if $d = \inf A$ and $d \in A$, we call d the minimum of A , and denote it as $d = \min A$.

Theorem 1.2.1 (Approximation Property of Supremum and Infimum)

Let $\emptyset \neq A \subseteq \mathbb{R}$.

1. $b = \sup A \implies \forall 0 < \epsilon \in \mathbb{R} \, \exists x \in A \, (b - \epsilon < x \leq b)$
2. $c = \inf A \implies \forall 0 < \epsilon \in \mathbb{R} \, \exists x \in A \, (c \leq x < c + \epsilon)$

Theorem 1.2.2 (Completeness Properties of \mathbb{R})

1. $\forall \emptyset \neq A \subseteq \mathbb{R}$, if A is bounded above, then A has a supremum in \mathbb{R}
2. $\forall \emptyset \neq A \subseteq \mathbb{R}$, if A is bounded below, then A has an infimum in \mathbb{R}

Theorem 1.2.3 (Well-Ordering Properties of \mathbb{Z} in \mathbb{R})

1. Every nonempty subset of \mathbb{Z} which is bounded above in \mathbb{R} has a maximum.
2. Every nonempty subset of \mathbb{Z} which is bounded below in \mathbb{R} has a minimum. In particular, every nonempty subset of \mathbb{N} has a minimum.

Theorem 1.2.4 (Floor and Ceiling Properties of \mathbb{Z} in \mathbb{R})

1. (Floor Properties) $\forall x \in \mathbb{R} \, \exists! n \in \mathbb{Z} \, (x - 1 < n \leq x)$
2. (Ceiling Properties) $\forall x \in \mathbb{R} \, \exists! n \in \mathbb{Z} \, (x \leq n < x + 1)$

Definition 1.2.3 (Floor and Ceiling Functions)

Given $x \in \mathbb{R}$ we define the floor of x to be the unique $n \in \mathbb{Z}$ with $x - 1 < n \leq x$ and denote the floor of x by $\lfloor x \rfloor$. The function $f : \mathbb{R} \rightarrow \mathbb{Z}$ given by $f(x) = \lfloor x \rfloor$ is called the floor function.

Similarly, we define the ceiling of x to be the unique $n \in \mathbb{Z}$ with $x \leq n < x + 1$ and denote the ceiling of x by $\lceil x \rceil$. The function $f : \mathbb{R} \rightarrow \mathbb{Z}$ given by $f(x) = \lceil x \rceil$ is called the ceiling function.

Theorem 1.2.5 (Archimedean Properties of \mathbb{Z} in \mathbb{R})

1. $\forall x \in \mathbb{R} \, \exists n \in \mathbb{Z} \, (n > x)$

$$2. \forall x \in \mathbb{R} \exists m \in \mathbb{Z} (m < x)$$

Theorem 1.2.6 (Density of \mathbb{Q})

$$\forall a, b \in \mathbb{R} (a < b) \quad \exists q \in \mathbb{Q} (a < q < b)$$

Chapter 2

Sequences

2.1 Limits of Sequences

Definition 2.1.1 (Sequence)

For $p \in \mathbb{Z}$, let $\mathbb{Z}_{\geq p} = \{k \in \mathbb{Z} | k \geq p\}$. A sequence in a set A is a function of the form $x : \mathbb{Z}_{\geq p} \rightarrow A$ for some $p \in \mathbb{Z}$. Given a sequence $x : \mathbb{Z}_{\geq p} \rightarrow A$, the k -th term of the sequence is the element $x_k = x(k) \in A$, and we denote the sequence x by

$$\langle x_k \rangle_{k \geq p} = \{x_k | k \geq p\} = \{x_p, x_{p+1}, x_{p+2}, \dots\}$$

Note that the range of the sequence $\langle x_k \rangle_{k \geq p}$ is the set $\{x_k\}_{k \geq p} = \{x_k | k \geq p\}$.

Remark

While the notation $\{x_k\}_{k \geq p}$ is more commonly used, since this set of notes works a lot between sequences and sets, we shall use the notation $\langle x_k \rangle_{k \geq p}$ to denote a sequence instead to make a clear distinction between the two.

Definition 2.1.2 (Subsequence)

Let $\langle x_k \rangle_{k \geq p}$ be a sequence. A subsequence of $\langle x_k \rangle_{k \geq p}$ is a sequence of the form $\langle x_{k_n} \rangle_{n \in \mathbb{N}}$ such that $k_1 < k_2 < k_3 < \dots$ and $x_{k_1} < x_{k_2} < x_{k_3} < \dots$, where $x_{k_l} = x_m$ for all $n \geq l \in \mathbb{N}$ and a unique $k \geq m \in \mathbb{Z}_{\geq p}$.

Remark

In other words, a subsequence $\langle x_{k_n} \rangle_{n \in \mathbb{N}}$ is constructed from $\langle x_k \rangle_{k \geq p}$ by "removing" from $x_p, x_{p+1}, x_{p+2}, \dots$ all the x_m 's except for those such that $m = k_l$ for some l .

Definition 2.1.3 (Extended Ordered Field)

Let F be an ordered field. We can define the extended ordered field \hat{F} to be the set $\hat{F} =$

$F \cup \{-\infty, \infty\}$, such that $\forall a \in F, -\infty < a < \infty$.

We also define, $\forall a \in F$:

- $a + \infty = \infty$,
- $a - \infty = -\infty$,
- if $a > 0$, then $a \cdot \infty = \infty$, and
- if $a < 0$, then $a \cdot \infty = -\infty$.

We define some indeterminant forms:

$$\infty - \infty, \infty \cdot 0, \frac{\infty}{\infty}, \frac{\infty}{0}, \frac{0}{\infty}$$

We extend the order relation j on F such that $-\infty < \infty$.

Definition 2.1.4 (Convergence, Divergence and Limits of a Sequence)

Let F be an extended ordered field. and $\langle x_k \rangle_{k \geq p}$ be a sequence in F . For $a \in F$, we say that the sequence $\langle x_k \rangle_{k \geq p}$ converges to a (or that the limit of $\langle x_k \rangle_{k \geq p}$ is equal to a), and we write $x_k \rightarrow a$ (as $k \rightarrow \infty$), or we write $\lim_{k \rightarrow \infty} x_k = a$, when

$$\forall 0 < \epsilon \in F \exists m \in \mathbb{Z} \forall k \in \mathbb{Z}_{\geq p} (k \geq m \implies |x_k - a| \leq \epsilon).$$

We say that the sequence $\langle x_k \rangle_{k \geq p}$ diverges (in F) when it does not converge (to any $a \in F$). We say that $\langle x_k \rangle_{k \geq p}$ diverges to infinity, or that the limit of $\langle x_k \rangle_{k \geq p}$ is equal to infinity, and we write $x_k \rightarrow \infty$ (as $k \rightarrow \infty$), or we write $\lim_{k \rightarrow \infty} x_k = \infty$, when

$$\forall r \in F \exists m \in \mathbb{Z} \forall k \in \mathbb{Z}_{\geq p} (k \geq m \implies x_k \geq r)$$

Similarly, we say that $\langle x_k \rangle_{k \geq p}$ diverges to $-\infty$, or that the limit of $\langle x_k \rangle_{k \geq p}$ is equal to negative infinity, and we write $x_k \rightarrow -\infty$ (as $k \rightarrow \infty$), or we write $\lim_{k \rightarrow \infty} x_k = -\infty$, when

$$\forall r \in F \exists m \in \mathbb{Z} \forall k \in \mathbb{Z}_{\geq p} (k \geq m \implies x_k \leq r)$$

Theorem 2.1.1 (Independence of Limit from Initial Terms)

Let $\langle x_k \rangle_{k \geq p}$ be a sequence in a subfield F of \mathbb{R} .

1. If $q \geq p$ and $y_k = x_k$ for all $k \geq q$, then $\langle x_k \rangle_{k \geq p}$ converges iff $\langle y_k \rangle_{k \geq q}$ converges, and in this case $\lim_{k \rightarrow \infty} x_k = \lim_{k \rightarrow \infty} y_k$.

(Note that in this statement, $\langle y_k \rangle_{k \geq q}$ is a subsequence of $\langle x_k \rangle_{k \geq p}$, such that it takes on all the elements of the sequence after some $q \geq p$.)

2. If $l \geq 0$ and $y_k = x_{k+l}$ for all $k \geq p$, then $\langle x_k \rangle_{k \geq p}$ converges iff $\langle y_k \rangle_{k \geq p}$ converges, and in this case $\lim_{k \rightarrow \infty} x_k = \lim_{k \rightarrow \infty} y_k$.

(Note that in this statement, $\langle x_k \rangle_{k \geq p}$ is a subsequence of $\langle y_k \rangle_{k \geq p}$ instead, such that $\langle x_k \rangle_{k \geq p}$ takes on all the values of $\langle y_k \rangle_{k \geq p}$ from $k + l$.)

Remark

Because of the above theorem, we often simply denote $\langle x_k \rangle_{k \geq p}$ as $\langle x_k \rangle$

Theorem 2.1.2 (Uniqueness of Limit)

Let $\langle x_k \rangle$ be a sequence in an ordered field F . If $\langle x_k \rangle$ has a limit (finite or infinite) then its limit is unique.

2.2 Limit Theorems

Theorem 2.2.1 (Basic Limits)

In any ordered field F , for $a \in F$ we have

$$\lim_{k \rightarrow \text{infy}} a = a, \quad \lim_{k \rightarrow \infty} k = \infty, \quad \lim_{k \rightarrow \infty} \frac{1}{k} = 0$$

Theorem 2.2.2 (Operations on Limits)

Let $\langle x_k \rangle$ and $\langle y_k \rangle$ be sequences in an ordered field F and let $c \in F$. SPS that $\langle x_k \rangle$ and $\langle y_k \rangle$ both converge with $x_k \rightarrow a$ and $y_k \rightarrow b$. Then

1. $cx_k \rightarrow ca$
2. $(x_k + y_k) \rightarrow a + b$
3. $(x_k - y_k) \rightarrow a - b$
4. $x_k y_k \rightarrow ab$
5. If $b \neq 0$, then $\frac{x_k}{y_k} \rightarrow \frac{a}{b}$

Theorem 2.2.3 (Extended Operations on Limits)

Let $\langle x_k \rangle$ and $\langle y_k \rangle$ be sequences in F . SPS that $\lim_{k \rightarrow \infty} x_k = u$ and $\lim_{k \rightarrow \infty} y_k = v$, where $u, v \in \hat{F}$.

1. If $u + v$ is defined in \hat{F} , then $\lim_{k \rightarrow \infty} (x_k + y_k) = u + v$.
2. If $u - v$ is defined in \hat{F} , then $\lim_{k \rightarrow \infty} (x_k - y_k) = u - v$.
3. If uv is defined in \hat{F} , then $\lim_{k \rightarrow \infty} x_k y_k = uv$.

4. If $\frac{u}{v}$ is defined in \hat{F} , then $\lim_{k \rightarrow \infty} \frac{x_k}{y_k} = \frac{u}{v}$

Theorem 2.2.4 (Monotonic Surjective Functions)

Let I and J be intervals in a subfield $F \subseteq \mathbb{R}$. SPS $f : I \rightarrow J$ is increasing and surjective. Let $\langle x_k \rangle$ be a sequence in I . Then

1. If $x_k \rightarrow a \in I$, then $f(x_k) \rightarrow f(a) \in J$.
2. If $x_k \rightarrow u \in F \cup \{\infty\}$ is the right endpoint of I , then $f(x_k) \rightarrow v \in F \cup \{\infty\}$ is the right endpoint in J .
3. If $x_k \rightarrow u \in F \cup \{-\infty\}$ is the left endpoint of I , then $f(x_k) \rightarrow v \in F \cup \{-\infty\}$ is the left endpoint in J .

Theorem 2.2.5 (Basic Elementary functions Acting on Limits)

Let $\langle x_k \rangle$ be a sequence in \mathbb{R} and let $b \in \mathbb{R}$. Then

1. $x_k \rightarrow a > 0 \implies x_k^b \rightarrow a^b$ and

$$x_k \rightarrow \infty \implies \lim_{k \rightarrow \infty} x_k^b = \begin{cases} \infty & b > 0 \\ 1 & b = 0 \\ 0 & b < 0 \end{cases}$$

2. $(x_k \rightarrow \infty \wedge b > 0) \implies b^{x_k} \rightarrow b^a$ and

$$(x_k \rightarrow \infty \wedge b > 0) \implies \lim_{k \rightarrow \infty} b^{x_k} = \begin{cases} \infty & b > 1 \\ 1 & b = 1 \\ 0 & 0 < b < 1 \end{cases}$$

3. $(x_k \rightarrow a > 0 \wedge b > 0) \implies \log_b x_k \rightarrow \log_b a$ and

$$(x_k \rightarrow \infty \wedge b > 0) \implies \lim_{k \rightarrow \infty} \log_b x_k = \begin{cases} \infty & b > 1 \\ 0 & b = 1 \\ -\infty & 0 < b < 1 \end{cases}$$

4. $x_k \rightarrow a \implies (\sin x_k \rightarrow \sin a \wedge \cos x_k \rightarrow \cos a)$ and
 $(x_k \rightarrow a (\forall t \in \mathbb{Z} a \neq \frac{\pi}{2} + 2\pi t)) \implies \tan x_k \rightarrow \tan a$

5. $x_k \rightarrow a \in [-1, 1] \implies (\arcsin x_k \rightarrow \arcsin a \wedge \arccos x_k \rightarrow \arccos a)$,
 $x_k \rightarrow a \implies \arctan x_k \rightarrow \arctan a$,
 $x_k \rightarrow \infty \implies \arctan x_k \rightarrow \frac{\pi}{2}$, and
 $x_k \rightarrow -\infty \implies \arctan x_k \rightarrow -\frac{\pi}{2}$

Theorem 2.2.6 (Comparison Theorem for Sequences)

Let $\langle x_k \rangle$ and $\langle y_k \rangle$ be sequences in a subfield $F \subseteq \mathbb{R}$. SPS that $x_k \leq y_k$ for all k . Then

1. $(x_k \rightarrow a \wedge y_k \rightarrow b) \implies a \leq b$
2. $x_k \rightarrow \infty \implies y_k \rightarrow \infty$
3. $y_k \rightarrow -\infty \implies x_k \rightarrow -\infty$

Theorem 2.2.7 (Squeeze Theorem for Sequences)

Let $\langle x_k \rangle$, $\langle y_k \rangle$ and $\langle z_k \rangle$ be sequences in a subfield $F \subseteq \mathbb{R}$.

1. $(\forall k \in \mathbb{N} \ x_k \leq y_k \leq z_k \wedge x_k \rightarrow a \wedge z_k \rightarrow a) \implies y_k \rightarrow a$
2. $(\forall k \in \mathbb{N} \ |x_k| \leq y_k \wedge y_k \rightarrow 0) \implies x_k \rightarrow 0$

Definition 2.2.1 (Bounds)

Let $\langle x_k \rangle$ be a sequence in an ordered set X . We say that

1. $\langle x_k \rangle$ is bounded above iff the set $\{x_k | n \in \mathbb{N}\}$ is bounded above;
2. $\langle x_k \rangle$ is bounded below iff the set $\{x_k | n \in \mathbb{N}\}$ is bounded below.

2.3 Bolzano-Weierstrass Theorem**Definition 2.3.1 (Increasing, Decreasing, and Monotonic Sequences)**

Let $\langle x_k \rangle$ be a sequence in a subfield $F \subseteq \mathbb{R}$. We say that

1. $\langle x_k \rangle$ is increasing iff $\forall k, l \in \mathbb{Z}_{\geq p} (k \leq l \implies x_k \leq x_l)$
2. $\langle x_k \rangle$ is strictly increasing iff $\forall k, l \in \mathbb{Z}_{\geq p} (k < l \implies x_k < x_l)$
3. $\langle x_k \rangle$ is decreasing iff $\forall k, l \in \mathbb{Z}_{\geq p} (k \leq l \implies x_k \geq x_l)$
4. $\langle x_k \rangle$ is strictly decreasing iff $\forall k, l \in \mathbb{Z}_{\geq p} (k < l \implies x_k > x_l)$

We say that $\langle x_k \rangle$ is monotonic when it is either increasing or decreasing only.

Theorem 2.3.1 (Monotone Convergence Theorem)

Let $\langle x_k \rangle$ be a sequence in \mathbb{R} .

1. SPS $\langle x_k \rangle$ is increasing. If $\langle x_k \rangle$ is bounded above, then $x_k \rightarrow \sup\{x_k\}$. If $\langle x_k \rangle$ is not bounded above, then $x_k \rightarrow \infty$.
2. SPS $\langle x_k \rangle$ is decreasing. If $\langle x_k \rangle$ is bounded below, then $x_k \rightarrow \inf\{x_k\}$. If $\langle x_k \rangle$ is not bounded below, then $x_k \rightarrow -\infty$.

Theorem 2.3.2 (Nested Interval Theorem)

Let I_0, I_1, I_2, \dots be nonempty, closed, and bounded intervals in \mathbb{R} . SPS $I_0 \supseteq I_1 \supseteq I_2 \supseteq \dots$, then $\bigcap_{k=0}^{\infty} I_k \neq \emptyset$. Moreover, if the lengths of these intervals satisfy $|I_k| \rightarrow 0$ as $k \rightarrow \infty$, then $\bigcap_{k=0}^{\infty} I_k$ is a single point.

Definition 2.3.2 (Rearrangement of a Sequence)

Let $\langle x_k \rangle_{k \geq p}$ be a sequence in a set X . Given a bijective function $F : \mathbb{Z}_{\geq q} \rightarrow \mathbb{Z}_{\geq p}$ such that $f(l) = k_l$ and let $y_l = x_{k_l}$ for $l \geq q$. Then the sequence $\langle y_l \rangle_{l \geq q}$ is called a rearrangement of the sequence $\langle x_k \rangle_{k \geq p}$.

Theorem 2.3.3 (Convergence of Subsequences and Rearrangements)

Let $\langle x_k \rangle$ be a sequence in a subfield $F \subseteq \mathbb{R}$. SPS that $x_k \rightarrow a$. Then

1. every subsequence of $\langle x_k \rangle$ converges to a ; and
2. every rearrangement of $\langle x_k \rangle$ converges to a .

Theorem 2.3.4 (Bolzano-Weierstrass Theorem)

Every bounded sequence of \mathbb{R} has a convergent subsequence.

2.4 Cauchy Sequences

Definition 2.4.1 (Cauchy)

Let $\langle x_k \rangle_{k \geq p}$ be a sequence in a subfield $F \subseteq \mathbb{R}$. We say that $\langle x_k \rangle$ is Cauchy when

$$\forall \epsilon > 0 \exists m \in \mathbb{Z} \forall k, l \in \mathbb{Z}_{k \geq p} (k, l \geq m \implies |x_k - x_l| \leq \epsilon)$$

Theorem 2.4.1 (Cauchy Criterion for Convergence)

Let $\langle x_k \rangle$ be a sequence of \mathbb{R} . Then $\langle x_k \rangle$ is Cauchy iff $\langle x_k \rangle$ converges (to some point $a \in \mathbb{R}$).

Chapter 3

Functions on \mathbb{R}

3.1 Two-Sided Limits

Definition 3.1.1 (Limit Point)

Let $U \subseteq F$ where F is an ordered field. Let $f : U \rightarrow F$. For $a \in F$ we say that a is a limit point of U when

$$\forall \epsilon > 0 \exists x \in U \ 0 < |x - a| < \epsilon$$

When a is a limit point of A , we make the following definitions.

1. For $b \in F$ we say that the limit of $f(x)$ as x tends to a is equal to b , and write $\lim_{x \rightarrow a} f(x) = b$ when

$$\forall \epsilon > 0 \exists \delta > 0 \forall x \in U \ (0 < |x - a| \leq \delta \implies |f(x) - b| \leq \epsilon).$$

2. We say that the limit of $f(x)$ as x tends to a is equal to infinity, and write $\lim_{x \rightarrow a} f(x) = \infty$ when

$$\forall r \in F \exists \delta > 0 \forall x \in U \ (0 < |x - a| \leq \delta \implies f(x) \geq r).$$

3. We say that the limit of $f(x)$ as x tends to a is equal to negative infinity, and write $\lim_{x \rightarrow a} f(x) = -\infty$ when

$$\forall r \in F \exists \delta > 0 \forall x \in U \ (0 < |x - a| \leq \delta \implies f(x) \leq r).$$

Definition 3.1.2 (Limit Point from Above and Below)

Let $U \subseteq F$ where F is an ordered field. Let $f : U \rightarrow F$.

For $a \in F$, we say that a is a **limit point from below** when

$$\forall \delta > 0 \exists x \in U \ a - \delta < x < a$$

When a is a limit point of U from below and $b \in F$, we define:

1. $\lim_{x \rightarrow a^-} f(x) = b \iff \forall \epsilon > 0 \exists \delta > 0 \forall x \in U \ (a - \delta \leq x < a \implies |f(x) - b| \leq \epsilon).$
2. $\lim_{x \rightarrow a^-} f(x) = \infty \iff \forall r \in F \exists \delta > 0 \forall x \in U \ (a - \delta \leq x < a \implies f(x) \geq r).$
3. $\lim_{x \rightarrow a^-} f(x) = -\infty \iff \forall r \in F \exists \delta > 0 \forall x \in U \ (a - \delta \leq x < a \implies f(x) \leq r).$

For $a \in F$, we say that a is a **limit point from above** when

$$\forall \delta > 0 \exists x \in U \ a < x \leq a + \delta$$

When a is a limit point of U from above and $b \in F$, we define:

1. $\lim_{x \rightarrow a^+} f(x) = b \iff \forall \epsilon > 0 \exists \delta > 0 \forall x \in U \ (a < x \leq a + \delta \implies |f(x) - b| \leq \epsilon).$
2. $\lim_{x \rightarrow a^+} f(x) = \infty \iff \forall r \in F \exists \delta > 0 \forall x \in U \ (a < x \leq a + \delta \implies f(x) \geq r).$
3. $\lim_{x \rightarrow a^+} f(x) = -\infty \iff \forall r \in F \exists \delta > 0 \forall x \in U \ (a < x \leq a + \delta \implies f(x) \leq r).$

Definition 3.1.3 (Infinity As A Limit Point)

Let $U \subseteq F$ where F is an ordered field. Let $f : U \rightarrow F$.

We say that infinity is a limit point (from below) when U is not bounded above, i.e. $\forall m \in F \exists x \in U \ x \geq m$. When U is not bounded above and $b \in F$, we make the following definitions:

1. $\lim_{x \rightarrow \infty} f(x) = b \iff \forall \epsilon > 0 \exists m \in F \forall x \in U \ (x \geq m \implies |f(x) - b| \leq \epsilon).$
2. $\lim_{x \rightarrow \infty} f(x) = \infty \iff \forall r \in F \exists m \in F \forall x \in U \ (x \geq m \implies f(x) \geq r).$
3. $\lim_{x \rightarrow \infty} f(x) = -\infty \iff \forall r \in F \exists m \in F \forall x \in U \ (x \geq m \implies f(x) \leq r).$

We say that negative infinity is a limit point (from above) when U is not bounded below, i.e. $\forall m \in F \exists x \in U \ x \leq m$. When U is not bounded below and $b \in F$, we make the following definitions:

1. $\lim_{x \rightarrow -\infty} f(x) = b \iff \forall \epsilon > 0 \exists m \in F \forall x \in U \ (x \leq m \implies |f(x) - b| \leq \epsilon).$
2. $\lim_{x \rightarrow -\infty} f(x) = \infty \iff \forall r \in F \exists m \in F \forall x \in U \ (x \leq m \implies f(x) \geq r).$
3. $\lim_{x \rightarrow -\infty} f(x) = -\infty \iff \forall r \in F \exists m \in F \forall x \in U \ (x \leq m \implies f(x) \leq r).$

Theorem 3.1.1 (Two-sided Limits)

Let F be a subfield of \mathbb{R} . Let $A \subseteq F$. Let $f : A \rightarrow F$. Let $a \in F$. SPS that a is a limit point of A both from above and below. Then $\forall u \in F$, we have $\lim_{x \rightarrow a} f(x) = u \iff \lim_{x \rightarrow a^-} f(x) = u = \lim_{x \rightarrow a^+} f(x)$.

Theorem 3.1.2 (Sequential Characterization of Limits of Functions)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, let $f : A \rightarrow F$ and $u \in F$.

1. When $a \in F$ is a limit point of A , $\lim_{x \rightarrow a} f(x) = u$ iff for every sequence $\langle x_k \rangle$ in $A \setminus \{a\}$ with $x_k \rightarrow a$ we have $f(x_k) \rightarrow u$.
2. When a is a limit point of A from below, $\lim_{x \rightarrow a^-} f(x) = u$ iff for every sequence $\langle x_k \rangle$ in $\{x \in A \mid x < a\}$ with $x_k \rightarrow a$ we have $f(x_k) \rightarrow u$.
3. When a is a limit point of A from above, $\lim_{x \rightarrow a^+} f(x) = u$ iff for every sequence $\langle x_k \rangle$ in $\{x \in A \mid x > a\}$ with $x_k \rightarrow a$ we have $f(x_k) \rightarrow u$.
4. When A is not bounded above, $\lim_{x \rightarrow \infty} f(x) = u$ iff for every sequence $\langle x_k \rangle$ in A with $x_k \rightarrow \infty$ we have $f(x_k) \rightarrow u$.
5. When A is not bounded below, $\lim_{x \rightarrow -\infty} f(x) = u$ iff for every sequence $\langle x_k \rangle$ in A with $x_k \rightarrow -\infty$ we have $f(x_k) \rightarrow u$.

Remark

It follows from the Sequential Characterization of Limits of Functions that all the theorems about limits of sequences (see [Chapter 2](#)) imply analogous theorems in the more general setting of limits of functions. We will state those theorems for easier reference.

Theorem 3.1.3 (Local Determination of Limits)

Let F be a subfield of \mathbb{R} , let $A, B \subseteq F$, let $f : A \rightarrow F$ and $g : B \rightarrow F$. SPS that $a \in F$ is a limit point of both sets A and B , and that for some $\delta > 0$ we have $C = \{x \in A \mid 0 < |x - a| \leq \delta\} \subseteq \{x \in B \mid 0 < |x - a| \leq \delta\}$ and that $f(x) = g(x)$ for all $x \in C$. Then for $u \in \hat{F}$

$$\lim_{x \rightarrow a} g(x) = u \iff \lim_{x \rightarrow a} f(x) = u$$

Analogous results hold for limits $x \rightarrow a^\pm$ and $x \rightarrow \pm\infty$.

Theorem 3.1.4 (Uniqueness of Limits)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, let $f : A \rightarrow F$ and let $a \in F$ be a limit point of A . For $u, v \in \hat{F}$,

$$\lim_{x \rightarrow a} f(x) = u \wedge \lim_{x \rightarrow a} f(x) = v \implies u = v$$

Analogous result holds for limits $x \in a^\pm$ and $x \rightarrow \pm\infty$.

Theorem 3.1.5 (Extended Operations on Limits)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, let $f, g : A \rightarrow F$, and let $a \in F$ be a limit point of A . Let $u, v \in \cap F$, and SPS that $\lim_{x \rightarrow a} f(x) = u$ and $\lim_{x \rightarrow a} g(x) = v$. Then

1. $u \pm v \in \hat{F} \implies \lim_{x \rightarrow a} (f \pm g)(x) = u \pm v$
2. $uv \in \hat{F} \implies \lim_{x \rightarrow a} (f \cdot g)(x) = uv$
3. $\frac{u}{v} \in \hat{F} \implies \lim_{x \rightarrow a} (\frac{f}{g})(x) = \frac{u}{v}$

Analogous results hold for limits $x \rightarrow a^\pm$ and $x \rightarrow \pm\infty$.

Theorem 3.1.6 (Basic Elementary Functions Acting on Limits)

For $A \subseteq \mathbb{R}$, let $f : A \rightarrow \mathbb{R}$ and $a, b \in \mathbb{R}$ with a as a limit point of A . We have

1. $\lim_{x \rightarrow a} f(x) = b > 0 \implies \lim_{x \rightarrow a} f(x)^c = b^c$

$$\lim_{x \rightarrow a} f(x) = \infty \implies \lim_{x \rightarrow a} f(x)^c = \begin{cases} \infty & c > 0 \\ 1 & c = 0 \\ 0 & c < 0 \end{cases}$$

$$\forall x \in A (f(x) > 0 \wedge \lim_{x \rightarrow a} f(x) = 0) \implies \lim_{x \rightarrow a} f(x)^c = \begin{cases} 0 & c > 0 \\ 1 & c = 0 \\ \infty & c < 0 \end{cases}$$
2. $\lim_{x \rightarrow a} f(x) = \infty \wedge c > 0 \implies \lim_{x \rightarrow a} c^{f(x)} = \begin{cases} \infty & c > 1 \\ 1 & c = 1 \\ 0 & c < 1 \end{cases}$

$$\lim_{x \rightarrow a} f(x) = b \wedge c > 0 \implies \lim_{x \rightarrow a} c^{f(x)} = c^b$$

$$\lim_{x \rightarrow a} f(x) = -\infty \wedge c > 0 \implies \lim_{x \rightarrow a} c^{f(x)} = \begin{cases} \infty & c < 1 \\ 1 & c = 1 \\ 0 & c > 1 \end{cases}$$
3. $\lim_{x \rightarrow a} f(x) = b > 0 \wedge c > 0 \implies \lim_{x \rightarrow a} \log_c f(x) = \log_c b$

$$\lim_{x \rightarrow a} f(x) = \infty \wedge c > 0 \implies \lim_{x \rightarrow a} \log_c f(x) = \begin{cases} \infty & c > 1 \\ 0 & c = 1 \\ -\infty & c < 1 \end{cases}$$

$$\forall x \in A (f(x) > 0 \wedge \lim_{x \rightarrow a} f(x) = 0 \wedge 1 \neq c > 0) \implies \lim_{x \rightarrow a} \log_c f(x) = \begin{cases} -\infty & c > 1 \\ \infty & c < 1 \end{cases}$$
4. $\lim_{x \rightarrow a} f(x) = b \implies (\lim_{x \rightarrow a} \sin f(x) = \sin b \wedge \lim_{x \rightarrow a} \cos f(x) = \cos b)$

The limits $\lim_{x \rightarrow \pm\infty} \sin x$, $\lim_{x \rightarrow \pm\infty} \cos x$, and $\lim_{x \rightarrow \pm\infty} \tan x$ do not exist.

5. $\forall x \in A (f(x) \in [-1, 1] \wedge \lim_{x \rightarrow a} f(x) = b) \implies \lim_{x \rightarrow a} \arcsin f(x) = \arcsin b$.
 $\lim_{x \rightarrow a} f(x) = b \in \mathbb{R} \implies \lim_{x \rightarrow a} \arctan f(x) = \arctan b$
 $\lim_{x \rightarrow a} f(x) = \infty \implies \lim_{x \rightarrow a} \arctan f(x) = \frac{\pi}{2}$ and
 $\lim_{x \rightarrow a} f(x) = -\infty \implies \lim_{x \rightarrow a} \arctan f(x) = -\frac{\pi}{2}$.

Similar results hold for limits $x \rightarrow a^\pm$ and $x \rightarrow \pm\infty$ (unless stated otherwise in the above statements).

Theorem 3.1.7 (Comparison Theorem for Functions)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, let $f, g : A \rightarrow F$ and let $a \in F$ be a limit point of A . SPS that $\forall x \in A f(x) \leq g(x)$. Then

1. $\exists u, v \in \hat{F} (\lim_{x \rightarrow a} f(x) = u \wedge \lim_{x \rightarrow a} g(x) = v) \implies u \leq v$.
2. $\lim_{x \rightarrow a} f(x) = \infty \implies \lim_{x \rightarrow a} g(x) = \infty$, and
3. $\lim_{x \rightarrow a} g(x) = -\infty \implies \lim_{x \rightarrow a} f(x) = -\infty$.

Similar results hold for when $x \rightarrow a^\pm$ and $x \rightarrow \pm\infty$.

Theorem 3.1.8 (Squeeze Theorem for Functions)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, let $f, g, h : A \rightarrow F$, let $b \in F$ and let a be a limit point of A . We have that

1. $\forall x \in A (f(x) \leq g(x) \leq h(x) \wedge \lim_{x \rightarrow a} f(x) = b = \lim_{x \rightarrow a} h(x)) \implies \lim_{x \rightarrow a} g(x) = b$.
2. $\forall x \in A (|f(x)| \leq g(x) \wedge \lim_{x \rightarrow a} g(x) = 0) \implies \lim_{x \rightarrow a} f(x) = 0$

3.2 Continuity

Definition 3.2.1 (Continuity)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, and let $f : A \rightarrow F$. For all $a \in A$, we say that f is continuous at a iff

$$\forall a \in A \forall \epsilon > 0 \exists \delta > 0 \forall x \in A (|x - a| \leq \delta \implies |f(x) - f(a)| \leq \epsilon)$$

(Note that δ depends on a, f , and, especially, ϵ in general).

f is said to be continuous (in A) when f is continuous at every point $a \in A$.

Definition 3.2.2 (Limit Point and Continuity)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, let $f : A \rightarrow F$ and let $a \in A$. Then

1. if a is not a limit point of A then f is continuous on a ; and
2. if a is a limit point of A , then f is continuous on a iff $\lim_{x \rightarrow a} f(x) = f(a)$.

Theorem 3.2.1 (Sequential Characterization of Continuity)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, let $f : A \rightarrow F$ and let $a \in A$. Then f is continuous at a iff for every sequence $\langle x_k \rangle$ in A with $x_k \rightarrow a$ we have $f(x_k) \rightarrow f(a)$.

Theorem 3.2.2 (Operations on Continuous Functions)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, let $f, g : A \rightarrow F$, let $a \in A$ and let $c \in F$. SPS that f and g are continuous at a . Then the functions cf , $f \pm g$ and fg are call continuous at a , and if $g(a) \neq 0$, then the function $\frac{f}{g}$ is continuous at a .

Theorem 3.2.3 (Composition of Continuous Functions)

Let F be a subfield of \mathbb{R} , let $A, B \subseteq \mathbb{R}$, and let $f : A \rightarrow \mathbb{R}$ and $g : B \rightarrow \mathbb{R}$. Let $h = g \circ f : C \rightarrow \mathbb{R}$ where $C = A \cap f^{-1}(B)$

1. If f is continuous at $a \in C$ and g is continuous at $f(a)$, then h is continuous at a .
2. If f is continuous (in A) and g is continuous (in B), then h is continuous (in C).

Corollary 3.2.3.1 (Continuity of Elementary Functions)

Every elementary function is continuous (in their respective domain).

Theorem 3.2.4 (Functions Acting on Limits)

Let F be a subfield of \mathbb{R} , let $A, B \subseteq F$, let $f : A \rightarrow F$ and $g : B \rightarrow F$, and let $h = g \circ f : C \rightarrow F$ where $C = A \cap f^{-1}(B)$. Let s be a limit point of C (hence also of A) and let b be a limit point of B . Let $c \in F$. SPS that $\lim_{x \rightarrow a} f(x) = a \wedge \lim_{y \rightarrow b} g(y) = c$. SPS either that $\forall x \in C \setminus \{a\} f(x) \neq b$ or g is continuous at b . Then $\lim_{x \rightarrow a} h(x) = c$.

Theorem 3.2.5 (Intermediate Value Theorem)

Let I be an interval in \mathbb{R} and let $f : I \rightarrow \mathbb{R}$ be continuous (in I). Let $a, b \in I$ with $a \leq b$ and let $y \in \mathbb{R}$.

$$\min\{f(a), f(b)\} \leq y \leq \max\{f(a), f(b)\} \implies \exists x \in [a, b] (f(x) = y)$$

Definition 3.2.3 (Maximum, Minimum and Extreme Values)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, and let $f : A \rightarrow F$. For $a \in A$, if $f(x) \leq f(a)$ for every $x \in A$, then we say that $f(a)$ is the maximum value of f or that f attains its maximum value at a . Similarly for $b \in A$, if $f(b) \leq f(x)$ for every $x \in A$, we say that $f(b)$ is the minimum value of f , or that f attains its minimum at b .

We say that f attains its extreme values in A when f attains its maximum value at some

point $a \in A$ and f attains its minimum value at some point $b \in A$.

Theorem 3.2.6 (Extreme Value Theorem)

Let $a, b \in \mathbb{R}$ with $a < b$, and let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then f attains its extreme values in $[a, b]$.

Definition 3.2.4 (Uniform Continuity)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, and let $f : A \rightarrow F$. We say that f is uniformly continuous in A when

$$\forall \epsilon > 0 \exists \delta > 0 \forall a \in A \forall x \in A (|x - a| \leq \delta \implies |f(x) - f(a)| \leq \epsilon)$$

Theorem 3.2.7 (Closed Bounded Intervals and Uniform Continuity)

Let $a, b \in \mathbb{R}$ with $a < b$ and let $f : [a, b] \rightarrow \mathbb{R}$. If f is continuous (on $[a, b]$) then f is uniformly continuous (on $[a, b]$).

Chapter 4

Differentiability on \mathbb{R}

4.1 The Derivative

Definition 4.1.1 (Differentiable on a Point)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, and let $a \in A$ be a limit point of A . We say that f is differentiable at a when the limit

$$\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

exists in F . In this case we call the limit the derivative of f at a , and we denote that by $f'(a)$, so we have

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}.$$

When $a \in A$ is a limit point of A from the right, we say that f is differentiable from the right at a and that $f'_+(a)$ is the derivative from the right of f at a , when

$$f'_+(a) = \lim_{x \rightarrow a^+} \frac{f(x) - f(a)}{x - a}.$$

Similarly, when $a \in A$ is a limit point of A from the left, we say that f is differentiable from the left at a and that $f'_-(a)$ is the derivative from the left of f at a when

$$f'_-(a) = \lim_{x \rightarrow a^-} \frac{f(x) - f(a)}{x - a}.$$

Definition 4.1.2 (Differentiable in a Domain)

We say that f is differentiable (in A) when f is differentiable at every point $a \in A$, or that f is in C^1 . In this case, the derivative of f is the function $f' : A \rightarrow F$ defined by

$$f'(x) = \lim_{u \rightarrow x} \frac{f(u) - f(x)}{u - x}.$$

Definition 4.1.3 (Differentiable n-times & n-th Derivative)

When f' is differentiable at a , we denote the derivative of f' at a by $f''(a)$, and we call $f''(a)$ the second derivative of f at a . When $f''(a)$ exists for every a in A , we say that f is twice differentiable (in A), or that it is in C^2 , and we call the function $f'' : A \rightarrow F$ is called the second derivative of f . Similarly, $f'''(a)$ is the derivative of f'' at a , and so on.

More generally, for any function $f : A \rightarrow F$, we define its derivative to be the function $f' : B \rightarrow F$ where $B = \{a \in A \mid f \text{ is differentiable at } a\}$, and we define its second derivative to be the function $f'' : C \rightarrow F$ where $C = \{a \in B \mid f' \text{ is differentiable at } a\}$, and so on.

Remark

Note that

$$\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}.$$

To be precise, the limit on the left exists in F iff the limit on the right exists in F , and in this case the two limits are equal.

Theorem 4.1.1 (Definition of Differentiability in ϵ - δ)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, let $f : A \rightarrow F$, and let $a \in A$ be a limit point of A . Then f is differentiable at a with derivative $f'(a) \iff$

$$\forall \epsilon > 0 \exists \delta > 0 \forall x \in A (|x - a| \leq \delta \implies |f(x) - f(a) - f'(a)(x - a)| \leq \epsilon)$$

Definition 4.1.4 (Linearization & Tangent Line)

When $f : A \rightarrow F$ is differentiable at a with derivative $f'(a)$, the function

$$l(x) = f(a) + f'(a)(x - a)$$

is called the linearization of f at a .

Note that the graph $y = l(x)$ of the linearization is the line through the point $(a, f(a))$ with slope $f'(a)$. This line is called the tangent line to the graph $y = f(x)$ at the point $(a, f(a))$.

Theorem 4.1.2 (Differentiability \implies Continuity)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, let $f : A \rightarrow F$ and let $a \in A$ be a limit point of A .

$$f \text{ is differentiable at } a \implies f \text{ is continuous at } a.$$

4.2 Differentiability Theorems

Theorem 4.2.1 (Local Determination of the Derivative)

Let F be a subfield of \mathbb{R} , let $A, B \subseteq F$, let $f : A \rightarrow F$ and $g : B \rightarrow F$, and let $a \in A \cap B$ be a limit point of both A and B . SPS that for some $\delta > 0$ we have $\{x \in A \mid |x - a| \leq \delta\} \subset \{x \in B \mid |x - a| \leq \delta\}$. If g is differentiable at a then so is f , and we have $f'(a) = g'(a)$.

Theorem 4.2.2 (Operations on Derivatives)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, let $f, g : A \rightarrow F$, let $a \in A$ be a limit point of A , and let $c \in F$. SPS that f and g are differentiable at a . Then

1. (Linearity) the functions cf and $f \pm g$ are differentiable at a with

$$(cf)'(a) = cf'(a), \quad (f \pm g)'(a) = f'(a) \pm g'(a)$$

2. (Product Rule) the function fg is differentiable at a with

$$(fg)'(a) = f'(a)g(a) + f(a)g'(a)$$

3. (Reciprocal Rule) if $g(a) \neq 0$ then the function $1/g$ is differentiable at a with

$$\left(\frac{1}{g}\right)'(a) = -\frac{g'(a)}{g(a)^2}$$

4. (Quotient Rule) if $g(a) \neq 0$ then the function f/g is differentiable at a with

$$\left(\frac{f}{g}\right)'(a) = \frac{f'(a)g(a) - f(a)g'(a)}{g(a)^2}$$

Theorem 4.2.3 (Chain Rule)

Let F be a subfield of \mathbb{R} , let $A, B \subseteq F$, let $f : A \rightarrow F$, let $g : B \rightarrow F$ and let $h = g \circ f : C \rightarrow F$ where $C = A \cap f^{-1}(B)$. Let $a \in C$ be a limit point of C (hence also for A) and let $b = f(a) \in B$. SPS that f is differentiable at a and g is differentiable at b . Then h is differentiable at a with

$$h'(a) = g'(f(a))f'(a)$$

4.3 Inverse Function Theorems**Theorem 4.3.1 (Monotonic Functions)**

Let F be a subfield of \mathbb{R} , let $A \subseteq F$ and let $f : A \rightarrow F$. Then f is monotonic iff f has the property that for all $a, b, c \in A$, if b lies between a and c , then $f(b)$ lies between $f(a)$ and $f(c)$.

Theorem 4.3.2 (Continuity and Strictly Monotonous Functions)

Let I be an interval in \mathbb{R} , let $f : I \rightarrow \mathbb{R}$ and let $J = f(I)$.

1. If f is continuous then its range $J = f(I)$ is an interval in \mathbb{R} .
2. If f is injective and continuous, then f is strictly monotonic.

3. If $f : I \rightarrow J$ is strictly monotonic, then so is its inverse $g : J \rightarrow I$.

Theorem 4.3.3 (Inverse Function Theorem)

Let I be an interval in \mathbb{R} , let $f : I \rightarrow \mathbb{R}$ and let $J = f(I)$.

1. If f is bijective and continuous, then its inverse g is continuous.
2. If f is bijective and continuous, and f is differentiable at a with $f'(a) \neq 0$, then its inverse g is differentiable at $b = f(a)$ with $g'(b) = \frac{1}{f'(a)}$.

Theorem 4.3.4 (Derivatives of the Basic Elementary Functions)

The basic elementary functions have the following derivatives.

1. $(x^a)' = ax^{a-1}$ where $a, x \in \mathbb{R}$ and x^{a-1} is defined.
2. $(a^x)' = (\ln a) \cdot a^x$ where $a > 0$ and $x \in \mathbb{R}$ and
 $(\log_a x)' = \frac{1}{\ln a} \cdot \frac{1}{x}$ where $0 < a \neq 1$ and $x > 0$, and in particular
 $(e^x)' = e^x$ for all $x \in \mathbb{R}$ and $(\ln x)' = \frac{1}{x}$ for all $x > 0$.
3. $(\sin x)' = \cos x$ and $(\cos x)' = -\sin x$ for all $x \in \mathbb{R}$, and
 $(\tan x)' = \sec^2 x$ and $(\sec x)' = \sec x \tan x$ for all $x \in \mathbb{R}$ with $x \neq \frac{\pi}{2} + k\pi, k \in \mathbb{Z}$,
 $(\cot x)' = -\csc^2 x$ and $(\csc x)' = -\cot x \csc x$ for all $x \in \mathbb{R}$ with $x \neq \pi + k\pi, k \in \mathbb{Z}$.
4. $(\arcsin x)' = \frac{1}{\sqrt{1-x^2}}$ and $(\arccos x)' = \frac{-1}{\sqrt{1-x^2}}$ for $|x| < 1$,
 $(\sec^{-1} x)'' = \frac{1}{x\sqrt{x^2-1}}$ and $(\csc^{-1} x)' = \frac{-1}{x\sqrt{x^2-1}}$ for $|x| > 1$, and
 $(\arctan x)' = \frac{1}{1+x^2}$ and $(\cot^{-1} x)' = -\frac{1}{1+x^2}$ for all $x \in \mathbb{R}$.

4.4 Mean Value Theorem

Definition 4.4.1 (Local Maximum and Minimum)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, let $f : A \rightarrow F$ and let $a \in A$. We say that f has a local maximum value at a when

$$\exists \delta > 0 \forall x \in A (|x - a| \leq \delta \implies f(x) \leq f(a))$$

Similarly, we say that f has a local minimum value at a when

$$\exists \delta > 0 \forall x \in A (|x - a| \leq \delta \implies f(x) \geq f(a))$$

Theorem 4.4.1 (Fermat's Theorem)

Let F be a subfield of \mathbb{R} , let $A \subseteq F$, let $f : A \rightarrow F$. SPS that $a \in A$ is a limit point of A , both from above and from below. SPS that f is differentiable at a and that f has a local maximum or minimum value at a . Then $f'(a) = 0$.

Theorem 4.4.2 (Rolle's Theorem)

SPS that $a, b \in \mathbb{R}$ with $a \neq b$. If $f : [a, b] \rightarrow \mathbb{R}$ is continuous on $[a, b]$ and differentiable on (a, b) , with $f(a) = 0 = f(b)$, then there exists a point $c \in (a, b)$ such that $f'(c) = 0$.

Remark

The continuity hypothesis and differentiability hypothesis in Rolle's Theorem cannot be relaxed at even one point in $[a, b]$.

Theorem 4.4.3 (Mean Value Theorem)

Let $a, b \in \mathbb{R}$ with $a \neq b$. If $f : [a, b] \rightarrow \mathbb{R}$ is differentiable on (a, b) and continuous on $[a, b]$, then there exists a point $c \in (a, b)$ with

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

Theorem 4.4.4 (Cauchy/Generalized Mean Value Theorem)

Let $a, b \in \mathbb{R}$ with $a \neq b$. If $f, g : [a, b] \rightarrow \mathbb{R}$ are differentiable on (a, b) and continuous on $[a, b]$, then there exists a point $c \in (a, b)$ such that

$$g'(c)(f(b) - f(a)) = f'(c)(g(b) - g(a))$$

Corollary 4.4.4.1 (Trichotomy of The Derivative)

Let $a, b \in \mathbb{R}$ with $a \neq b$. Let $f : [a, b] \rightarrow \mathbb{R}$. SPS that f is differentiable in (a, b) and continuous at a and b .

1. $\forall x \in (a, b) f'(x) \geq 0 \implies f$ is increasing on $[a, b]$.
2. $\forall x \in (a, b) f'(x) > 0 \implies f$ is strictly increasing on $[a, b]$.
3. $\forall x \in (a, b) f'(x) \leq 0 \implies f$ is decreasing on $[a, b]$.
4. $\forall x \in (a, b) f'(x) < 0 \implies f$ is strictly decreasing on $[a, b]$.
5. $\forall x \in (a, b) f'(x) = 0 \implies f$ is constant on $[a, b]$.
6. If $g : [a, b] \rightarrow \mathbb{R}$ is continuous at a and b and differentiable in (a, b) with $g'(x) = f'(x)$ for all $x \in (a, b)$, then for some $c \in \mathbb{R}$ we have $g(x) = f(x) + c$ for all $x \in (a, b)$.

Corollary 4.4.4.2 (The Second Derivative Test)

Let I be an interval in \mathbb{R} , let $f : I \rightarrow \mathbb{R}$ and let $a \in I$. SPS that f is differentiable in I with $f'(a) = 0$.

1. If $f''(a) > 0$ then f has a local minimum at a .
2. If $f''(a) < 0$ then f has a local maximum at a .

4.5 l'Hôpital's Rule

Theorem 4.5.1 (l'Hôpital's Rule)

Let I be a non degenerate interval in \mathbb{R} . Let $a \in I$, or let a be an endpoint of I . Let $f, g : I \setminus \{a\} \rightarrow \mathbb{R}$. SPS that f and g are differentiable in $I \setminus \{a\}$ with $g'(x) \neq 0$ for all $x \in I \setminus \{a\}$. SPS further that $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$ or $\pm \infty$, and suppose that $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} \in \hat{\mathbb{R}}$. Then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

Chapter 5

Integrability on \mathbb{R}

5.1 The Riemann Integral

Definition 5.1.1 (Partition & Subintervals)

A partition of the closed interval $[a, b]$ is a set $X = \{x_0, x_1, \dots, x_n\}$ with

$$a = x_0 < x_1 < x_2 < \dots < x_n = b.$$

The intervals $[x_{i-1}, x_i]$ are called the subintervals of $[a, b]$, and we write

$$\Delta_i x = x_i - x_{i-1}$$

for the size of the i^{th} subinterval. Note that

$$\sum_{i=1}^n \Delta_i x = b - a$$

The size (or norm) of the partition X , denoted by ΔX is

$$\Delta X = \max\{\Delta_i x | 1 \leq i \leq n\}$$

i.e. ΔX is the largest subinterval in X .

Definition 5.1.2 (The Riemann Sum)

Let X be a partition of $[a, b]$, and let $f : [a, b] \rightarrow \mathbb{R}$ be bounded. A Riemann sum for f on X is a sum of the form

$$S = \sum_{i=1}^n f(t_i) \Delta_i x \quad \text{for some } t_i \in [x_{i-1}, x_i].$$

The points t_i are called sample points.

Definition 5.1.3 ((Riemann) Integrable)

Let $f : [a, b] \rightarrow \mathbb{R}$ be bounded. We say that f is (Riemann) integrable on $[a, b]$ when

$$\exists I \in \mathbb{R} \forall \epsilon > 0 \exists \delta > 0 \forall X = \{x_0, x_1, \dots, x_n\}$$

$$\Delta X < \delta \implies \left(\forall t_i \in [x_{i-1}, x_i] \quad \left| \sum_{i=1}^n f(t_i) \Delta_i x - I \right| < \epsilon \right)$$

where $X = \{x_0, x_1, \dots, x_n\}$ is a partition on $[a, b]$.

The number I is unique, and it is called the (Riemann) integral of f on $[a, b]$, and we write

$$I = \int_a^b f, \text{ or } I = \int_a^b f(x) dx$$

5.2 Upper and Lower Riemann Sums**Definition 5.2.1 (Upper and Lower Riemann Sums)**

Let X be a partition for $[a, b]$ and let $f : [a, b] \rightarrow \mathbb{R}$ be bounded. The upper Riemann sum for f on X , denoted by $U(f, X)$, is

$$U(f, X) = \sum_{i=1}^n M_i \Delta_i X \quad \text{where } M_i = \sup\{f(t) \mid t \in [x_{i-1}, x_i]\}$$

and the lower Riemann sum for f on X , denoted $L(f, X)$, is

$$L(f, X) = \sum_{i=1}^n m_i \Delta_i X \quad \text{where } m_i = \inf\{f(t) \mid t \in [x_{i-1}, x_i]\}$$

Remark

- $U(f, X)$ and $L(f, X)$ are not necessarily Riemann sums themselves, since we do not always have $M_i = f(t_i)$ or $m_i = f(s_i)$, for any $t_i, s_i \in [x_{i-1}, x_i]$.
- If f is increasing, then $M_i = f(x_i)$ and $m_i = f(x_{i-1})$, and in this case $U(f, X)$ and $L(f, X)$ are Riemann sums.
- Similarly, if f is decreasing, then $M_i = f(x_{i-1})$ and $m_i = f(x_i)$, and $U(f, X)$ and $L(f, X)$ are Riemann sums.
- If f is continuous, then by the **Extreme Value Theorem**, we have $M_i = f(t_i)$ and $m_i = f(s_i)$ for some $t_i, s_i \in [x_{i-1}, x_i]$, and so in this case $U(f, X)$ and $L(f, X)$ are again Riemann sums.

Theorem 5.2.1 (Upper and Lower Riemann Sums and the Riemann Sum)

Let X be a partition of $[a, b]$ and let $f : [a, b] \rightarrow \mathbb{R}$ be bounded. Then

$$U(f, X) = \sup \left\{ \sum_{i=1}^n f(t_i) \Delta_i x \mid t_i \in [x_{i-1}, x_i] \right\}$$

$$L(f, X) = \inf \left\{ \sum_{i=1}^n f(t_i) \Delta_i x \mid t_i \in [x_{i-1}, x_i] \right\}$$

In particular, for every Riemann sum S for f on X we have

$$L(f, X) \leq \sum_{i=1}^n f(t_i) \Delta_i x \leq U(f, X)$$

for every $t_i \in [x_{i-1}, x_i]$.

Theorem 5.2.2 (Partition Refinement)

Let $f : [a, b] \rightarrow \mathbb{R}$ be bounded with upper and lower bounds M and m . Let $X = \{x_0, x_1, \dots, x_n\}$ and $Y = \{y_0, y_1, \dots, y_m\}$ be partitions of $[a, b]$ such that $\{x_0, x_1, \dots, x_n\} \subseteq \{y_0, y_1, \dots, y_m\}$ (or $m \geq n$). Then

$$0 \leq L(f, Y) - L(f, X) \leq (M - m) \Delta X$$

$$0 \leq U(f, X) - U(f, Y) \leq (M - m) \Delta X$$

Remark

Let X and Y be partitions of $[a, b]$ with $X \subset Y$. Then

$$L(f, X) \leq L(f, Y) \leq U(f, Y) \leq U(f, X)$$

Definition 5.2.2 (Upper and Lower Integral)

Let $f : [a, b] \rightarrow \mathbb{R}$ be bounded. The upper integral of f on $[a, b]$, denoted by $U(f)$, is given by

$$U(f) = \inf \{U(f, X) \mid X \text{ is a partition of } [a, b]\}$$

and the lower integral of f on $[a, b]$, denoted by $L(f)$, is given by

$$L(f) = \sup \{L(f, X) \mid X \text{ is a partition of } [a, b]\}.$$

Theorem 5.2.3 (Upper Integral \geq Lower Integral)

Let $f : [a, b] \rightarrow \mathbb{R}$. We always have $L(f) \leq U(f)$.

Theorem 5.2.4 (Equivalent Definitions of Integrability)

Let $f : [a, b] \rightarrow \mathbb{R}$ be bounded. Then TFAE:

1. f is integrable on $[a, b]$.
2. $\forall \epsilon > 0 \exists$ partition X $U(f, X) - L(f, X) < \epsilon$.
3. $L(f) = U(f)$.

5.3 Evaluating Integrals of Continuous Functions

Theorem 5.3.1 (Continuous Functions are Integrable)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then f is integrable on $[a, b]$.

Lemma 5.3.2 (Summation Formulas)

We have

$$\sum_{i=1}^n 1 = n, \quad \sum_{i=1}^n i = \frac{n(n+1)}{2}, \quad \sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}, \quad \sum_{i=1}^n i^3 = \frac{n^2(n+1)^2}{4}$$

5.4 Basic Properties of Integrals

Theorem 5.4.1 (Linearity)

Let f and g be integrable on $[a, b]$ and let $c \in \mathbb{R}$. Then $f + g$ and cf are both integrable on $[a, b]$ and

$$\int_a^b (f + g) = \int_a^b f + \int_a^b g$$

and

$$\int_a^b cf = c \int_a^b f$$

Theorem 5.4.2 (Comparison Theorem for Integrals)

Let f and g be integrable on $[a, b]$. If $f(x) \leq g(x)$ for all $x \in [a, b]$ then

$$\int_a^b f \leq \int_a^b g.$$

Theorem 5.4.3 (Additivity)

Let $a < b < c$ and let $f : [a, c] \rightarrow \mathbb{R}$ be bounded. Then f is integrable on $[a, c]$ iff f is integrable both on $[a, b]$ and on $[b, c]$, and in this case

$$\int_a^b f + \int_b^c f = \int_a^c f.$$

Definition 5.4.1 (Integral at a Point & Integral from the Right)

We define $\int_a^a f = 0$ and for $a < b$ we define $\int_b^a f = -\int_a^b f$.

Note

Using the above definition, the Additivity Theorem extends to the case that $a, b, c \in \mathbb{R}$ are not in increasing order: for any $a, b, c \in \mathbb{R}$, if f is integrable on $[\max\{a, b, c\}, \min\{a, b, c\}]$ then

$$\int_a^b f + \int_b^c f = \int_a^c f.$$

Theorem 5.4.4 (Estimation)

Let f be integrable on $[a, b]$. Then $|f|$ is integrable on $[a, b]$ and

$$\left| \int_a^b f \right| \leq \int_a^b |f|$$

5.5 Fundamental Theorem of Calculus**Note**

For a function F , defined on an interval containing $[a, b]$, we write

$$[F(x)]_a^b = F(b) - F(a).$$

Theorem 5.5.1 (The Fundamental Theorem of Calculus)

1. Let f be integrable on $[a, b]$. Define $F : [a, b] \rightarrow \mathbb{R}$ by

$$F(x) = \int_a^x f = \int_a^x f(t) dt$$

Then F is continuous on $[a, b]$. Moreover, if f is continuous at a point $x \in [a, b]$, then F is differentiable at x and

$$F'(x) = f(x).$$

2. Let f be integrable on $[a, b]$. Let F be differentiable on $[a, b]$ with $F' = f$. Then

$$\int_a^b f = [F(x)]_a^b = F(b) - F(a)$$

Definition 5.5.1 (Antiderivative)

A function F such that $F' = f$ on an interval is called an antiderivative of f on the interval.

Chapter 6

Series of \mathbb{R}

6.1 Introduction

Definition 6.1.1 (Series)

Let $\{a_n\}_{n \geq k}$ be a sequence. The series $\sum_{n \geq k} a_n$ is defined to be the sequence $\{S_l\}_{l \geq k}$ where

$$S_l = \sum_{n=k}^l a_n = a_k + a_{k+1} + a_{k+2} + \dots + a_l$$

The term S_l is called the l^{th} partial sum of the series $\sum_{n \geq k} a_n$. The sum of the series, denoted by

$$S = \sum_{n=k}^{\infty} a_n = a_k + a_{k+1} + a_{k+2} + \dots,$$

is the limit of the sequence of the partial sums, if it exists, and we say that the series converges when the sum exists and is finite.

Example 6.1.1 (Geometric Series)

For $a \neq 0$, the series $\sum_{n \geq k} a r^n$ converges iff $|r| < 1$, and in this case

$$\sum_{n=k}^{\infty} a r^n = \frac{a r^k}{1 - r}.$$

Theorem 6.1.1 (First Finitely Many Terms Do Not Affect Convergence)

Let $\{a_n\}_{n \geq k}$ be a sequence. Then for any integer $m \geq k$, the series $\sum_{n \geq k} a_n$ converges iff

the series $\sum_{n \geq m} a_n$ converges, and in this case

$$\sum_{n=k}^{\infty} a_n = (a_k + a_{k+1} + \dots + a_{m-1}) + \sum_{n=m}^{\infty} a_n.$$

Definition 6.1.2 (Error)

When we approximate a value x by the value y , the error in our approximation is $|x - y|$.

Theorem 6.1.2 (Linearity)

If $\sum a_n$ and $\sum b_n$ are convergent series then

1. for any $c \in \mathbb{R}$, $\sum ca_n$ converges and $\sum_{n=k}^{\infty} ca_n = c \sum_{n=k}^{\infty} a_n$, and
2. the series $\sum(a_n + b_n)$ converges and $\sum_{n=k}^{\infty} (a_n + b_n) = \sum_{n=k}^{\infty} a_n + \sum_{n=k}^{\infty} b_n$

Theorem 6.1.3 (Series of Positive Terms)

Let $\sum a_n$ be a series.

1. If $a_n \geq 0$ for all $n \geq k$ then either $\sum a_n$ converges or $\sum_{n=k}^{\infty} a_n = \infty$.
2. If $a_n \leq 0$ for all $n \geq k$ then either $\sum a_n$ converges or $\sum_{n=k}^{\infty} a_n = -\infty$.

Theorem 6.1.4 (Cauchy Criterion)

Let $\sum a_n$ be a series. Then $\sum a_n$ converges iff

$$\forall \epsilon > 0 \exists N \in \mathbb{Z}_{\geq k} \forall l, m \in \mathbb{Z}_{\geq k} \left(m > l > N \implies \left| \sum_{n=l+1}^m a_n \right| < \epsilon \right)$$

6.2 Convergence Tests

Theorem 6.2.1 (Divergence Test)

If $\sum a_n$ converges then $\lim_{n \rightarrow \infty} a_n = 0$. Equivalently, if $\lim_{n \rightarrow \infty} a_n$ either does not exist, or exists but is not equal to 0, then $\sum a_n$ diverges.

Remark

Converse of the Divergence Test is false. E.g. $\sum \frac{1}{n}$ (Harmonic Series) diverges but $\lim \frac{1}{n} = 0$.

Theorem 6.2.2 (Integral Test)

Let $f(x)$ be positive and decreasing for $x \geq k$, and let $a_n = f(n)$ for all integers $n \geq k$. Then $\sum a_n$ converges iff $\int_k^\infty f(x) dx$ converges, and in this case, for any $l \geq k$ we have

$$\int_{l+1}^\infty f(x) dx \leq \sum_{n=l+1}^\infty a_n \leq \int_l^\infty f(x) dx$$

Corollary 6.2.2.1 (p-Series)

The series

$$\sum_{k=1}^\infty \frac{1}{k^p}$$

converges iff $p > 1$.

Theorem 6.2.3 (Comparison Test)

SPS that $0 \leq a_n \leq b_n$ for all $n \geq k$. Then if $\sum b_n$ converges, then so does $\sum a_n$ and in this case,

$$\sum_{n=k}^\infty a_n \leq \sum_{n=k}^\infty b_n.$$

Theorem 6.2.4 (Limit Comparison Test)

Let $a_n \geq 0$ and let $b_n > 0$ for all $n \geq k$. SPS that $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = r$. Then

1. if $r = \infty$ and $\sum a_n$ converges, then so does $\sum b_n$.
2. if $r = 0$ and $\sum b_n$ converges, then so does $\sum a_n$, and
3. if $0 < r < \infty$, then $\sum a_n$ converges iff $\sum b_n$ converges.

Theorem 6.2.5 (Ratio Test)

Let $a_n > 0$ for all $n \geq k$. SPS $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = r$. Then

1. if $r < 1$, then $\sum a_n$ converges, and
2. if $r > 1$, then $\lim_{n \rightarrow \infty} a_n = \infty$ so $\sum a_n = \infty$

Definition 6.2.1 (Limit Supremum)

The limit supremum of a sequence of real numbers $\{x_k\}$ is defined to be

$$\limsup_{k \rightarrow \infty} x_k := \lim_{n \rightarrow \infty} \left(\sup_{k > n} x_k \right).$$

Theorem 6.2.6 (Root Test)

Let $a_n \geq 0$ for all $n \geq k$. Let $r = \limsup_{n \rightarrow \infty} \sqrt[n]{a_n}$. Then

1. if $r < 1$, then $\sum a_n$ converges, and
2. if $r \not< 1$, then $\lim_{n \rightarrow \infty} a_n = \infty$ so $\sum a_n = \infty$.

Remark

The Root and Ratio Tests are inconclusive when $r = 1$.

Definition 6.2.2 (Alternating Sequence)

A sequence $\{a_n\}_{n \geq k}$ is said to be alternating when either we have $a_n = (-1)^n |a_n|$ for all $n \geq k$ or we have $a_n = (-1)^{n+1} |a_n|$ for all $n \geq k$.

Theorem 6.2.7 (Alternating Series Test)

Let $\{a_n\}_{n \geq k}$ be an alternating series. If $\{|a_n|\}$ is decreasing with $\lim_{n \rightarrow \infty} |a_n| = 0$, then $\sum_{n \geq k} a_n$ converges, and in this case

$$\left| \sum_{n=k}^{\infty} a_n \right| \leq |a_k|$$

Definition 6.2.3 (Converge Absolutely & Converge Conditionally)

A series $\sum_{n \geq k} a_n$ is said to converge absolute when $\sum_{n \geq k} |a_n|$ converges.

The series is said to converge conditionally if $\sum_{n \geq k} a_n$ converges but $\sum_{n \geq k} |a_n|$ diverges.

Theorem 6.2.8 (Absolute Converges \implies Convergence)

If $\sum |a_n|$ converges then so does $\sum a_n$.

6.3 Fubini's Theorem for Series**Definition 6.3.1 (Multiplication of Series)**

SPS that $\sum_{n \geq 0} |a_n|$ and $\sum_{n \geq 0} b_n$ both converge, and define $c_n = \sum_{k=0}^n a_k b_{n-k}$. Then $\sum_{n \geq 0} c_n$ converges and

$$\sum_{n=0}^{\infty} c_n = \left(\sum_{n=0}^{\infty} a_n \right) \left(\sum_{n=0}^{\infty} b_n \right)$$

Theorem 6.3.1 (Fubini's Theorem for Series)

Let $a_{n,m} \in \mathbb{R}$ for all $n, m \geq 0$. SPS that $\sum_{m \geq 0} |a_{n,m}|$ converges for each $n \geq 0$ and that

$\sum_{n \geq 0} \left(\sum_{m=0}^{\infty} |a_{n,m}| \right)$ converges. Then $\sum_{m \geq 0} a_{n,m}$ converges for all $n \geq 0$, $\sum_{n \geq 0} \left(\sum_{m=0}^{\infty} a_{n,m} \right)$ converges, $\sum_{n \geq 0} a_{n,m}$ converges for all $m \geq 0$, $\sum_{m \geq 0} \left(\sum_{n=0}^{\infty} a_{n,m} \right)$ converges, and

$$\sum_{n=0}^{\infty} \left(\sum_{m=0}^{\infty} a_{n,m} \right) = \sum_{m=0}^{\infty} \left(\sum_{n=0}^{\infty} a_{n,m} \right)$$

Chapter 7

Sequences and Series of Functions

7.1 Pointwise Convergence

Definition 7.1.1 (Pointwise Convergence of Sequences of Functions)

Let $I \subset \mathbb{R}$, let $f : I \rightarrow \mathbb{R}$, and for each integer $n \geq k$ let $f_n : I \rightarrow \mathbb{R}$. We say that the sequence of functions $\{f_n\}_{n \geq k}$ converges pointwise to f on I , and write $f_n \rightarrow f$ pointwise on I , when $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ for all $x \in I$, i.e.

$$\forall x \in I \forall \epsilon > 0 \exists N \geq k \forall n \geq k (n > N \implies |f_n(x) - f(x)| < \epsilon)$$

Remark

Not that by the *Cauchy Criterion for Convergence (for sequences)*, the sequence $\{f_n\}_{n \geq k}$ converges pointwise to some function $f(x)$ on I iff

$$\forall x \in I \forall \epsilon > 0 \exists N \geq k \forall n, m \geq k (n, m > N \implies |f_n(x) - f_m(x)| < \epsilon)$$

7.2 Uniform Convergence

Definition 7.2.1 (Uniform Convergence of Sequences of Functions)

Let $I \subset \mathbb{R}$, let $f : I \rightarrow \mathbb{R}$, and for each integer $n \geq k$, let $f_n : I \rightarrow \mathbb{R}$. We say that the sequence of functions $\{f_n\}_{n \geq k}$ converges uniformly to f on I , and write $f_n \rightarrow f$ uniformly on I , when

$$\forall \epsilon > 0 \exists N \geq k \forall x \in I \forall n \geq k (n > N \implies |f_n(x) - f(x)| < \epsilon)$$

Theorem 7.2.1 (Cauchy Criterion for Uniform Convergence - Sequence of Functions)

Let $\{f_n\}_{n \geq k}$ be a sequence of functions on $I \subset \mathbb{R}$. Then $\{f_n\}_{n \geq k}$ converges uniformly to some function f on I

$$\iff \forall \epsilon > 0 \exists N \geq k \forall x \in I \forall n, m \geq N (n, m > N \implies |f_n(x) - f_m(x)| < \epsilon)$$

Theorem 7.2.2 (Uniform Convergence, Limits and Continuity - Sequence of Functions)

SPS that $f_n \rightarrow f$ uniformly on I . Let x be a limit point of I . If $\lim_{y \rightarrow x} f_n(y)$ exists for each n , then

$$\lim_{y \rightarrow x} \lim_{n \rightarrow \infty} f_n(y) = \lim_{n \rightarrow \infty} \lim_{y \rightarrow x} f_n(y)$$

In particular, if each f_n is continuous on I , then so is f .

Remark

The above theorem allows us to “interchange” limit notations.

Theorem 7.2.3 (Uniform Convergence and Integration - Sequence of Functions)

SPS that $f_n \rightarrow f$ uniformly on $[a, b]$. If each f_n is integrable on $[a, b]$ then so is f . In this case, if $g_n(x) = \int_a^x f_n(t) dt$ and $g(x) = \int_a^x f(t) dt$, then $g_n \rightarrow g$ uniformly on $[a, b]$. In particular, we have

$$\lim_{n \rightarrow \infty} \int_a^b f_n(x) dx = \int_a^b \lim_{n \rightarrow \infty} f_n(x) dx$$

Remark

The above theorem can be interpreted as if we have a sequence of integrable functions, f_n on a closed domain $[a, b]$, then the function that the sequence of functions converge uniformly to, i.e. f , is also integrable on $[a, b]$.

Moreover, we can “interchange” the limit and integration signs in the function.

Theorem 7.2.4 (Uniform Convergence and Differentiation - Sequence of Functions)

Let $\{f_n\}$ be a sequence of functions on $[a, b]$. SPS that each f_n is differentiable on $[a, b]$, $\{f'_n\}$ converges uniformly on $[a, b]$, and $\{f_n(c)\}$ converges for some $c \in [a, b]$. Then $\{f_n\}$ converges uniformly on $[a, b]$, $\lim_{n \rightarrow \infty} f_n(x)$ is differentiable, and

$$\frac{d}{dx} \lim_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} \frac{d}{dx} f_n(x)$$

Remark

The above theorem allows us to “interchange” the process of differentiation and taking the limit of a sequence of function.

7.3 Series of Functions

Definition 7.3.1 (Series of Functions)

Let $\{f_n\}_{n \geq k}$ be a sequence of functions on $I \subseteq \mathbb{R}$. The series of functions $\sum_{n \geq k} f_n(x)$ is defined to be the sequence $\{S_l(x)\}$ where $S_l(x) = \sum_{n=k}^l f_n(x)$. The function $S_l(x)$ is called the l^{th} partial sum of the series. We say that the series $\sum_{n \geq k} f_n(x)$ converges pointwise (or uniformly) on I when the sequence $\{S_l\}$ converges, pointwise (or uniformly) on I . In this case, the sum of the series of functions is defined to be the function

$$f(x) = \sum_{n=k}^{\infty} f_n(x) = \lim_{n \rightarrow \infty} S_l = \lim_{n \rightarrow \infty} \sum_{n=k}^l f_n(x)$$

Note

Many of the following theorems have similar proofs to the earlier two sections.

Theorem 7.3.1 (Cauchy Criterion for Uniform Convergence - Series of Functions)

The series $\sum_{n \geq k} f_n(x)$ converges uniformly (to some function f) on I

$$\iff \forall \epsilon > 0 \exists N \geq k \forall x \in I \forall m, l \geq k \left(m > l > N \implies \left| \sum_{n=l+1}^m f_n(x) \right| < \epsilon \right)$$

Theorem 7.3.2 (Uniform Convergence, Limits and Continuity - Series of Functions)

SPS that $\sum_{n \geq k} f_n(x)$ converges uniformly on I . Let x be a limit point of I . If $\lim_{y \rightarrow x} f_n(y)$ exists for all $n \geq k$, then

$$\lim_{y \rightarrow x} \sum_{n=k}^{\infty} f_n(y) = \sum_{n=k}^{\infty} \lim_{y \rightarrow x} f_n(y)$$

In particular, if each $f_n(x)$ is continuous on I , then so is $\sum_{n=k}^{\infty} f_n(x)$.

Theorem 7.3.3 (Term-by-Term Integration)

SPS that $\sum_{n \geq k} f_n(x)$ converges uniformly on $[a, b]$. If each $f_n(x)$ is integrable on $[a, b]$, then so is $\sum_{n \geq k} f_n(x)$. In this case, if $g_n(x) := \int_a^x f_n(t) dt$ and $g(x) := \int_a^x \sum_{n=k}^{\infty} f_n(t) dt$, then $\sum_{n \geq k} g_n(x)$ converges uniformly to $g(x)$ on I . In particular, we have

$$\int_a^b \sum_{n=k}^{\infty} f_n(x) dx = \sum_{n=k}^{\infty} \int_a^b f_n(x) dx$$

Theorem 7.3.4 (Term-by-Term Differentiation)

SPS that each $f_n(x)$ is differentiable on $[a, b]$, $\sum_{n \geq k} f'_n(x)$ converges uniformly on $[a, b]$, and $\sum_{n \geq k} f_n(c)$ converges for some $c \in [a, b]$. Then $\sum_{n \geq k} f_n(x)$ converges uniformly on $[a, b]$, and

$$\frac{d}{dx} \sum_{n=k}^{\infty} f_n(x) = \sum_{n=k}^{\infty} \frac{d}{dx} f_n(x)$$

Theorem 7.3.5 (Weierstrass M-Test)

SPS that f_n is bounded with $|f_n(x)| \leq M_n$ for all $n \geq k$ and $x \in I$, and $\sum_{n \geq k} M_n$ converges. Then $\sum_{n \geq k} f_n(x)$ converges uniformly on I .

7.4 Power Series

Definition 7.4.1 (Power Series)

A power series centered at a is a series of the form $\sum_{n \geq 0} a_n(x - a)^n$ for some real numbers a_n , where we use the convention that $(x - a)^0 = 1$.

Lemma 7.4.1 (Abel's Formula)

Let $\{a_n\}$ and $\{b_n\}$ be sequences. Then we have

$$\sum_{n=m}^l a_n b_n + \sum_{[p=m]}^{l-1} \left(\sum_{n=m}^p a_n \right) (b_{p-1} - b_p) = \left(\sum_{n=m}^l a_n \right) b_l$$

Theorem 7.4.2 (The Interval and Radius of Convergence)

Let $\sum_{n \geq 0} a_n(x - a)^n$ be a power series and let $R = \frac{1}{\limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}} \in [0, \infty)$. Then the set of $x \in \mathbb{R}$ for which the power series converges is an interval I centered at a of radius R . Indeed

1. $|x - a| > R \implies \lim_{n \rightarrow \infty} a_n(x - a)^n \neq 0 \implies \sum_{n \geq 0} a_n(x - a)^n$ diverges,
2. $|x - a| < R \implies \sum_{n \geq 0} a_n(x - a)^n$ converges absolutely,
3. $0 < r < R \implies \sum_{n \geq 0} a_n(x - a)^n$ converges uniformly in $[a - r, a + r]$.

Theorem 7.4.3 (Abel's Theorem)

Let $\sum_{n \geq 0} a_n(x - a)^n$ be a power series and let $R = \frac{1}{\limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}} \in [0, \infty)$. If $\sum_{n \geq 0} a_n(x - a)^n$ converges when $x = a + R$, then the convergence is uniform on $[a, a + R]$, and, similarly, if $\sum_{n \geq 0} a_n(x - a)^n$ converges when $x = a - R$ then the convergence is uniform on $[a - R, a]$.

Definition 7.4.2 (Interval and Radius of Convergence)

The number R in the previous 2 theorems is called the radius of convergence, and the interval I is called the interval of convergence of the power series.

7.5 Operations on Power Series**Theorem 7.5.1 (Continuity of Power Series)**

SPS that the power series $\sum a_n(x-a)^n$ converges in an interval I . Then the sum $f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n$ is continuous on I .

Proof

This follows from *uniform convergence* of $\sum a_n(x-a)^n$ in closed subintervals of I .

Theorem 7.5.2 (Addition and Subtraction of Power Series)

SPS that the power series $\sum a_n(x-a)^n$ and $\sum b_n(x-a)^n$ both converge in the interval I . Then $\sum (a_n + b_n)(x-a)^n$ and $\sum (a_n - b_n)(x-a)^n$ both converge in I , and for all $x \in I$ we have

$$\left(\sum_{n=0}^{\infty} a_n(x-a)^n \right) \pm \left(\sum_{n=0}^{\infty} b_n(x-a)^n \right) = \sum_{n=0}^{\infty} (a_n \pm b_n)(x-a)^n$$

Proof

This follows from *linearity*.

Theorem 7.5.3 (Multiplication of Power Series)

SPS that the power series $\sum a_n(x-a)^n$ and $\sum b_n(x-a)^n$ both converge in an open interval I with $a \in I$. Let $c_n = \sum_{k=0}^n a_k b_{n-k}$. Then $\sum c_n(x-a)^n$ converges in I and for all $x \in I$ we have

$$\sum_{n=0}^{\infty} c_n(x-a)^n = \sum_{k=0}^n a_k b_{n-k} = \left(\sum_{n=0}^{\infty} a_n(x-a)^n \right) \left(\sum_{n=0}^{\infty} b_n(x-a)^n \right)$$

Proof

This follows from *Multiplication of Series*, since the power series converge absolutely on I .

Theorem 7.5.4 (Division of Power Series)

SPS that the power series $\sum a_n(x-a)^n$ and $\sum b_n(x-a)^n$ both converge in an open interval I with $a \in I$ and that $b_0 \neq 0$. Define c_n by

$$c_0 = \frac{a_0}{b_0}, \text{ and for } n > 0, c_n = \frac{a_n}{b_n} - \frac{b_n c_0}{b_0} - \frac{b_{n-1} c_1}{b_{n-1}} - \dots - \frac{b_1 c_{n-1}}{b_1}.$$

Then there is an open interval J with $a \in J$ such that $\sum c_n(x-a)^n$ converges in J and for all $x \in J$,

$$\sum_{n=0}^{\infty} = \frac{\sum_{n=0}^{\infty} a_n(x-a)^n}{\sum_{n=0}^{\infty} b_n(x-a)^n}.$$

Theorem 7.5.5 (Composition of Power Series)

Let $f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n$ in an open interval I with $a \in I$, and let $g(y) = \sum_{m=0}^{\infty} b_m(y-b)^m$ in an open interval J with $b \in J$ and with $a_0 \in J$. Let K be an open interval with $a \in K$ such that $f(K) \subset J$. For each $m \geq 0$, let $c_{n,m}$ be the coefficients, found by multiplying power series, such that $\sum_{n=0}^{\infty} c_{n,m}(x-a)^n = b_m \left(\sum_{n=0}^{\infty} a_n(x-a)^n - b \right)^m$. Then $\sum_{m \geq 0} c_{n,m}$ converges for all $m \geq 0$, and for all $x \in K$, $\sum_{n \geq 0} \left(\sum_{m=0}^{\infty} c_{n,m} \right) (x-a)^n$ converges and

$$\sum_{n=0}^{\infty} \left(\sum_{m=0}^{\infty} c_{n,m} \right) (x-a)^n = g(f(x))$$

Proof

This follows from *Fubini's Theorem for Series* since

$$g(f(x)) = \sum_{m=0}^{\infty} b_m(f(x)-b)^m = \sum_{m=0}^{\infty} \left(\sum_{n=0}^{\infty} a_n(x-a)^n - b \right)^m = \sum_{m=0}^{\infty} \left(\sum_{n=0}^{\infty} c_{n,m}(x-a)^n \right).$$

Theorem 7.5.6 (Integration of Power Series)

SPS that $\sum a_n(x-a)^n$ converges in the interval I . Then $\forall x$ in I , the sum $f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n$ is integrable on $[\min\{a, x\}, \max\{a, x\}]$ and

$$\int_a^x \sum_{n=0}^{\infty} a_n(t-a)^n dt = \sum_{n=0}^{\infty} \int_a^x a_n(t-a)^n dt = \sum_{n=0}^{\infty} \frac{a_n}{n+1} (x-a)^{n+1}.$$

Theorem 7.5.7 (Differentiation of Power Series)

SPS that $\sum a_n(x-a)^n$ converges in the interval I . Then the sum $f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n$ is differentiable in I and

$$f'(x) = \sum_{n=1}^{\infty} a_n(x-a)^{n-1}.$$

7.6 Taylor Series

Theorem 7.6.1 (Coefficients of the Taylor Series)

SPS that $\sum_{n=0}^{\infty} a_n(x-a)^n$ in an open interval I centered at a . Then f is infinitely differentiable at a and for all $n \geq 0$, we have

$$a_n = \frac{f^{(n)}(a)}{n!},$$

where $f^{(n)}(a)$ denotes the n -th derivative of f at a .

Proof

By repeatedly applying *Differentiation on Power Series*, we can obtain our desired result.

Definition 7.6.1 (Taylor Series and Polynomial)

Given a function $f(x)$ whose derivatives of all order exists at $x = a$, we define the Taylor series of $f(x)$ centered at a to be the power series

$$T(x) = \sum_{n=0}^{\infty} a_n(x-a)^n \quad \text{where } a_n = \frac{f^{(n)}(a)}{n!}$$

and we define the l^{th} Taylor Polynomial of $f(x)$ centered at a to be the l^{th} partial sum

$$T_l(x) = \sum_{n=0}^l a_n(x-a)^n \quad \text{where } a_n = \frac{f^{(n)}(a)}{n!}.$$

Theorem 7.6.2 (Taylor's Theorem)

Let $f(x)$ be infinitely differentiable in an open interval I with $a \in I$. Let $T_l(x)$ be the l^{th} Taylor polynomial for $f(x)$ centered at a . Then for all $x \in I$ there exists a number c between a and x such that

$$f(x) - T_l(x) = \frac{f^{(l+1)}(c)}{(l+1)!}(x-a)^{l+1}.$$

Theorem 7.6.3 ((?) Accuracy of the Maclaurin Expansion)

The functions e^x , $\sin x$ and $(1+x)^p$ are all exactly equal to the sum of their Taylor series centered at 0 in the interval of convergence.

Chapter 8

Topology in Euclidean Space

Appendix A

ZF Set Theory and the Axiom of Choice

A.1 Introduction

Example A.1.1 (Russel's Paradox)

Let X be the set of all sets, and let $S = \{A \in X \mid A \notin A\}$.

Note for example that $Z \notin Z \implies Z \in S$, and $X \in X \implies X \notin S$.

Thus we have $S \in S \iff S \notin S$.

To ensure that mathematical paradoxes (like the above) can no longer arise, mathematicians considered the following questions, and with these questions, rough answers are provided:

1. What exactly is an allowable mathematical object?

A: Every mathematical object is a mathematical set, and a mathematical set can be constructed using certain rules, for e.g. the now widely accepted Zermelo-Fraenkel Set Theory and the Axiom of Choice. While the Axiom of Choice is still highly criticized even today (e.g. the highly controversial **Banach-Tarski Paradox**), the Zermelo-Fraenkel Set Theory is widely welcomed, but not without critics. We shall call the Zermelo-Fraenkel Set Theory and the Axiom of Choice as the ZFC Axioms of Set Theory.

2. What exactly is an allowable mathematical statement?

A: Every mathematical statement can be expressed in a formal symbolic language, which uses symbols rather than words from any spoken language.

3. What exactly is allowable in a mathematical proof?

A: Every mathematical proof is a finite list of ordered pairs $(\mathcal{S}_n, \mathcal{F}_n)$ (which we can think of as proven theorems), where each \mathcal{S}_n is a finite set of formulas (called the *premises*) and each \mathcal{F}_n is a single formula (called the *conclusion*), which that each pair $(\mathcal{S}_n, \mathcal{F}_n)$ can be obtained from previous pairs $(\mathcal{S}_i, \mathcal{F}_i)$ with $i < n$, using certain proof rules.

In the remainder of this appendix, we shall look more into the first 2 questions.

A.2 ZFC Axioms of Set Theory

Definition A.2.1 (Mathematical Symbols)

We allow ourselves to use only the following symbols from the following symbol set:

\neg	<i>not</i>
\wedge	<i>and</i>
\vee	<i>or</i>
\implies	<i>implies</i>
\iff	<i>if and only if</i>
$=$	<i>equals</i>
\in	<i>is an element of</i>
\forall	<i>for all</i>
\exists	<i>there exists</i>
$() \ \{ \} \ \square$	<i>parenthesis</i>

along with some variable symbols such as x, y, z, u, v, w, \dots or x_1, x_2, x_3, \dots

Definition A.2.2 (Formula)

A formula (in the formal symbolic language of first order set theory) is a non-empty finite string of symbols, from the above list, which can be obtained using finitely many applications following the three rules below:

1. If x and y are variable symbols, then each of the following strings are formulas.

$$x = y, \quad x \in y$$

2. If F and G are formulas then each of the following strings are formulas.

$$\neg F, \quad (F \wedge G), \quad (F \vee G), \quad (F \implies G), \quad (F \iff G)$$

3. If x is a variable symbol and F is a formula then each of the following is a formula.

$$\forall x \in F, \quad \exists x \in F$$

Definition A.2.3 (Free or Bounded Variable)

Let x be a variable symbol and let F be a formula. For each occurrence of the symbol x , which does not immediately follow a quantifier, in the formula F , we define whether the occurrence of x is free or bound inductively as follows:

1. If F is a formula of one of the forms $y = z$ or $y \in z$, where y and z are variable symbols (possibly equal to x), then every occurrence of x in F is free, and no occurrence is bound.
2. If F is a formula of one of the forms $\neg H, (H \wedge G), (H \vee G), (H \implies G), (H \iff G)$, where G and H are formulas, then each occurrence of the symbol x is either an occurrence in the formula G or an occurrence in the formula H , and each free (respectively, bound) occurrence of x in G remains free (respectively, bound) in F , and similarly for each free (or bound) occurrence of x in H . In other words, wlog, if x is bounded in G , then it is bounded in F , and vice versa.
3. If F is a formula of one of the forms $\forall y \in G$ or $\exists y \in G$, where G is a formula and y is a variable symbol. If y is different from x , then each free (or bound) occurrence of x in G remains free (or bound) in the formula G , and if $y = x$ then every free occurrence of x in G becomes bound in F , and every bound occurrence of x in G remains bound in F .

Definition A.2.4 (Is Bound By and Binds)

When a quantifier symbol occurs in a given formula F , and is followed by the variable symbol x and then by the formula G , any free occurrence of x in G will become bound in the given formula F (by the 3rd definition above). We shall say that the occurrence of x is bound by (that occurrence of) the quantifier symbol, or that (the occurrence of) the quantifier symbol binds the occurrence of x .

Definition A.2.5 (Free Variable, Statement, Statement About)

A **free variable** in a formula F is any variable symbol that has at least one free occurrence in F . A formula F with no free variables is called a **statement**. When the free variables in F all lie in the set $\{x_1, x_2, \dots, x_n\}$, we shall write F as $F(x_1, x_2, \dots, x_n)$ and we shall say that F is a **statement about** the variables x_1, x_2, \dots, x_n .

Definition A.2.6 (Unique Existence)

When $F(x)$ is a statement about x , we sometimes write $F(y)$ as a short form for the formula $\forall x(x = y \implies F(x))$, and we sometimes write

$$\exists! y \quad F(y)$$

which we read as "there exists a unique y such that $F(y)$ ", as a short form for the formula

$$(\exists y \quad F(y) \wedge \forall z \quad F(z)) \implies z = y$$

which is, in turn, for the formula

$$\exists y \left(\forall x (x = y \implies F(x)) \wedge \forall z (\forall x (x = z \implies F(x)) \implies z = y) \right)$$

Remark (The ZFC Axioms of Set Theory (informal))

Every mathematical set can be constructed using specific rules, which we shall use the ZFC Axioms of Set Theory. Below is a list of the ZFC Axioms, stated informally.

- *Empty Set Axiom:* There exists an empty set \emptyset with no elements.
- *Extension Axiom:* 2 sets are equal if and only if they have the same elements.
- *Separation Axiom:* If u is a set and $F(x)$ is a statement about x , $\{x \in u : F(x)\}$ is a set.
- *Pair Axiom:* If u and v are sets then $\{u, v\}$ is a set.
- *Union Axiom:* If u is a set then $\bigcup_{v \in u} v$ is a set.
- *Power Set Axiom:* If u is a set then $\mathcal{P}(u) = \{v : v \subseteq u\}$ is a set.
- *Axiom of Infinity:* If we define the natural numbers to be the sets $0 = \emptyset, 1 = \{0\}, 2 = \{0, 1\}, 3 = \{0, 1, 2\}$ and so on, then $\mathbb{N} = \{0, 1, 2, 3, \dots\}$ is a set.
- *Replacement Axiom:* If u is a set and $F(x, y)$ is a statement about x and y with the property that $\forall x \exists! y F(x, y)$ then $\{y : \exists x \in u F(x, y)\}$ is a set.
- *Axiom of Choice:* Given a set u of non-empty pairwise disjoint sets, there exists a set which contains exactly one element from each of the sets in u .

Definition A.2.7 (Empty Set Axiom)

The Empty Set Axiom is the formula

$$\exists u \forall x \neg x \in u$$

Definition A.2.8 (Extension Axiom)

The Extension Axiom is the formula

$$\forall u \forall v \left(u = v \iff \forall x (x \in u \iff x \in v) \right)$$

Theorem A.2.1 (Uniqueness of the Empty Set)

The empty set is unique.

Definition A.2.9 (\emptyset)

We denote the unique empty set by \emptyset .

Definition A.2.10 (Subset)

Given sets u and v , we say that u is a **subset** of v , and write $u \subseteq v$, when $\forall x(x \in u \implies x \in v)$

Definition A.2.11 (Separation Axiom)

For any statement $F(x)$ about x , the following formula is an axiom.

$$\forall u \exists v \forall x (x \in v \iff (x \in u \wedge F(x)))$$

More generally, for any statement $F(x, u_1, u_2, \dots, u_n)$ about x, u_1, u_2, \dots, u_n where $n \geq 0$, the following formula is an axiom.

$$\forall u \forall u_1 \dots \forall u_n \exists v \forall x (x \in v \iff (x \in u \wedge F(x, u_1, \dots, u_n)))$$

Any axiom of this form is called the *Separation Axiom*.

Note

It is important to realize that a Separation Axiom only allows us to construct a subset of a given set u . So, e.g., we cannot use the Separation Axiom to show that the collection $S = \{x : \neg x \in x\}$, which is used to formulate *Russel's Paradox*, is a set.

Definition A.2.12 (Pair Axiom)

The Pair Axiom is the formula

$$\forall u \forall v \exists w \forall x (x \in w \iff (x = u \vee x = v))$$

Definition A.2.13 (Union Axiom)

The Union Axiom is the formula

$$\forall u \exists w \forall x (x \in w \iff \exists v (v \in u \wedge x \in v))$$

Definition A.2.14 (Union)

Given a set u , by the Union Axiom there exists a set w with the property that $\forall x (x \in w \iff \exists v (v \in u \wedge x \in v))$, and by the Extension Axiom, this set w is unique. We call the set w the **union** of the elements in u , and denote it by

$$\cup u = \bigcup_{v \in u} v.$$

Given two sets u and v , we define the union of u and v to be the set

$$u \cup v := \bigcup \{u, v\}.$$

Given three sets u , v , and w , note that $\{z\} = \{z, z\}$ is a set and so $\{x, y, z\} = \{x, y\} \cup \{z\}$ is also a set. More generally, if u_1, u_2, \dots, u_n are sets then $\{u_1, u_2, \dots, u_n\}$ is a set and we define the union of the sets u_1, u_2, \dots, u_n to be

$$u_1 \cup u_2 \cup \dots \cup u_n = \bigcup_{k=1}^n u_k = \bigcup \{u_1, u_2, \dots, u_n\}$$

Definition A.2.15 (Intersection)

Given a set u , we define the intersection of the elements in u to be the set

$$\bigcap u = \left\{ x \in \bigcup u \mid \forall v (v \in u \implies x \in v) \right\}$$

Given two sets u and v , we define the intersection of u and v to be the set

$$u \cap v = \bigcap \{u, v\}$$

and more generally, given sets u_1, u_2, \dots, u_n , we define the intersection of u_1, u_2, \dots, u_n to be the set

$$u_1 \cap u_2 \cap \dots \cap u_n = \bigcap_{k=1}^n u_k = \bigcap \{u_1, u_2, \dots, u_n\}$$

Definition A.2.16 (Power Set Axiom)

The Power Set Axiom is the formula

$$\forall u \exists w \forall v (v \in w \iff v \subseteq u)$$

Definition A.2.17 (Power Set)

Given a set u , the set w is with the property that $\forall v (v \in w \iff v \subseteq u)$ (which exists by the Power Set Axiom and is unique by the Extension Axiom) is called the power set of u and is denoted by $\mathcal{P}(u)$, so we have

$$\mathcal{P}(u) = \{v \mid v \subseteq u\}$$

Definition A.2.18 (Ordered Pair)

Given two sets x and y , we define the ordered pair (x, y) to be the set

$$(x, y) = \{\{x\}, \{x, y\}\}.$$

Given two sets u and v , note that if $x \in u$ and $y \in v$ then we have $\{x\} \in \mathcal{P}(u \cup v)$ and $\{x, y\} \in \mathcal{P}(u \cup v)$ and so $(x, y) = \{\{x\}, \{x, y\}\} \in \mathcal{P}(\mathcal{P}(u \cup v))$. We define the product $u \times v$ to be the set

$$u \times v = \{(x, y) \mid x \in u \wedge y \in v\},$$

i.e.

$$u \times v = \left\{ z \in \mathcal{P}(\mathcal{P}(u \cup v)) \mid \exists x \exists y ((x \in u \wedge y \in v) \wedge z = (x, y)) \right\}$$

Definition A.2.19 (Successor, Inductive)

We define

$$0 = \emptyset, \quad 1 = \{0\}, \quad 2 = \{0, 1\} = 1 \cup \{1\}, \quad 3 = \{0, 1, 2\} = 2 \cup \{2\},$$

and so on. For a set x , we define the successor of x to be the set

$$x + 1 = x \cup \{x\}.$$

A set u is called inductive when it has the property that

$$(0 \in u \wedge \forall x(x \in u \implies x + 1 \in u))$$

Definition A.2.20 (Axiom of Infinity)

The Axiom of Infinity is the formula

$$\exists u(0 \in u \wedge \forall x(x \in u \implies x + 1 \in u))$$

so the Axiom of Infinity states that there exists an inductive set.

Theorem A.2.2 (Existence & Uniqueness of an Inductive Set)

$\exists w := \{x | x \in v \text{ for every inductive set } v\}$

Moreover, this set w is an inductive set.

Definition A.2.21 (Natural Numbers)

The unique set w in the above theorem is called the set of natural numbers, and we denote it by \mathbb{N} . We write

$$\begin{aligned} \mathbb{N} &= \{x | x \in v \text{ for every inductive set } v\} \\ &= \{0, 1, 2, 3, \dots\} \end{aligned}$$

For $x, y \in \mathbb{N}$, we write $x \dot{\prec} y$ when $x \in y$ and write $x \leq y$ when $x < y \vee x = y$.

Remark

For a formula F , we write $\forall x \in u F$ as a shorthand notation for the formula $\forall x(x \in u \implies F)$. Similarly, we write $\exists x \in u F$ as a shorthand notation for $\exists x(x \in u \wedge F)$.

Theorem A.2.3 (Principle of Induction)

Let $F(x)$ be a statement about x . SPS that

1. $F(0)$, and
2. $\forall x \in \mathbb{N}(F(x) \implies F(x + 1))$.

Then $\forall x \in \mathbb{N} F(x)$

Remark

The expression $F(0)$ is short for $\forall x(x = 0 \implies F(x))$, which in turn is short for $\forall x(\forall y \neg y \in x \implies F(x))$. Similarly, $F(x + 1)$ is short for the formula $\forall y(y = x + 1 \implies F(y))$, where $F(y)$ is short for $\forall x(x = y \implies F(x))$.

Definition A.2.22 (Replacement Axiom)

Given a statement $F(x, y)$ about x and y , the following formula is an axiom:

$$\forall u \left(\forall x \exists! y F(x, y) \implies \exists w \forall y (y \in w \iff \exists x \in u F(x, y)) \right)$$

where $\exists! y F(x, y)$ is short for $\exists y (F(x, y) \wedge \forall z (F(x, z) \implies z = y))$ with $F(x, z)$ short for the formula $\forall y (y = z \implies F(x, y))$. More generally, given a statement $F(x, y, u_1, \dots, u_n)$ about x, y, u_1, \dots, u_n with $n \geq 0$, the following formula is an axiom:

$$\forall u \forall u_1 \dots \forall u_n \left(\forall x \exists! y F(x, y, u_1, \dots, u_n) \implies \exists w \forall y (y \in w \iff \exists x \in u F(x, y, u_1, \dots, u_n)) \right)$$

An axiom of this form is called a Replacement Axiom.

Definition A.2.23 (Axiom of Choice)

The Axiom of Choice is the formula given by

$$\forall u \left(\left(\neg \phi \in u \wedge \forall x \in u \forall y \in u (\neg x = y \implies x \cap y = \emptyset) \right) \implies \exists w \forall v \in u \exists! x \in v x \in w \right)$$

From this point on, we will be using upper-case letters to denote sets, instead of lower-case as per the statements above.

A.3 Relations, Equivalence Relations, Functions and Recursion

Definition A.3.1 (Binary Relation)

A binary relation R on a set X is a subset $R \subseteq X \times X$. More generally, a binary relation is any set R whose elements are ordered pairs. For a binary relation R , we usually write xRy instead of $(x, y) \in R$.

Definition A.3.2 (Domain, Range, Image, Inverse Image, Inverse, Composition)

Let R and S be binary relations.

The domain of R is

$$\text{Domain}(R) = \{x \mid \exists y xRy\}$$

and the range of R is

$$\text{Range}(R) = \{x | \exists y \, xRy\}.$$

For any set A , the image of A under R is

$$R(A) = \{y | \exists x \in A \, xRy\}$$

and the inverse image of A under R is

$$R^{-1}(A) = \{x | \exists y \in A \, xRy\}.$$

The inverse of R is

$$R^{-1} = \{(y, x) | (x, y) \in R\}$$

and the composition S composed with R is

$$S \circ R = \{(x, z) | \exists y \, xRy \wedge ySz\}$$

Theorem A.3.1 (Domain, Range, Image and Inverse Image as Sets)

Let A be a set and let R be a binary relation. Then $\text{Domain}(R)$, $\text{Range}(R)$, $R(A)$ and $R^{-1}(A)$ are sets.

Theorem A.3.2 (Inverse and Composition as Binary Relations)

Let A be a set and let R and S be binary relations. Then R^{-1} and $S \circ R$ are binary relations.

Definition A.3.3 (Equivalence Relation)

An equivalence relation on a set X is a binary relation R on X such that

1. R is **reflexive**, i.e. $\forall x \in X \, xRx$
2. R is **symmetric**, i.e. $\forall x, y \in X \, (xRy \implies yRx)$, and
3. R is **transitive**, i.e. $\forall x, y, z \in X \, ((xRy \wedge yRz) \implies xRz)$.

Definition A.3.4 (Equivalence Class)

Let R be an equivalence relation on the set X . For $a \in X$, the equivalence class of a modulo R is the set

$$[a]_R = \{x \in X | xRa\}$$

Definition A.3.5 (Partition)

A partition of a set X is a set S of non-empty pairwise disjoint sets whose union is X , that is a set S such that

1. $\forall X, Y \in S \, (X \neq Y \implies X \cap Y = \emptyset)$
2. $\bigcup S = X$.

Theorem A.3.3 (Correspondence of Equivalence Relations and Partitions)

Given a set X , we have the following correspondence between equivalence relations on X and partitions of X .

1. Given an equivalence relation R on X , the set of all equivalence classes

$$S_R = \{[a]_R | a \in X\}$$

is a partition of X .

2. Given a partition S of X , the relation R_S on X is defined by

$$R_S = \{(x, y) \in X \times X | \exists A \in S (x \in A \wedge y \in A)\}$$

is an equivalence relation on X .

3. Given an equivalence relation R on X we have $R_{S_R} = R$, and a given partition S of X , we have $S_{R_S} = S$.

Note (Set of All Equivalence Classes)

Given an equivalence relation R on X , the set of all equivalence classes, which we denote by S_R in the above theorem, is usually denoted by X/R , so

$$X/R = \{[a]_R | a \in X\}$$

Definition A.3.6 (Set of Representatives)

Let R be an equivalence relation. A set of representatives for R is a subset of X which contains exactly one element from each equivalence class in X/R .

Remark

Notice that the AC is equivalent to the statement that every equivalence relation has a set of representatives.

Definition A.3.7 (Function)

Get sets X and Y , a function from X to Y is a binary relation $f \subseteq X \times Y$ with the property that

$$\forall x \in X \exists! y \in Y (x, y) \in f$$

More generally, a function is a binary relation with the property that

$$\forall x \in \text{Domain}(f) \exists! y (x, y) \in f.$$

For a function f , we usually write $y = f(x)$ instead of xy . It is customary to use the notation $f : X \rightarrow Y$ when $X = \text{Domain}(f)$ and Y is any set with $\text{Range}(f) \subseteq Y$.

Definition A.3.8 (One-to-one & Onto)

Let $f : X \rightarrow Y$. The function f is called *one-to-one* (or *injective*) when

$$\forall y \in Y \exists \text{ at most one } x \in X \ y = f(x)$$

and f is called *onto* (or *surjective*) when

$$\forall y \in Y \exists \text{ at least one } x \in X \ y = f(x)$$

Definition A.3.9 (Left and Right Inverses)

Let $f : X \rightarrow Y$. Let I_X and I_Y denote the identity function on X and Y respectively. A *left inverse* of f is a function $g : Y \rightarrow X$ such that $g \circ f = I_X$. A *right inverse* of f is a function $H : X \rightarrow Y$ such that $f \circ H = I_Y$. Note that if f has a left inverse g and a right inverse H , then we have $g = g \circ I_Y = g \circ f \circ H = I_X \circ H = H$. In this case, we say that g is the (unique two-sided) inverse of f .

Theorem A.3.4 (Surjective and Injective VS Inverses)

Let $f : X \rightarrow Y$. Then

1. f is one-to-one if and only if f has a left inverse.
2. f is onto if and only if f has a right inverse.
3. f is one-to-one and onto if and only if f has a (two-sided) inverse.

Definition A.3.10 (Invertible)

A function $f : X \rightarrow Y$ is called *invertible* (or *bijective*) when it is one-to-one and onto, or equivalently, when it has a (unique two-sided) inverse.

Theorem A.3.5 (The Recursion Theorem)

1. Let A be a set, let $a \in A$, and let $g : A \times \mathbb{N} \rightarrow A$. Then there exists a unique function $f : \mathbb{N} \rightarrow A$ such that

$$f(0) = a \text{ and } f(n+1) = g(f(n), n) \text{ for all } n \in \mathbb{N}$$

2. Let A and B be sets, let $g : A \rightarrow B$, and let $h : A \times B \times \mathbb{N} \rightarrow B$. Then there exists a unique function $f : A \times \mathbb{N} \rightarrow B$ such that for all $a \in A$ we have

$$f(a, 0) = g(a) \text{ and } f(a, n+1) = h(a, f(a, n), n) \text{ for all } n \in \mathbb{N}$$

A.4 Construction of Integers, Rational, Real and Complex Numbers

Definition A.4.1 (Sum and Product)

By Part(2) of the *Recursion Theorem*, there is a unique function $s : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ such that for all $a, b \in \mathbb{N}$ we have

$$s(a, 0) = a, \quad s(1, b + 1) = s(a, b) + 1.$$

We call $s(a, b)$ the sum of a and $b \in \mathbb{N}$ and write it as

$$a + b = s(a, b).$$

Also, there is a unique function $p : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ such that for all $a, b \in \mathbb{N}$ we have

$$p(a, 0) = 0, \quad p(a, b + 1) = p(a, b) + a$$

We call $p(a, b)$ the product of a and b in \mathbb{N} , and we write it as

$$a \cdot b = p(a, b)$$

Definition A.4.2 (Integers)

We define the set of integers to be the set

$$\mathbb{Z} = (\mathbb{N} \times \mathbb{N})/R$$

where R is the equivalence relation given by

$$(a, b)R(c, d) \iff a + d = b + c$$

For $[(a, b)]$ and $[(c, d)]$ in \mathbb{Z} , we define

$$\begin{aligned} [(a, b)] \leq [(c, d)] &\iff b + c \leq a + d \\ [(a, b)] + [(c, d)] &\iff [(a + c, b + d)] \\ [(a, b)] \cdot [(c, d)] &= [(ac + bd, ad + bc)] \end{aligned}$$

For $n \in \mathbb{N}$ we write $n = [(n, 0)]$ and $-n = [(0, n)]$, so that every element of \mathbb{Z} can be written as $\pm n$ for some $n \in \mathbb{N}$, and we can identify \mathbb{N} with a subset of \mathbb{Z}

Definition A.4.3 (Rational Numbers)

We define the set of rational numbers to be the set

$$\mathbb{Q} = (\mathbb{N} \times \mathbb{Z}^+)/R$$

where $\mathbb{Z}^+ = \{x \in \mathbb{N} | x \neq 0\}$ and R is the equivalence relation given by

$$(a, b)R(c, d) \iff ad = bc$$

For $[(a, b)]$ and $[(c, d)]$ in \mathbb{Q} we define

$$\begin{aligned} [(a, b)] \leq [(c, d)] &\iff a \cdot d \leq b \cdot c \\ [(a, b)] + [(c, d)] &\iff [(a \cdot d + b \cdot c, b \cdot d)] \\ [(a, b)] \cdot [(c, d)] &= [(ac, bd)] \end{aligned}$$

For $a \in \mathbb{N}$ and $b \in \mathbb{Z}^+$, it is customary to write $\frac{a}{b} = [(a, b)]$. Also for $a \in \mathbb{Z}$ we write $a = [(a, 1)]$, and we identify \mathbb{Z} with a subset of \mathbb{Q}

Definition A.4.4 (Real Numbers)

We define the set of real numbers to be the set

$$\mathbb{R} = \{x \subseteq \mathbb{Q} | x \neq \emptyset, x \neq \mathbb{Q}, \forall a \in x \forall b \in \mathbb{Q} (b \leq a \implies b \in x), \forall a \in x \exists b \in x a < b\}$$

For $x, y \in \mathbb{R}$ we define

$$\begin{aligned} x \leq y &\iff x \subseteq y \\ x + y &= \{a + b | a, b \in \mathbb{Q}, a \in x, b \in y\} \end{aligned}$$

For $0 \leq x, y \in \mathbb{R}$ we define

$$x \cdot y = \{a \cdot b | 0 \leq a, b \in \mathbb{Q}, a \in x, b \in y\} \cup \{c \in \mathbb{Q} | c < 0\},$$

and YOU can try to, similarly, define $x \cdot y$ in the case that $x \not\leq 0$ and $y \not\leq 0$.

Definition A.4.5 (Complex Numbers)

We define the set of complex numbers to be the set

$$\mathbb{C} = \mathbb{R} \times \mathbb{R}.$$

We define addition and multiplication in \mathbb{C} by

$$\begin{aligned} (a, b) + (c, d) &= (a + c, b + d) \\ (a, b) \cdot (c, d) &= (ac - bd, ad + bc). \end{aligned}$$

We write $i = (0, 1)$. For $x \in \mathbb{R}$ we write $x = (x, 0)$ and identify \mathbb{R} with a subset of \mathbb{C} .

Appendix B

Functions and Cardinality

B.1 Functions

Definition B.1.1 (Range, Image, and Inverse Image)

Let X and Y be sets and let $f : X \rightarrow Y$. Recall (see *Function in Appendix A*) that the domain of f and the range of f are the sets

$$\text{Domain}(f) = X, \quad \text{Range}(f) = f(X) = \{f(x) | x \in X\}$$

For $A \subseteq X$, the image of A under f is the set

$$f(A) = \{f(x) | x \in A\}$$

For $B \subseteq Y$, the inverse image of B under f is the set

$$f^{-1}(B) = \{x \in X | f(x) \in B\}$$

Definition B.1.2 (Composite Function)

Let X , Y and Z be sets. Let $f : X \rightarrow Y$ and let $g : Y \rightarrow Z$. We define the composite function $g \circ f : X \rightarrow Z$ by $(g \circ f)(x) = g(f(x))$ for all $x \in X$

Definition B.1.3 (Bijection)

Let X and Y be sets. Let $f : X \rightarrow Y$. We say that f is a bijection, or that f is bijective, if f is both one-to-one and onto (or that f is both injective and surjective).

Theorem B.1.1 (Bijectiveness and Inverse of the Composite Function)

Let X , Y and Z be sets. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$. Then

1. if f and g are both injective then so is $g \circ f$,

2. if f and g are both surjective then so is $g \circ f$, and
3. if f and g are both invertible then so is $g \circ f$, and in this case $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$

Definition B.1.4 (Identity Function)

For a set X , we define the identity function on X to be the function $I_X : X \rightarrow X$ given by $I_X(x) = x$ for all $x \in X$. Note that for $f : X \rightarrow Y$ we have $f \circ I_X = f$ and $I_Y \circ f = f$.

Theorem B.1.2 (Bijectiveness and Invertability of Functions)

Let X and Y be nonempty sets and let $f : X \rightarrow Y$. Then

1. f is injective if and only if f has a left inverse,
2. f is surjective if and only if f has a right inverse, and
3. f is bijective if and only if f has a left inverse g and a right inverse h , and in this case we have $g = h = f^{-1}$.

Corollary B.1.2.1 (Relationship between Injection and Surjection)

Let X and Y be sets. Then there exists an injective map $f : X \rightarrow Y$ if and only if there exists a surjective map $g : Y \rightarrow X$.

B.2 Cardinality

Definition B.2.1 (Equal Cardinality)

Let A and B be sets. We say that A and B have the same cardinality, and write $|A| = |B|$, when there exists a bijective map $f : A \rightarrow B$.

We say that the cardinality of A is less than or equal to the cardinality of B , and write $|A| \leq |B|$, when there exists an injective map $f : A \rightarrow B$.

We say that the cardinality of A is less than the cardinality of B , and write $|A| < |B|$, when $|A| \leq |B| \wedge |A| \neq |B|$ (i.e. there exists an injective map from A to B but no surjective map from A to B).

We also write $|A| \geq |B|$ when $|B| \leq |A|$ and $|A| > |B|$ when $|B| < |A|$.

Definition B.2.2 (Properties for Cardinality of Sets)

For all sets A , B , and C ,

1. $|A| = |A|$,
2. if $|A| = |B|$, then $|B| = |A|$,
3. if $|A| = |B|$ and $|B| = |C|$, then $|A| = |C|$,

4. $|A| \leq |B| \iff (|A| = |B| \vee |A| < |B|)$, and
5. $|A| \leq |B| \wedge |B| \leq |C| \implies |A| \leq |C|$.

Definition B.2.3 (Finiteness and Countability of Sets)

Let A be a set. For each $n \in \mathbb{N}$, let $S_n = \{0, 1, 2, \dots, n-1\}$. For $n \in \mathbb{N}$, we say that the cardinality of A is equal to n , or that A has n elements, and write $|A| = n$, when $|A| = |S_n|$. We say that A is finite when $|A| = n$ for some $n \in \mathbb{N}$. We say that A is infinite when A is not finite. We say that A is countable when $|A| = |\mathbb{N}|$.

Remark

Note that a set A is said to be countable when A is of the form $A = \{a_0, a_1, a_2, \dots\}$ where all its elements are distinct.

Theorem B.2.1

Let A be a set. Then the following are equivalent.

1. A is infinite.
2. A contains a countable subset.
3. $|\mathbb{N}| \leq |A|$
4. There exists a map $f : A \rightarrow A$ which is injective but not surjective.

Corollary B.2.1.1

Let A and B be sets.

1. If A is countable then A is infinite.
2. When $|A| \leq |B|$, if B is finite then so is A , and if A is infinite, so is B .
3. If $|A| = n$ and $|B| = m$, then $|A| \leq |B|$ iff $n \leq m$.
4. If $|A| = n$ and $|B| = m$, then $|A| \leq |B| \iff n \leq m$.
5. When one of the two sets A or B is finite. If $|A| \leq |B| \wedge |B| \leq |A| \implies |A| = |B|$.

Theorem B.2.2 ($|\mathbb{N}|$ as a Threshold for Finiteness and Countability)

Let A be a set. $|A| \leq |\mathbb{N}| \iff A$ is finite or countable.

Theorem B.2.3

Let A be a set. Then

1. $|A| < |\mathbb{N}| \iff A$ is finite,
2. $|\mathbb{N}| < |A| \iff A$ is neither finite nor countable, and

$$3. |A| \leq |\mathbb{N}| \wedge |\mathbb{N}| \leq |A| \implies |A| = |\mathbb{N}|.$$

Definition B.2.4 (Countability and \aleph_0)

Let A be a set. When A is countable we write $|A| = \aleph_0$.

When A is finite we write $|A| < \aleph_0$.

When A is infinite we write $|A| \geq \aleph_0$.

When A is either finite or countable we write $|A| \leq \aleph_0$, and say that A is at most countable.

When A is neither finite nor countable we write $|A| > \aleph_0$, and say that A is uncountable.

Theorem B.2.4 (Set Cartesian Product and Union, and \mathbb{Q} are Countable)

1. If A and B are countable sets, then so is $A \times B$.
2. If A and B are countable sets, then so is $A \cup B$.
3. If A_0, A_1, A_2, \dots are countable sets, then so is $\bigcup_{k=0}^{\infty} A_k$.
4. \mathbb{Q} is countable.

Remark

For a set A , we let 2^A denote the set of all functions from A to $S_2 = \{0, 1\}$, i.e.

$$2^A = \{f | f : A \rightarrow S_2\}$$

Theorem B.2.5 (\mathbb{R} as an Uncountable Set)

1. For every set A , $|\mathcal{P}(A)| = |2^A|$.
2. For every set A , $|A| < |\mathcal{P}(A)|$.
3. \mathbb{R} is uncountable.

Theorem B.2.6 (Cantor-Schröder-Bernstein Theorem)

Let A and B be sets.

$$|A| \leq |B| \wedge |B| \leq |A| \implies |A| = |B|.$$