# PMATH451 — Measure and Integration

Class notes for Fall 2019

by

Johnson Ng

BMath (Hons), Pure Mathematics major, Actuarial Science Minor

University of Waterloo

# Table of Contents

Τa	Table of Contents	2
Li	ist of Definitions	4
Li	ist of Theorems	5
Li	ist of Procedures	6
Li	ist of Homework	7
Pı	Preface	9
1	Lecture 1 Sep 04th, 2019	11
	1.1 Motivation for the Study of Measures	11
	1.2 Algebras and $\sigma$ -Algebra of Sets	15
2	Lecture 2 Sep 06th 2019	19
	2.1 Algebra and $\sigma$ -algebra of Sets (Continued)	19
	2.2 Measures	25
3	Lecture 3 Sep 9th 2019	27
	3.1 Measures (Continued)	27
4	Lecture 4 Sep 11th 2019	33
	4.1 Measures (Continued 2)	33
5	Lecture 5 Sep 13th 2019	39
	5.1 Measures (Continued 3)	39
	5.2 The Outer Measure	41

6	Lecture 6 Sep 16th 2019	45
	6.1 The Outer Measure (Continued)	45
	6.2 The Lebesgue-Stieltjes Outer Measure	49
7	Lecture 7 Sep 18th 2019	51
	7.1 The Lebesgue-Stieltjes Outer Measure (Continued)	51
A	Deep Dives into Proofs	57
	A.1 Proving that $\mathfrak M$ is closed under countable unions in Carathéodory's Theorem $\ \ldots \ \ldots \ \ldots$	57
В	Common Themes and Tricks	61
	B.1 Re-represent an arbitrary union using disjoint sets	61
Bi	ibliography	63
In	ndex	64

# List of Definitions

1	■ Definition (Algebra of Sets)	15
2	$\blacksquare$ Definition ( $\sigma$ -Algebra of Sets)	10
3	$\blacksquare$ Definition (Generator of a $\sigma$ -algebra)	20
4	$\blacksquare$ Definition (Borel $\sigma$ -algebra)	2
5	■ Definition (Infinitely Often)	24
6	E Definition (Almost always)	24
7	E Definition (Characteristic Function)	25
8	■ Definition (Measure)	25
9	■ Definition (Measure Space)	27
10	$\blacksquare$ Definition (Finitivity, $\sigma$ -finitivity, and Semi-finitivity of a Measure)	29
11	■ Definition (Null Set of a Measure)	36
12	■ Definition (Complete Measure Space)	36
13	■ Definition (Outer Measure)	4
14	$\blacksquare$ Definition ( $\mu^*$ -measurability)	4
15	■ Definition (Metric Outer Measure)	46
16	■ Definition (Lebesgue-Stieltjes Outer Measure)	50

# List of Theorems

1	♦ Proposition (Properties of Algebra of Sets)	15
2	<b><math>\bullet</math></b> Proposition (Closure of $\sigma$ -algebras under Countable Intersection)	19
3	<b><math>\delta</math></b> Proposition (Existence of the 'Smallest' $\sigma$ -algebra on a Set)	20
4	$\bullet$ Proposition (Other Formulations of the Borel $\sigma$ -algebra (aka Proposition 1.2))	22
5	■ Theorem (Properties of a Measure)	30
6	■ Theorem (Extending the Measurable Sets)	37
7	♦ Proposition (Lebesgue's Outer Measure)	41
8	■ Theorem (Carathéodory's Theorem)	44
9	■Theorem (Carathéodory's Second Theorem)	51
10	♦ Proposition (Lebesgue-Stieltjes Outer Measure on Half-open Intervals)	54

# List of Procedures

# List of Homework

🌣 Homework (Homework 1)	24
♣ Homework (Homework 2 – A review on limsup and liminf)	25
<b>♦</b> Homework (Homework 3)	31
♣ Homework (Homework 4)	45
🍫 Homework (Homework 5)	45
to Homework (Homework 6)	46



Assignment problems is introduced in class as we go, so we have the special environment homework for these in this note.

I included a special chapter in the appendix (see Appendix A) that records and provides insights into what drove the direction(s) of certain proofs. This is an attempt to resolve the problem of proofs being overly obscure with its motivations. Contents presented in this appendix are typically like rough work, and so are typically much longer than the presented proof.

I also made an appendix for some of the common themes and tricks (see Appendix B) that are seen repeatedly in this topic I think it is invaluable that they are noted down, because the ideas that these commonalities carry forward.

# 1.1 Motivation for the Study of Measures

Recall Riemann integration.

# **■** Definition (Riemann Integration)

Let  $f:[a,b] \to \mathbb{R}$  be a **bounded** function. We call

$$P = \{a = x_0 < x_1 < \dots < x_n = b\} \subseteq [a, b]$$

a partition of [a, b], and

$$\Delta x_i = x_i - x_{i-1}$$

as the length of the  $i^{th}$  interval for i = 1, ..., n.

Let

$$M_i = \sup\{f(x) : x \in [x_{i-1}, x_i]\}$$

be the supremum of f on the i<sup>th</sup> interval, and

$$m_i = \inf\{f(x) : x \in [x_{i-1}, x_i]\}$$

be the infimum of f on the  $i^{th}$  interval. We define the Riemann upper sum as

$$U(f,P) = \sum_{i} M_i \Delta x_i,$$

and the Riemann lower sum as

$$L(f,P) = \sum_{i} m_i \Delta x_i.$$




Figure 1.1: Idea of Riemann integration

We define the Riemann upper integral as

$$\int_{a}^{b} f \, dx = \inf_{P} U(f, P)$$

and the Riemann lower integral as

$$\int_{a_{-}}^{b} f \, dx = \sup L(f, P).$$

We say that f is Riemann integrable if

$$\overline{\int_a^b} f \, dx = \int_a^b f \, dx,$$

and we write the integral of f as

$$\int_{a}^{b} f \, dx = \overline{\int_{a}^{b}} f \, dx = \int_{a}^{b} f \, dx.$$

As hyped up as one does earlier in university about Riemann integration, there are functions that are not Riemann integrable!

#### Example 1.1.1

Consider a function  $f : [0,1] \to \mathbb{R}$  given by

$$f(x) = \begin{cases} 1 & x \in \mathbb{Q} \\ 0 & x \notin \mathbb{Q} \end{cases}.$$

Then

$$\overline{\int_a^b} f \, dx = 1 \text{ and } \underline{\int_a^b} f \, dx = 0.$$

Thus *f* is not Riemann integrable.

### **66** Note 1.1.1 (Shortcomings of the Riemann integral)

1. We cannot characterize functions that are Riemann integrable, i.e. we do not have a list of characteristics that we can check against to see if a function is Riemann integrable.

*This remained an open problem in the earlier 1920s.* 

- 2. The Riemann integral behaves badly when it comes to pointwise limits of functions. The next example shall illustrate this.
- 3. The Riemann integral is awkward when f is unbounded. In particular, we used to hack our way around by looking at whether the Riemann integral converges to some value the function approaches the unbounded point, and then "conclude" that the integral is the limit of that convergence.
- 4. Recall that the Fundamental Theorem of Calculus states that

$$\frac{d}{dx} \int_{a}^{x} f(t) \, dt = f(x).$$

We know that this works for Riemann integrals. By the first shortcoming, the problem here is that we do not fully know what are the functions that the Fundamental Theorem is true for.

- 5. In PMATH450, we saw that Fourier developed the Fourier series, which is an extremely useful tool in solving Differential Equations using sines and cosines. However, the convergence of the Fourier series remains largely unexplained by Fourier, and we have but developed some roundabout ways of showing some convergence.
- 6. Consider the set R if Riemann integrable functions on the interval [a, b]. The set R has a natural metric:

$$d(f,g) = \int_{a}^{b} |f - g| \, dx.$$

However, the metric space (R, d) is **not complete**. This means many of our favorite results in PMATH351 are not usable!

7. There are many functions that seem like they should have an integral, but turned out that they did not under Riemann integration.

Example 1.1.2 (Pointwise Limits of Riemann Integrable Functions is not necessarily Riemann Integrable)

Let  $\mathbb{Q} = \{x_n\}_{n \in \mathbb{N}}$ . Then consider a sequence of functions

$$f_n(x) = \begin{cases} 1 & x \in \{x_1, \dots, x_n\} \\ 0 & x \notin \{x_1, \dots, x_n\} \end{cases}.$$

It is rather clear that

$$\overline{\int_a^b} f \, dx = \int_a^b f \, dx = 0.$$

However, the pointwise limit of the  $f_n$ 's, and that is

$$\lim_{n \to \infty} f_n(x) = f(x) = \begin{cases} 1 & x \in \mathbb{Q} \\ 0 & x \notin \mathbb{Q} \end{cases},$$

is, as mentioned in the last example, not Riemann integrable.

\*

To address the shortcomings of the Riemann integral, Henri Lebesgue developed the Lebesgue integral, of which we have seen in PMATH450.

Instead of dividing the *x*-axis, Lebesgue decided to divide the *y*-axis first.

If the range of a function f is [c,d], where c,d can be infinite, then we partition the interval such that

$$P = \{c = y_0 < y_1 < \ldots < y_n = d\},\$$

and we define

$$E_i = \{x : f(x) \in [y_{i-1}, y_i]\}.$$

Then if  $A_i$  is the area of the "rectangle" for the  $i^{th}$  interval of [c,d], we have

$$y_{i-1} \cdot \ell(E_i) \le A_i \le y_i \cdot \ell(E_i),$$

where  $\ell(E_i)$  is the Lebesgue measure of the set  $E_i$ . Then if we let  $\int_a^b f$  denote the Lebesgue integral of f, we would expect

$$\sum_{i=1}^n y_{i-1} \cdot \ell(E_i) \le \int_a^b f \le \sum_{i=1}^n y_i \cdot \ell(E_i).$$

However, to truly understand what this means, we need to understand what the Lebesgue measure is.

Furthermore, recall that in PMATH450, we saw that not all sets, in  $\mathbb{R}$  for example, are measurable, and for 'good' reasons, there always exists non-measurable sets.

# Algebras and $\sigma$ -Algebra of Sets

# **Definition 1 (Algebra of Sets)**

Given X, a non-empty collection of subsets of X, i.e.  $\emptyset \neq \mathcal{A} \subseteq \mathcal{P}(X)$ , is called an algebra of sets of X provided that:

1. 
$$A_1, \ldots, A_n \in \mathcal{A} \implies \bigcup_{i=1}^n A_i \in \mathcal{A}$$
; and

2. 
$$A \in \mathcal{A} \implies A^C \in \mathcal{A}$$
.

### **♦** Proposition 1 (Properties of Algebra of Sets)

If  $\mathcal{A}$  is an algebra of sets of X, then

3.  $\emptyset$ ,  $X \in \mathcal{A}$ ;

4. 
$$A, B \in \mathcal{A} \implies A \setminus B = \{x \in X \mid x \in A \land x \notin B\} \in \mathcal{A} ; and$$

5. 
$$A_1, \ldots, A_n \in \mathcal{A} \implies \bigcap_{i=1}^n A_i \in \mathcal{A}$$
.

#### Proof

3. 
$$\mathcal{A} \neq \emptyset \implies \exists A \in \mathcal{A} \implies A^C \in \mathcal{A} \implies A \cup A^C = X \in \mathcal{A} \implies \emptyset = X^C \in \mathcal{A}.$$

4. 
$$A, B \in \mathcal{A} \implies A^C \in \mathcal{A} \implies A^C \cup B \in \mathcal{A} \implies A \setminus B = (A^C \cup B)^C \in \mathcal{A}$$
.

5. (**De Morgan's Law**) Notice that  $(A_1 \cap A_2 \cap ... \cap A_n)^C = A_1^C \cup A_2^C \cup A_2^C$  $\dots A_n^C \in \mathcal{A}$  since  $A_i^C \in \mathcal{A}$ . Thus the complement

$$A_1 \cap A_2 \cap \ldots \cap A_n \in \mathcal{A}$$
.

For this course, we shall use the convention that

- the 'ambient' space *X* is always non-empty;
- $\mathcal{P}(X)$ , the power set of X, has nontrivial elements; and
- we denote  $A^C = \{x \in X : x \notin A\}$  for  $A \subseteq X$ .

# **E** Definition 2 ( $\sigma$ -Algebra of Sets)

Given X and  $\emptyset \neq \mathcal{A} \subseteq \mathcal{P}(X)$ , we say that  $\mathcal{A}$  is a  $\sigma$ -algebra of sets of X if it is an algebra of sets and

$$\forall A_n \in \mathcal{A}, n \in \mathbb{N}, \quad \bigcup_{n \in \mathbb{N}} A_n \in \mathcal{A}.$$

# Example 1.2.1

- 1.  $\mathcal{P}(X)$  is a *σ*-algebra.
- 2. Consider X as an infinite set. We say that a set A is **cofinite** if  $A^C$  is finite. Let

$$\mathcal{A} := \{ A \in \mathcal{P}(X) \mid A \text{ is finite or cofinite } \}.$$

Then  $\mathcal{A}$  is an algebra of sets:

- finite union of finite sets remains finite;
- finite union of finite and cofinite sets remains cofinite; and
- complement of finite sets are the cofinite sets and vice versa.

However,  $\mathcal{A}$  is **not** a  $\sigma$ -algebra: consider  $A_n = \{2^n\} \subseteq X = \mathbb{N}$ , which we then realize that

$$\bigcup_{n\in\mathbb{N}} A_n = \text{ set of all even numbers },$$

but the set of all even numbers is clearly not finite, and its complement, which is the set of all odd numbers, is not finite.

3. Consider X as an uncountable set. We say that a set A is co-countable if  $A^C$  is countable. <sup>1</sup> The set

$$\mathcal{A} := \{ A \subseteq X \mid A \text{ is countable or co-countable } \}$$

is a  $\sigma$ -algebra:

• countable union of countable sets is countable;

 $<sup>^{1}</sup>$  Recall that a set A is said to be countable if there is a one-to-one correspondence between elements of A and the natural numbers.

- countable union of countable and co-countable sets is co-countable; and
- complement of countable sets are co-countable and vice versa.



# 2.1 Algebra and $\sigma$ -algebra of Sets (Continued)

We've seen some examples of  $\sigma$ -algebras. Let's now look at some other important properties of  $\sigma$ -algebras.

# **\bullet** Proposition 2 (Closure of $\sigma$ -algebras under Countable Intersection)

Let X be a set,  $\mathcal{A}$  a  $\sigma$ -algebra on X. If  $A_n \in \mathcal{A}$  for each  $n \in \mathbb{N}$ , then  $\bigcap_n A_n \in \mathcal{A}$ .

This follows rather similarly to **\langle** Proposition 1 where we used **De Morgan's Law**.

#### Proof

We observe that

$$A_n \in \mathcal{A} \implies A_n^C \in \mathcal{A}$$

$$\implies \bigcup_n A_n^C \in \mathcal{A}$$

$$\implies \bigcap_n A_n = \left(\bigcup_n A_n^C\right)^C \in \mathcal{A}.$$

Let  $\mathcal{A}_{\alpha} \subseteq \mathcal{P}(X)$ , where  $\alpha$  is from some index set. We denote

$$\bigcap_{\alpha} \mathcal{A}_{\alpha} = \{ A \subseteq X : A \in \mathcal{A}_{\alpha}, \, \forall \alpha \}.$$

# **♦** Proposition 3 (Existence of the 'Smallest' *σ*-algebra on a Set)

Let X be a set and  $\{\mathcal{A}_{\alpha}\}_{\alpha}$  as a collection of  $\sigma$ -algebras on X. Then  $\bigcap_{\alpha} A_{\alpha}$  is a  $\sigma$ -algebra.

### Proof

$$A \in \bigcap_{\alpha} \mathcal{A}_{\alpha} \implies \forall \alpha, A \in \mathcal{A}_{\alpha}$$
$$\implies \forall \alpha, A^{C} \in \mathcal{A}_{\alpha}$$
$$\implies A^{C} \in \bigcap_{\alpha} \mathcal{A}_{\alpha}$$

and

$$\forall n \in \mathbb{N}, A_n \in \bigcap_{\alpha} \mathcal{A}_{\alpha} \implies \forall n \in \mathbb{N}, \forall \alpha, A_n \in \mathcal{A}_{\alpha}$$

$$\implies \forall \alpha, \bigcup_{n} A_n \in \mathcal{A}_{\alpha}$$

$$\implies \bigcup_{n} A_n \in \bigcap_{\alpha} \mathcal{A}_{\alpha}.$$

Due to the above proposition, the following definition is well-defined.

#### **E** Definition 3 (Generator of a $\sigma$ -algebra)

Let X be a set, and  $\xi \subseteq \mathcal{P}(X)$  has some non-trivial set(s). Consider all  $\sigma$ -algebras  $\mathcal{A}_{\alpha}$  with the property that  $\xi \subseteq \mathcal{A}_{\alpha}$ . Then we say that  $\bigcap_{\alpha} \mathcal{A}_{\alpha}$ 

is the  $\sigma$ -algebra generated by  $\xi$ , and we denote this generated  $\sigma$ -algebra as

$$\mathfrak{M}(\xi) = \bigcap_{\alpha} \mathcal{A}_{\alpha}.$$

#### Remark 2.1.1

- 1. It is clear from the definition that if  $\mathcal{A}$  is a  $\sigma$ -algebra on X and  $\xi \subseteq \mathcal{A}$ , then  $\mathfrak{M}(\xi) \subseteq \mathcal{A}$ .
- 2. We often say that  $\mathfrak{M}(\xi)$  is the "smallest  $\sigma$ -algebra containing  $\xi$ ".

The following is an example of such a  $\sigma$ -algebra.

# **E** Definition 4 (Borel $\sigma$ -algebra)

Let X be a metric space (or topological space). The  $\sigma$ -algebra generated by the open subsets of X is called the Borel  $\sigma$ -algebra, of which we denote by  $\mathfrak{B}(X)$ .

#### Remark 2.1.2 (Some sets in $\mathfrak{B}(X)$ )

Given an arbitrary metric space (or topological space) X. It is often hard to firmly grasp what kind of sets are in the Borel  $\sigma$ -algebra  $\mathfrak{B}(X)$ . The following are some examples that are in  $\mathfrak{B}(X)$ .

- 1. Let  $\{O_n\}_{n\in\mathbb{N}}$  denote a countable collection of open sets. By  $\lozenge$  Proposi*tion* 2,  $\bigcap_n O_n \in \mathfrak{B}(X)$ . We call these countable union of open sets as  $G_\delta$ sets.
- 2. Let  $\{\mathcal{F}_n\}_{n\in\mathbb{N}}$  denote a countable collection of closed sets. By  $\Diamond$  Proposition 2,  $\bigcup_n \mathcal{F}_n \in \mathfrak{B}(X)$ . We call these countable intersection of closed sets as  $F_{\sigma}$  sets.
- 3. Let  $\{H_n\}$  be a countable collection of  $G_\delta$  sets. Then  $\bigcup_n H_n \in \mathfrak{B}(X)$ . *These are called the*  $G_{\delta\sigma}$  *sets.*
- 4. Let  $\{K_n\}$  be a countable collection of  $F_{\sigma}$  sets. Then  $\bigcap_n K_n \in \mathfrak{B}(X)$ . These are called the  $F_{\sigma\delta}$  sets.

We can continue constructing the  $G_{\delta\sigma...}$  and  $F_{\sigma\delta...}$  similarly, and all these sets belong to the Borel  $\sigma$ -algebra  $\mathfrak{B}(X)$ .

# • Proposition 4 (Other Formulations of the Borel σ-algebra (aka Proposition 1.2))

The following collection of sets are all equal:

- 1.  $\mathfrak{B}_1 = \mathfrak{B}(\mathbb{R})$ ;
- 2.  $\mathfrak{B}_2 = \sigma$ -algebra generated by open intervals (e.g. (a, b));
- 3.  $\mathfrak{B}_3 = \sigma$ -algebra generated by closed intervals (e.g. [a, b]);
- 4.  $\mathfrak{B}_4 = \sigma$ -algebra generated by half-open intervals (e.g. (a, b]);
- 5.  $\mathfrak{B}_5 = \sigma$ -algebra generated by  $(-\infty, a)$  and  $(b, \infty)$ ; and
- 6.  $\mathfrak{B}_6 = \sigma$ -algebra generated by  $(-\infty, a]$  and  $[b, \infty)$ .

As commented before, it is often hard knowing that is in a Borel  $\sigma$ -algebra, and what is not, despite knowing what its generator is. However, when talking about containments, this is a fairly straightforward discussion thanks to its closure under countable unions and Proposition 2. We simply need to talk about the generators.

#### Proof

 $\mathfrak{B}_2 \subseteq \mathfrak{B}_1$  Given an arbitrary generator (a,b) in  $\mathfrak{B}_2$ , we know that (a,b) is an open set, and clearly  $(a,b) \subseteq \mathbb{R}$ . Thus  $(a,b) \in \mathfrak{B}_1$ , so  $\mathfrak{B}_2 \subseteq \mathfrak{B}_1$ .

 $\mathfrak{B}_3 \subseteq \mathfrak{B}_2$  Given an arbitrary generator [a, b] of  $\mathfrak{B}_2$ , we have

$$[a,b] = \bigcap_{n} \left(a - \frac{1}{n}, b + \frac{1}{n}\right) \in \mathfrak{B}_2.$$

Thus  $\mathfrak{B}_3 \subseteq \mathfrak{B}_2$ .

 $\mathfrak{B}_4 \subseteq \mathfrak{B}_3$  Given an arbitrary generator (a, b] of  $\mathfrak{B}_4$ ,

$$(a,b] = \bigcup_{n} \left[ a + \frac{1}{n}, b \right] \in \mathfrak{B}_3.$$

Thus  $\mathfrak{B}_4 \subseteq \mathfrak{B}_3$ .

 $\mathfrak{B}_5 \subseteq \mathfrak{B}_4$  Given an arbitrary generator  $(-\infty, a)$  for  $\mathfrak{B}_5$ ,

$$(-\infty, a) = \bigcup_{n} \left(-\infty, a - \frac{1}{n}\right) \in \mathfrak{B}_4.$$

On the other hand, for  $(b, \infty)$  in  $\mathfrak{B}_5$ ,

$$(b,\infty) = \bigcup_{n} (b,n) \in \mathfrak{B}_4.$$

 $\mathfrak{B}_6 \subseteq \mathfrak{B}_5$  We have that

$$(-\infty, a] = \bigcap_{n} \left(-\infty, a + \frac{1}{n}\right) \in \mathfrak{B}_{5}$$

and

$$[b,\infty)=\bigcap_n\left(b-\frac{1}{n},\infty\right)\in\mathfrak{B}_5.$$

 $\mathfrak{B}_1 \subseteq \mathfrak{B}_6$  Let  $c < d \in \mathbb{R}$ . Notice that

$$(-\infty, d] \cap [c, \infty) = [c, d] \in \mathfrak{B}_6.$$

Furthermore,

$$(c,d) = \bigcup_{n} \left[ c + \frac{1}{n}, d - \frac{1}{n} \right] \in \mathfrak{B}_6.$$

Recall that given an open set  $O \subseteq \mathbb{R}$ , we have

$$O = \bigcup \{ (c,d) \subseteq O : c,d \in \mathbb{Q} \},\$$

which shows that O is a countable union of open sets (with rational endpoints). It follows that  $O \in \mathfrak{B}_6$  and so  $\mathfrak{B}_1 \subseteq \mathfrak{B}_6$ . 

#### Exercise 2.1.1

Show that  $\mathfrak{B}(\mathbb{R}^2)$  is generated by open rectangles  $(a,b) \times (c,d)$ .

### **■** Definition 5 (Infinitely Often)

Given  $E_n \subseteq X$  for  $n \in \mathbb{N}$ , we say that  $x \in E_n$  infinitely often (i.o.) if

$${n:x\in E_n}$$

is an **infinite set**. We typically let

$$A := \{x \in X : x \in E_n \ i.o. \ \}$$

be the set of x's that are in the  $E_n$ 's infinitely often.

# **■** Definition 6 (Almost always)

Given  $E_n \subseteq X$  for  $n \in \mathbb{N}$ , we say that  $x \in E_n$  almost always (a.a.) if

$${n:x \notin E_n}$$

is a *finite set*. We typically let

$$B := \{x \in X : x \in E_n \ a.a. \}$$

be the set of x's that are in the  $E_n$ 's almost always.

#### **\$\phi\_0** Homework (Homework 1)

Let X be a set,  $\mathcal{A}$  a  $\sigma$ -algebra on X, and  $E_n \in \mathcal{A}$  for  $n \in \mathbb{N}$ . Prove that

$$A := \{x \in X : x \in E_n \ i.o. \ \}$$

and

$$B := \{x \in X : x \in E_n \ a.a. \}$$

are both in  $\mathcal{A}$ .

#### **■** Definition 7 (Characteristic Function)

*Let*  $E \subseteq X$ *. We call the function* 

$$\chi_E(x) = \begin{cases} 1 & x \in E \\ 0 & x \notin E \end{cases}$$

the characteristic function of E.

# **‡** Homework (Homework 2 − A review on limsup and liminf)

Let  $E_n \subseteq X$  for  $n \in \mathbb{N}$ , and

$$A := \{x \in X : x \in E_n \ i.o. \}$$
  
$$B := \{x \in X : x \in E_n \ a.a. \}.$$

Show that

$$\chi_A(x) = \limsup_n \chi_{E_n}(x)$$

$$\chi_B(x) = \liminf_n \chi_{E_n}(x).$$

#### Remark 2.1.3

Due to the above result, some people write

$$A = \limsup E_n$$
$$B = \liminf E_n.$$

# Measures

# **Definition 8 (Measure)**

Let X be a set and  $\mathcal{A}$  a  $\sigma$ -algebra of subsets of X. A function  $\mu:\mathcal{A}\to$  $[0, \infty]$  is called a *measure* on  $\mathcal{A}$  provided that:

1. 
$$\mu(\emptyset) = 0$$
; and

2. if  $E_n \in \mathcal{A}$  for each  $n \in \mathbb{N}$ , and  $\{E_n\}$  is disjoint, we have

$$\mu\left(\bigcup_n E_n\right) = \sum_n \mu(E_n).$$



# 3.1 Measures (Continued)

# **E** Definition 9 (Measure Space)

Let X be a set,  $\mathfrak{M}$  a  $\sigma$ -algebra of subsets of X and  $\mu: \mathfrak{M} \to [0, \infty]$ . We call the 3-tuple  $(X, \mathfrak{M}, \mu)$  a measure space.

#### Remark 3.1.1

If  $\mu(X) = 1$ , we also call  $(X, \mathfrak{M}, \mu)$  a probability space, and  $\mu$  is called a probability measure.

#### Example 3.1.1

1. (Counting Measure) Let X be a set and  $\mathfrak{M} = \mathcal{P}(X)$ . For  $E \in \mathfrak{M}$ , define

$$\mu(E) = \begin{cases} |E| & E \text{ is finite} \\ \infty & \text{otherwise} \end{cases}.$$

We verify that  $\mu$  is indeed a measure:

- (a) We have that  $\mu(\emptyset) = |\emptyset| = 0$ .
- (b) Let  $\{E_n\}_{n=1}^{\infty} \subseteq \mathfrak{M}$  be a pairwise disjoint set. Notice that if any of the sets are infinite, say  $E_{N_0}$  is infinite, then

$$\mu(E_{N_0}) = \infty = |E_{N_0}|.$$

Since  $\bigcup_{n=1}^{\infty} E_n$  is infinite in this case, we have

$$\mu\left(\bigcup_{n=1}^{\infty} E_n\right) = \infty = \left|\bigcup_{n=1}^{\infty} E_n\right|.$$

On the other hand, if all the sets are finite, then since the  $E_n$ 's are disjoint, we have

$$\mu\left(\bigcup_{n=1}^{\infty} E_n\right) = \left|\bigcup_{n=1}^{\infty} E_n\right| = \sum_{n=1}^{\infty} |E_n| = \sum_{n=1}^{\infty} \mu(E_n).$$

We call  $\mu$  a counting measure.

2. Let *X* be an uncountable set. Recall that in Example 1.2.1, we showed that

$$\mathfrak{M} := \{ A \subseteq X \mid A \text{ is countable or co-countable } \}$$

is a  $\sigma$ -algebra. There are many measures that we can define on this  $\sigma$ -algebra. For instance,

$$\nu(E) = \begin{cases} 0 & E \text{ is countable} \\ 1 & E \text{ is uncountable} \end{cases},$$

and

$$\delta(E) = \begin{cases} 0 & E \text{ is countable} \\ \infty & E \text{ is uncountable} \end{cases}.$$

Verifying that both  $\nu$  and  $\delta$  are indeed measures shall be left to the reader as a straightforward exercise.

3. Let's make a non-example. Let X be an infinite set, and  $\mathfrak{M} = \mathcal{P}(X)$ . Define

$$\mu(E) = \begin{cases} 0 & E \text{ is finite} \\ \infty & E \text{ is infinite} \end{cases}.$$

Consider  $X = \mathbb{N}$  and a sequence of sets with singletons,

$$E_n = \{2n+1\}, \quad \text{for } n \in \mathbb{N}.$$

Clearly,

$$\bigcup_{n=1}^{\infty} E_n = \text{ set of all odd numbers },$$

and clearly

$$\mu\left(\bigcup_{n=1}^{\infty}E_n\right)=\infty.$$

However, notice that

$$\mu(E_n) = 0$$
 for each  $n \in \mathbb{N}$ .

Since each of the  $E_n$ 's are pairwise disjoint, we should have

$$\infty = \mu\left(\bigcup_{n=1}^{\infty} E_n\right) = \sum_{n=1}^{\infty} \mu(E_n) = 0,$$

which is impossible. Thus  $\mu$  is **not** a measure.



### Remark 3.1.2 (Finite additivity)

Given a finite set of pairwise disjoint sets  $\{E_n\}_{n=1}^N\subseteq\mathfrak{M}$  for some  $\sigma$ -algebra  $\mathfrak{M}$  of some set X. By the definition of a  $\sigma$ -algebra, we may set  $E_n = \emptyset$  for n > N. Then

$$\mu\left(\bigcup_{n=1}^{N} E_n\right) = \mu\left(\bigcup_{n=1}^{\infty} E_n\right) = \sum_{n=1}^{\infty} \mu(E_n) = \sum_{n=1}^{N} \mu(E_n).$$

We call this the finite additivity of a measure.



# **Ξ** Definition 10 (Finitivity, $\sigma$ -finitivity, and Semi-finitivity of a Measure)

Let  $(X, \mathfrak{M}, \mu)$  be a measure space.

- 1. We say that  $\mu$  is **finite** if  $\mu(E) < \infty$  for every  $E \in \mathfrak{M}$ .
- 2. If  $X = \bigcup_{n=1}^{\infty} X_n$  with  $X_n \in \mathfrak{M}$ , we say that  $\mu$  is  $\sigma$ -finite if

$$\mu(X_n) < \infty$$
 for every  $n \in \mathbb{N}$ .

3. We say that  $\mu$  is semi-finite if for every  $E \in \mathfrak{M}$  with  $\mu(E) = \infty$ ,

 $\exists F \subseteq E \in \mathfrak{M} \text{ such that }$ 

$$0 < \mu(F) < \infty$$
.

#### Exercise 3.1.1

- 1. Show that the counting measure is finite iff the ambient space X is a finite set.
- 2. Show that  $\delta$  in Example 3.1.1 is neither finite,  $\sigma$ -finite, nor semi-finite.

#### **■**Theorem 5 (Properties of a Measure)

Let  $(X, \mathfrak{M}, \mu)$  be a measure space. Then

- 1. (Monotonicity) If  $E \subseteq F$  and  $E, F \in \mathfrak{M}$ , then  $\mu(E) \leq \mu(F)$ .
- 2. (Subadditivity) If  $\{E_n\}_{n=1}^{\infty} \subseteq \mathfrak{M}$ , then

$$\mu\left(\bigcup_n E_n\right) \leq \sum_n \mu(E_n).$$

3. (Continuity from below) If  $\{E_n\}_{n=1}^{\infty} \subseteq \mathfrak{M}$  is an increasing sequence of sets, i.e.

$$E_1 \subseteq E_2 \subseteq \ldots \subseteq E_n \subseteq \ldots$$

then

$$\mu\left(\bigcup_{n=1}^{\infty} E_n\right) = \lim_{n \to \infty} \mu(E_n).$$

4. (Continuity from above) If  $\{E_n\}_{n=1}^{\infty} \subseteq \mathfrak{M}$  is a decreasing sequence of sets, i.e.

$$E_1 \supseteq E_2 \supseteq \ldots \supseteq E_n \supseteq \ldots$$

and  $\exists n_0 \in \mathbb{N}$  such that  $\mu(E_{n_0}) < \infty$ , then

$$\mu\left(\bigcap_{n=1}^{\infty}E_n\right)=\lim_{n\to\infty}\mu(E_n).$$

# Remark 3.1.3 (A comment on the condition for the 4<sup>th</sup> statement)

It may seem that the extra condition of a finite measure seem extravagant. However, it is necessary, as demonstrated below.

Consider  $X = \mathbb{N}$ , with  $\mu$  as the counting measure. Then, consider the sequence of sets

$$E_{1} = \{1, 2, 3, \ldots\},\$$

$$E_{2} = \{2, 3, 4, \ldots\},\$$

$$E_{3} = \{3, 4, 5, \ldots\},\$$

$$\vdots$$

$$E_{n} = \{n, n+1, n+2, \ldots\},\$$

$$\vdots$$

Then  $\bigcap_{n=1}^{\infty} E_n = \emptyset$ , which then  $\mu\left(\bigcap_{n=1}^{\infty} E_n\right) = 0$ . However,

$$\mu(E_n) = \infty$$
 for each  $n \in \mathbb{N}$ .

### **♦** Homework (Homework 3)

Let  $(X, \mathfrak{M}, \mu)$  be a measure space. Let  $\{E_n\}_{n=1}^{\infty} \subseteq \mathfrak{M}$ , and

$$A := \{x \in X \mid x \in E_n \ i.o. \ \}.$$

*Prove that*  $\sum_{n=1}^{\infty} \mu(E_n) < \infty$  *implies that*  $\mu(A) = 0$ .

# *∠ Lecture 4 Sep 11th 2019*

# 4.1 Measures (Continued 2)

We shall now prove **P**Theorem 5.



1. Notice that

$$F = (F \cap E) \cup (F \setminus E),$$

and  $F \cap E$  and  $F \setminus E$  are disjoint. Thus

$$\mu(F) = \mu(F \cap E) + \mu(F \setminus E) = \mu(E) + \mu(F \setminus E).$$

Since  $\mu(F \setminus E) \ge 0$ , we have

$$\mu(F) \ge \mu(E)$$
.

2. Consider a sequence of sets defined as such: 1

$$F_{1} = E_{1}$$

$$F_{2} = E_{2} \setminus E_{1}$$

$$\vdots$$

$$F_{n} = E_{n} \setminus \bigcup_{j=1}^{n-1} E_{j}.$$

First, note that  $F_n \subseteq E_n$  for each  $n \in \mathbb{N}$ . So by the last part, we have

$$\mu(F_n) \le \mu(E_n)$$
 for each  $n \in \mathbb{N}$ .

¹ ★ This is a common technique in measure theory. We will see this repeatedly so in this course.

Secondly,

$$\bigcup_{n=1}^{\infty} F_n = \bigcup_{n=1}^{\infty} E_n.$$

Also,  $\{F_n\}_{n=1}^{\infty}$  is a pairwise disjoint collection of sets. It follows that

$$\mu\left(\bigcup_{n=1}^{\infty} E_n\right) = \mu\left(\bigcup_{n=1}^{\infty} F_n\right) = \sum_{n=1}^{\infty} \mu(F_n) \le \sum_{n=1}^{\infty} \mu(E_n).$$

3. Consider a sequence of sets defined as such:

$$F_1 = E_1$$

$$F_2 = E_2 \setminus E_1$$

$$F_3 = E_3 \setminus E_2$$

$$\vdots$$

$$F_n = E_n \setminus E_{n-1}.$$

We see that

• 
$$\bigcup_{n=1}^{\infty} F_n = \bigcup_{n=1}^{\infty} E_n$$
;

• 
$$\bigcup_{n=1}^{N} F_n = \bigcup_{n=1}^{N} E_n = E_N$$
; and

•  $\{F_n\}_n$  is a collection pairwise disjoint sets.

Thus we have

$$\mu\left(\bigcup_{n=1}^{\infty} E_n\right) = \mu\left(\bigcup_{n=1}^{\infty} F_n\right) = \sum_{n=1}^{\infty} \mu(F_n)$$
$$= \lim_{N \to \infty} \sum_{n=1}^{N} \mu(F_n) = \lim_{N \to \infty} \mu\left(\bigcup_{n=1}^{N} F_n\right)$$
$$= \lim_{N \to \infty} \mu(E_N).$$

4. First, it is important that we notice that

$$\bigcap_{n=1}^{\infty} E_n = \bigcap_{n=m}^{\infty} E_n$$

for any  $m \in \mathbb{N}$ , since  $\{E_n\}_n$  is a decreasing sequence of sets.

Suppose  $n_0 \in \mathbb{N}$  is such that  $\mu(E_{n_0}) < \infty$ . Consider a sequence of sets defined as follows: for  $n_0 \le j \in \mathbb{N}$ , we let  $F_j = E_{n_0} \setminus E_j$ . Then

we have

$$\emptyset = F_{n_0} \subseteq F_{n_0+1} \subseteq \ldots \subseteq F_{n_0+k} \subseteq \ldots,$$

i.e.  $\{F_n\}_{n=n_0}^{\infty}$  is an increasing sequence of sets. By the last part, we have

$$\mu\left(\bigcup_{n=n_0}^{\infty} F_n\right) = \lim_{n \to \infty} \mu(F_{n_0+n}) = \lim_{n \to \infty} \mu(E_{n_0} \setminus E_{n_0+n})$$
$$= \mu(E_{n_0}) - \lim_{n \to \infty} \mu(E_{n_0+n})$$
$$= \mu(E_{n_0}) - \lim_{n \to \infty} \mu(E_n).$$

Furthermore, we observe that

$$\bigcup_{n=1}^{\infty} F_n = E_{n_0} \setminus \bigcap_{n=n_0}^{\infty} E_n.$$

Thus

$$\mu\left(\bigcup_{n=n_0}^{\infty} F_n\right) = \mu\left(E_{n_0} \setminus \bigcap_{n=n_0}^{\infty} E_n\right) = \mu(E_{n_0}) - \mu\left(\bigcap_{n=n_0}^{\infty} E_n\right)$$
$$= \mu(E_{n_0}) - \mu\left(\bigcap_{n=1}^{\infty} E_n\right).$$

It follows that indeed

$$\mu\left(\bigcap_{n=1}^{\infty} E_n\right) = \lim_{n \to \infty} \mu(E_n).$$

#### Exercise 4.1.1

Let  $(X, \mathfrak{M}, \mu)$  be a measure space. Show that

- 1.  $\mu$  is finite iff  $\mu(X) < \infty$ .
- 2.  $\mu$  is  $\sigma$ -finite implies that  $\mu$  is semi-finite.

# Solution

1. This is rather simple.

 $(\Longrightarrow)$   $\mu$  is finite implies that each  $E \in \mathfrak{M}$  has a finite measure. In particular,  $X \in \mathfrak{M}$ , and so  $\mu(X) < \infty$ .

( ⇐ )  $\forall E \in \mathfrak{M}, E \subseteq X$ , thus by the first item in Pheorem 5, we have  $\mu(E) \leq \mu(X) < \infty$ . Thus  $\mu$  is finite.

2.  $\mu$  being  $\sigma$ -finite means that if  $X = \bigcup_{n=1}^{\infty} X_n$  where  $X_n \in \mathfrak{M}$ , then  $\mu(X_n) < \infty$  for each n. Let  $E \in \mathfrak{M}$  such that  $\mu(E) = \infty$ . If we take

$$E_n = X_n \cap E$$
,

then  $\mu(E_n) < \infty$  for each  $n \in \mathbb{N}$ . Then, taking a union of any finite number of these  $E_n$ 's will give us a subset of E with a finite measure. Hence,  $\mu$  is indeed semi-finite.

### **■** Definition 11 (Null Set of a Measure)

Let  $(X, \mathfrak{M}, \mu)$  be a measure space. The set

$$\mathcal{N} := \{ N \in \mathfrak{M} : \mu(N) = 0 \}$$

is called the  $\mu$ -null set, or the null set of the measure  $\mu$ .

#### Remark 4.1.1

1. If  $N_i \in \mathcal{N}$ , then  $\bigcup_{n=1}^{\infty} N_i \in \mathcal{N}$ .

<sup>2</sup> Requires elab

2. If  $N \in \mathcal{N}$ , and  $E \in \mathfrak{M}$  and  $E \subseteq N$ , then  $E \in \mathcal{N}$ .

It is important to note there that the highlighted condition is required, since not all subsets of N are measurable.

3. N is **not** a  $\sigma$ -algebra. If we picked an X such that  $\mu(X) \neq 0$ , then  $\emptyset \in N$  but  $X \notin N$ .

# **■** Definition 12 (Complete Measure Space)

Let  $(X, \mathfrak{M}, \mu)$  be a measure space. We say that the space is **complete** if  $N \in \mathcal{N}$  and  $E \subseteq N$ , then  $E \in \mathfrak{M}$ . In this case, we also say that  $\mu$  is a **complete measure** on  $\mathfrak{M}$ .

#### **Remark 4.1.2**

By the first item in  $\blacksquare$  Theorem 5, we have that if  $\mu(E) = 0$ , and so  $E \in \mathcal{N}$  as

## **■** Theorem 6 (Extending the Measurable Sets)

Let  $(X, \mathfrak{M}, \mu)$  be a measure space and

$$\mathcal{N} := \{ N \in \mathfrak{M} \mid \mu(N) = 0 \}.$$

Consider

$$\overline{\mathfrak{M}} := \{ E \cup F \mid E \in \mathfrak{M}, F \subseteq N \in \mathcal{N} \}.$$

Then  $\overline{\mathfrak{M}}$  is a  $\sigma$ -algebra which contains  $\mathfrak{M}.$  Furthermore, if we define  $\overline{\mu}:$  $\overline{\mathfrak{M}} \to [0,\infty] \, as$ 

$$\overline{\mu}(E \cup F) = \mu(E),$$

then  $\overline{\mu}$  is a well-defined measure on  $\overline{\mathfrak{M}}$ .

*Moreover, if*  $\nu : \overline{\mathbb{M}} \to [0, \infty]$  *is any measure such that*  $\nu(E) = \mu(E)$  *for* all  $E \in \mathfrak{M}$ , then  $v = \overline{\mu}$ .

## *∠ Lecture 5 Sep 13th 2019*

## 5.1 Measures (Continued 3)

## Proof (Extending the Measurable Sets)

 $\overline{\mathbb{M}}$  is a  $\sigma$ -algebra Since  $\emptyset \in \mathbb{M}$  and  $\emptyset \subseteq N$  for any  $N \in \mathcal{N}$ , it is clear that  $\emptyset \in \overline{\mathbb{M}}$ .

Now, for  $E \cup F \in \overline{\mathfrak{M}}$ , if we suppose  $F \subseteq N \in \mathcal{N}$ , then

$$(E \cup F)^C = (E \cup N)^C \cup (N \setminus E \cup F) \in \overline{\mathfrak{M}}$$

since  $E \cup N \in \mathfrak{M}$  and  $N \setminus (E \cup F) \in \mathcal{N}$ .

Let  $\{E_n \cup F_n\}_{n=1}^{\infty} \subseteq \overline{\mathfrak{M}}$ . Then we observe that

$$\bigcup_{n=1}^{\infty} (E_n \cup F_n) = \bigcup_{n=1}^{\infty} E_n \cup \bigcup_{n=1}^{\infty} F_n \in \overline{\mathbb{M}}.$$

Well-definedness of  $\overline{\mu}$  Let  $E_1 \cup F_1 = E_2 \cup F_2 \in \overline{\mathfrak{M}}$ . Suppose  $F_1 \subseteq N_1, F_2 \subseteq N_2 \in \mathcal{N}$ . WTS

$$\mu(E_1) = \overline{\mu}(E_1 \cup F_1) = \overline{\mu}(E_2 \cup F_2) = \mu(E_2)$$

Notice that

$$E_1 \subseteq E_1 \cup F_1 = E_2 \cup F_2 \subseteq E_2 \cup N_2,$$

and

$$E_2 \subseteq E_2 \cup F_2 = E_1 \cup F_1 \subseteq E_1 \cup N_1$$
.

By Prheorem 5, in particular, by subadditivity, we have that

$$\mu(E_1) \le \mu(E_2 \cup N_2) \le \mu(E_2) + 0 = \mu(E_2)$$

and

$$\mu(E_2) \le \mu(E_1 \cup N_1) \le \mu(E_1) + 0 = \mu(E_1).$$

It follows that  $\mu(E_1) = \mu(E_2)$ , as required.

#### $\overline{\mu}$ is a measure

1. Since  $\emptyset \in \mathfrak{M}$  and  $\emptyset \in \mathcal{N}$ ,  $\overline{\mu}$  is defined for  $\emptyset$ , and

$$\overline{\mu}(\emptyset) = \mu(\emptyset) = 0.$$

2. Let  $\{E_n \cup F_n\}_{n=1}^{\infty} \subseteq \overline{\mathbb{M}}$  be a pairwise disjoint collection. We observe that

$$\overline{\mu}\left(\bigcup_{n=1}^{\infty}(E_n \cup F_n)\right) = \overline{\mu}\left(\bigcup_{n=1}^{\infty}E_n \cup \bigcup_{n=1}^{\infty}F_n\right)$$
$$= \mu\left(\bigcup_{n=1}^{\infty}E_n\right)$$
$$= \sum_{n=1}^{\infty}\mu(E_n),$$

and

$$\sum_{n=1}^{\infty} \overline{\mu}(E_n \cup F_n) = \sum_{n=1}^{\infty} \mu(E_n).$$

Hence

$$\overline{\mu}\left(\bigcup_{n=1}^{\infty}(E_n\cup F_n)\right)=\sum_{n=1}^{\infty}\overline{\mu}(E_n\cup F_n).$$

 $v = \overline{\mu}$  Let  $E \cup F \in \overline{\mathfrak{M}}$ . Suppose  $F \subseteq N \in \mathfrak{M}$  By monotonicity,

$$\overline{\mu}(E \cup F) = \mu(E) = \nu(E) \le \nu(E \cup F).$$

By subadditivity,

$$\nu(E \cup F) \le \nu(E) + \nu(F) \le \mu(E) + \nu(N) \le \overline{\mu}(E \cup F) + \mu(N) = \overline{\mu}(E \cup F) + 0.$$

Thus, indeed,

$$\nu(E \cup F) = \overline{\mu}(E \cup F).$$

### *5.2 The Outer Measure*

In this section, we will show that one way we can construct a measure is by using an outer measure.

## **■** Definition 13 (Outer Measure)

Given a set X, a function

$$\mu^*: \mathcal{P}(X) \Longrightarrow [0, \infty]$$

is called an outer measure if

- 1.  $\mu^*(\emptyset) = 0$ ;
- 2. (monotonicity) if  $E \subseteq F$ , then  $\mu^*(E) \le \mu^*(F)$ ; and
- 3. (countable subadditivity) if  $\{A_n\}_n \subseteq \mathcal{P}(X)$ , then

$$\mu^* \left( \bigcup_{n=1}^{\infty} A_n \right) \le \sum_{n=1}^{\infty} \mu^*(A_n).$$

Coming from PMATH450, we have seen an example of an outer measure.

## **♦** Proposition 7 (Lebesgue's Outer Measure)

Given  $E \subseteq \mathbb{R}$ , consider

$$\mu^*(E) := \inf \left\{ \sum_{n=1}^{\infty} (b_n - a_n) : E \subseteq \bigcup_{n=1}^{\infty} (a_n, b_n) \right\}.$$

 $\mu^*$  is Lebesgue's outer measure.

#### Proof

- 1. It is clear that  $\mu^*(\emptyset) = \emptyset$ , since we can pick all  $(a_n, b_n) = \emptyset$ .
- 2. Suppose  $A \subseteq B \subseteq \mathbb{R}$ . It is clear that any collection of intervals whose union contain B will contain A, but there are such collections for A that do not contain B. This means that

$$\mu^*(A) \le \mu^*(B)$$

by the property of the infimum.

3. Let  $E = \bigcup_{i=1}^{\infty} E_i$ . WTS  $\mu^*(E) \le \sum_{i=1}^{\infty} \mu^*(E_i)$ .

Now if  $\mu^*(E_i) = \infty$  for any i, then the inequality is trivially true. Thus, wma  $\mu^*(E_i) < \infty$  for all i.

<sup>1</sup> Let  $\varepsilon > 0$ . By the definition of the infimum, for each i, we can pick a countable sequence  $\{(a_n^i, b_n^i)\}_{n=1}^{\infty} \subseteq \mathcal{P}(X)$  such that  $E_1 \subseteq \bigcup_{n=1}^{\infty} (a_n^i, b_n^i)$  and

<sup>1</sup> This is also a common trick in measure theory.

$$\sum_{n=1}^{\infty} (b_n^i - a_n^i) \le \mu^*(E_i) + \frac{\varepsilon}{2^i}.$$

Then

$$E = \bigcup_{i=1}^{\infty} E_i = \bigcup_{i=1}^{\infty} \bigcup_{n=1}^{\infty} (a_n^i, b_n^i).$$

And so it follows that

$$\mu^*(E) \le \sum_{i=1}^{\infty} \sum_{n=1}^{\infty} (b_n^i - a_n^i)$$

$$\le \sum_{i=1}^{\infty} \mu^*(E_i) + \sum_{i=1}^{\infty} \frac{\varepsilon}{2^i}$$

$$= \sum_{i=1}^{\infty} \mu^*(E_i) + \varepsilon.$$

Since  $\varepsilon$  was arbitrary, it follows that

$$\mu^*(E) \le \sum_{i=1}^{\infty} \mu^*(E_i).$$

Show that had we defined Lebesgue's outer measure with closed intervals, i.e.

$$\tilde{\mu}^*(E) := \inf \left\{ \sum_{n=1}^{\infty} (b_n - a_n) : E \subseteq \bigcup_{n=1}^{\infty} [a_n, b_n] \right\},$$

 $\tilde{\mu}^*$  is still an outer measure.

*In fact, we can do so for half-open intervals.* 

#### Example 5.2.1 (Lebesgue-Stieltjes Outer Measure)

Let  $F : \mathbb{R} \to \mathbb{R}$  be an increasing function that is continuous from the right. Let

$$\mu^*(E) := \inf \left\{ \sum_{n=1}^{\infty} (F(b_n) - F(a_n)) : E \subseteq \bigcup_{n=1}^{\infty} (a_n, b_n) \right\}.$$

Then  $\mu^*$  is an outer measure.

#### Remark 5.2.1

Again, we could have defined the above outer measure using open or closed intervals.

## Example 5.2.2 (Lebesgue's Outer Measure on $\mathbb{R}^2$ )

Let  $E \subseteq \mathbb{R}^2$ , and

$$\mu^*(E) := \inf \left\{ \sum_{n=1}^{\infty} A(R_n) : E \subseteq \bigcup_{n=1}^{\infty} R_n \right\},$$

where *A* is the 'area' function, and  $R_n = (a_n, b_n) \times (c_n, d_n)$  are open rectangles. Then  $\mu^*$  is an outer measure.

#### Remark 5.2.2

- 1. Again, we can define the above outer measure using closed rectangles, or half-open rectangles.
- 2. We can continue defining an outer measure for  $\mathbb{R}^3$  using cubes, for  $\mathbb{R}^4$ using hypercubes, and so on.

We want to now show that given an outer measure, we can always construct a measure. This is known as Carathéodory's Theorem.

This requires the following definition:

## **E** Definition 14 ( $\mu^*$ -measurability)

A set  $A \subseteq X$  is said to be  $\mu^*$ -measurable if  $\forall E \subseteq X$ ,

$$\mu^*(E) = \mu^*(E \cap A) + \mu^*(E \cap A^C).$$

#### Remark 5.2.3

1. By subadditivity, we always have

$$\mu^*(E) \le \mu^*(E \cap A) + \mu^*(E \cap A^C),$$

since 
$$E = (E \cap A) \cup (E \cap A^C)$$
.

2. Note that  $E \cap A^C = E \setminus A$ . In a sense, A is said to be  $\mu^*$ -measurable if it can slice any subset of X such that we have additivity of the sliced parts. We may also say that A is a 'universal slicer'.

## **■**Theorem 8 (Carathéodory's Theorem)

If  $\mu^*$  is an outer measure on a set X, let

$$\mathfrak{M} := \{ A \subseteq X : A \text{ is } \mu^*\text{-measureable} \}.$$

*Then*  $\mathfrak{M}$  *is a*  $\sigma$ *-algebra, and we set* 

$$\mu:\mathfrak{M}\to[0,\infty]$$

such that

$$\mu(A) = \mu^*(A).$$

Then  $\mu$  is a complete measure on  $\mathfrak{M}$ .

## 6.1 The Outer Measure (Continued)

#### ♣ Homework (Homework 4)

Let  $\mathfrak{M}$  be an algebra of sets on X, and whenever  $\{A_n\}_{n\in\mathbb{N}}\subseteq\mathfrak{M}$  is a disjoint collection of sets, then  $\bigcup_n A_n\in\mathfrak{M}$ . Then  $\mathfrak{M}$  is a  $\sigma$ -algebra.

#### **\$**<sup>a</sup> Homework (Homework 5)

Recall that Lebesgue's Outer Measure on  $\mathbb{R}$  is defined as

$$\mu^*(E) := \inf \left\{ \sum_{n=1}^{\infty} (b_n - a_n) : E \subseteq \bigcup_{n=1}^{\infty} (a_n, b_n) \right\}.$$

Prove that we can equivalently define

$$\mu^*(E) := \inf \left\{ \sum_{n=1}^{\infty} (b_n - a_n) : E \subseteq \bigcup_{n=1}^{\infty} (a_n, b_n) \right\}.$$

Similarly, Lebesgue's Outer Measure on  $\mathbb{R}^2$  is defined as

$$\mu_2^*(E) = \inf \left\{ \sum_{n=1}^{\infty} (b_n - a_n)(d_n - c_n) : E \subseteq \bigcup_{n=1}^{\infty} (a_n, b_n) \times (c_n, d_n) \right\}.$$

Prove that we can equivalently define

$$\mu_2^*(E) = \inf \left\{ \sum_{n=1}^{\infty} (b_n - a_n)(d_n - c_n) : E \subseteq \bigcup_{n=1}^{\infty} (a_n, b_n] \times (c_n, d_n] \right\}.$$

#### **■** Definition 15 (Metric Outer Measure)

Let (X, d) be a metric space, and  $A, B \subseteq X$ , and

$$d(A, B) = \inf \{ d(x, y) : x \in A, y \in B \}.$$

An outer measure,  $\mu^*$ , on X is called a *metric outer measure* if whenever d(A,B) > 0, then

$$\mu^*(A \cup B) = \mu^*(A) + \mu^*(B).$$

#### \* Homework (Homework 6)

*Prove that Lebesgue's Outer Measure on*  $\mathbb{R}$  *is a metric outer measure.* 

## Proof (Carathéodory's Theorem)

#### $\mathfrak{M}$ is a $\sigma$ -algebra

 $\emptyset \in \mathfrak{M}$  Given any  $E \subseteq X$ , we observe that

$$\mu^*(E \cap \emptyset) + \mu^*(E \cap \emptyset^C) = \mu^*(\emptyset) + \mu^*(E \cap X)$$
$$= 0 + \mu^*(E) = \mu^*(E).$$

 $A \in \mathfrak{M} \implies A^{C} \in \mathfrak{M}$  Observe that given any  $E \subseteq X$ ,

$$\mu^*(E \cap A^C) + \mu^*(E \cap (A^C)^C) = \mu^*(E \cap A^C) + \mu^*(E \cap A) = \mu^*(E).$$

Thus  $A^C \in \mathfrak{M}$ .

 $^{\rm 1}$  To show that  $\mathfrak M$  is closed under countable unions, we break the work into several steps.

<sup>1</sup> For a deep dive, see Appendix A.1.

 $A, B \in \mathfrak{M} \implies A \cup B \in \mathfrak{M}$  Since  $A \in \mathfrak{M}$ , we have

$$\mu^*(E) = \mu^*(E \cap A) + \mu^*(E \cap A^C).$$

Since  $B \in \mathfrak{M}$ ,

$$\mu^{*}(E) = \mu^{*}(E \cap A) + \mu^{*}(E \cap A^{C})$$

$$= \mu^{*}(E \cap A \cap B) + \mu^{*}(E \cap A \cap B^{C})$$

$$+ \mu^{*}(E \cap A^{C} \cap B) + \mu^{*}(E \cap A^{C} \cap B^{C})$$

$$= \mu^{*}(E \cap A \cap B) + \mu^{*}(E \cap A \cap B^{C})$$

$$+ \mu^{*}(E \cap A^{C} \cap B) + \mu^{*}(E \cap (A \cup B)^{C})$$

Notice that

$$E \cap (A \cup B) = [E \cap A \cap B] \cup [E \cap A^C \cap B] \cup [E \cap A \cap B^C].$$

Thus

$$\begin{split} \mu^*(E) &= \mu^*(E \cap A) + \mu^*(E \cap A^C) \\ &= \mu^*(E \cap A \cap B) + \mu^*(E \cap A \cap B^C) \\ &+ \mu^*(E \cap A^C \cap B) + \mu^*(E \cap (A \cup B)^C) \\ &\geq \mu^*(E \cap (A \cup B)) + \mu^*(E \cap (A \cup B)^C). \end{split}$$

Thus  $A \cup B \in \mathfrak{M}$ .

Consequently, by induction, we have that  $\forall \{A_n\}_n \subseteq \mathfrak{M}$ ,

$$\bigcup_{n=1}^{N} A_n \in \mathfrak{M}$$

for all  $N \in \mathbb{N}$ .

Now  $\mathfrak M$  is an algebra of sets. By Homework 4, it suffices for us to prove the following to show that  $\mathfrak M$  is a  $\sigma$ -algebra of sets.

 $\forall \{A_n\}_n \subseteq \mathfrak{M} \text{ disjoint, } \Longrightarrow \bigcup_n A_n \in \mathfrak{M} \text{ Let } B_N = \bigcup_{n=1}^N A_n. \text{ We first}$ require the following lemma:

 $\forall E \subseteq X, \ \mu^*(E \cap B_N) = \sum_{n=1}^N \mu^*(E \cap A_n)$  Notice that for any  $n \in \mathbb{N}$ ,  $A_n \in \mathfrak{M}$ , and so

$$\mu^*(E \cap B_N) = \mu^*(E \cap B_N \cap A_n) + \mu^*(E \cap B_N \cap A_n^C)$$
  
= \mu^\*(E \cap A\_n) + \mu^\*(E \cap B\_{N-1}).

The desired result follows by induction.  $\dashv$ 

Let  $B = \bigcup_{n=1}^{\infty} A_n$ . Then

$$\mu^*(E \cap B) \le \sum_{n=1}^{\infty} \mu^*(E \cap A_n)$$

by subadditivity.

Now  $B_N \subseteq B$  for each  $N \in \mathbb{N}$ . This implies that  $B_N^C \supseteq B^C$ , and so by monotonicity,

$$\mu^*(E\cap B_N^C) \geq \mu^*(E\cap B^C).$$

Thus, for every  $N \in \mathbb{N}$ ,

$$\mu^*(E) = \mu^*(E \cap B_N) + \mu^*(E \cap B_N^C)$$
 
$$\geq \sum_{n=1}^N \mu^*(E \cap A_n) + \mu^*(E \cap B^C).$$

It follows that

$$\mu^{*}(E) \ge \sum_{n=1}^{\infty} \mu^{*}(E \cap A_{n}) + \mu^{*}(E \cap B^{C})$$
$$\ge \mu^{*}(E \cap B) + \mu^{*}(E \cap B^{C}).$$

With Homework 4,  $\mathfrak{M}$  is a  $\sigma$ -algebra.

#### $\mu$ is a measure

- $\bullet \quad \mu(\emptyset) = \mu^*(\emptyset) = 0.$
- Let  $\{A_n\}_n \subseteq \mathfrak{M}$  be a disjoint collection of sets, and  $B = \bigcup_{n=1}^{\infty} A_n$ . Then by a similar argument as the end of the last 'part',

$$\mu(B) = \mu^{*}(B)$$

$$\geq \sum_{n=1}^{\infty} \mu^{*}(B \cap A_{n}) + \mu^{*}(B \cap B^{C})$$

$$= \sum_{n=1}^{\infty} \mu^{*}(B \cap A_{n}) + 0$$

$$= \sum_{n=1}^{\infty} \mu^{*}(A_{n}) = \sum_{n=1}^{\infty} \mu(A_{n}).$$

Thus

$$\mu(B) = \mu\left(\bigcup_{n=1}^{\infty} A_n\right) = \sum_{n=1}^{\infty} \mu(A_n).$$

 $\mu$  is complete Let  $A \in \mathcal{N}$  and  $B \subseteq A$ . By monotonicity,  $\mu(B) =$  $\mu^*(B) \le \mu^*(A) = 0$ . Then

$$\mu^*(E\cap B) + \mu^*(E\cap B^C) = 0 + \mu^*(E\cap B^C) \leq \mu^*(E)$$

by monotonicity. Thus  $B \in \mathfrak{M}$ . Thus  $\mu$  is complete.

We would like to make sure that

- 1. there are many sets that are measurable; and
- 2. the notion of a measure covers our notion of length.

We shall see this with the Metric Outer Measure, and that the measurable sets is at least the Borel set.

#### 6.2 The Lebesgue-Stieltjes Outer Measure

This outer measure is motivated by probability theory. The idea is that we consider the measure space  $(\Omega, \mathfrak{M}, P)$ , where  $\Omega$  is the sample space set,  $\mathfrak{M}$  is a  $\sigma$ -algebra on  $\Omega$ , and P is the probability measure, i.e.  $P(\Omega) = 1$ .

We then define a random variable, which is a function  $X : \Omega \to \mathbb{R}$ . The cumulative distribution function (cdf) is defined as

$$F_X(t) := P(\{\omega : X(\omega) \le t\}),$$

and it has these properties:

- 1.  $F_X$  is increasing; and
- 2.  $F_X$  is right-continuous.

#### Example 6.2.1

Let  $\Omega = \{H, T\}$ , and define the probability measure as

$$P({H}) = \frac{1}{2} = P({T}).$$

We can define

$$X(T) = 0$$
 and  $X(H) = 1$ .

Then

$$P(\{\omega : X(\omega) = 1\}) = P(\{H\}) = \frac{1}{2}$$

and

$$P(\{\omega: X(\omega) = 0\}) = P(\{T\}) = \frac{1}{2}.$$

In the context of probability, we often see the shorthand

$$P(X = t) = P(\{\omega : X(\omega) = t\}).$$




Figure 6.1: Simple example of a cdf

## **■** Definition 16 (Lebesgue-Stieltjes Outer Measure)

Let  $F:\mathbb{R}\to\mathbb{R}$  be an increasing function that is continuous from the right. Let

$$\mu^*(E) := \inf \left\{ \sum_{n=1}^{\infty} (F(b_n) - F(a_n)) : E \subseteq \bigcup_{n=1}^{\infty} (a_n, b_n] \right\}.$$

Then  $\mu^*$  is an outer measure.

#### Exercise 6.2.1

We mentioned that the above is indeed an outer measure in Example 5.2.1. Prove this.

## *∠ Lecture 7 Sep 18th 2019*

## 7.1 The Lebesgue-Stieltjes Outer Measure (Continued)

#### **■** Theorem 9 (Carathéodory's Second Theorem)

Let (X, d) be a metric space, and  $\mu^*$  a Metric Outer Measure. Then every Borel set is  $\mu^*$ -measurability.

#### Proof

By Carathéodory's Theorem,

$$\mathfrak{M} = \{ A \subseteq X : A \text{ is } \mu^*\text{-measurable} \}$$

is a  $\sigma$ -algebra. Then our statement says that  $\mathfrak{B}(X)\subseteq \mathfrak{M}$ . Thus, it suffices for us to show that if  $U\in \mathfrak{B}(X)$ , i.e. if  $U\subseteq X$  is open, then  $U\in \mathfrak{M}$ . In particular, WTS  $\forall E\subseteq X$ ,

$$\mu^*(E) = \mu^*(E \cap U) + \mu^*(E \cap U^C).$$

Again, by subadditivity,

$$E = (E \cap U) \cup (E \cap U^C) \implies \mu^(E) \leq \mu^*(E \cap U) + \mu^*(E \cap U^C).$$

Thus it suffices for us to show that

$$\mu^*(E) \ge \mu^*(E \cap U) + \mu^{(E)}(E \cap U^{C}).$$

Now if  $\mu^*(E) = \infty$ , this is trivially true. WMA  $\mu^*(E) < \infty$ .

<sup>1</sup> Consider  $\{A_k\}_k \subseteq \mathcal{P}(X)$  such that

$$A_k := \left\{ x \in E \cap U : d(x, E \cap U^C) \geq \frac{1}{k} \right\} \subseteq E \cap U.$$

It is clear that  $A_1 \subseteq A_2 \subseteq A_3 \subseteq ...$ , i.e.  $\{A_k\}_k$  is an increasing sequence of sets. Also,  $\bigcup_k A_k = E \cap U$ .

For each  $k \in \mathbb{N}$ , notice that  $A_k \cup (E \cap U^C) \subseteq E$ . Thus by subadditivity and additivity over disjoint sets, for every k, we have

$$\mu^*(E) \ge \mu^*(A_k \cup (E \cap U^C)) = \mu^*(A_k) + \mu^*(E \cap U^C).$$

Since  $\{A_k\}_k$  is an increasing sequence of sets, it follows that

$$\mu^*(E) \ge \lim_{k \to \infty} \mu^*(A_k) + \mu^*(E \cap U^C).$$

Claim:  $\lim_{k\to\infty} \mu^*(A_k) = \mu^*(E\cap U)$  Since  $A_k \subseteq E\cap U$ , by subadditivity,

$$\mu^*(A_k) \leq \mu^*(E \cap U).$$

It remains to prove the other inequality.

<sup>2</sup> Let 
$$D_1 = A_1$$
,  $D_2 = A_2 \setminus A_1$ , ...  $D_n = A_n \setminus A_{n-1}$ . Then, notice that 
$$E \cap U = \bigcup_n D_n = A_1 \cup D_2 \cup D_3 \cup \dots$$
$$= A_2 \cup D_3 \cup D_4 \cup \dots$$
$$= A_n \cup D_{n+1} \cup D_{n+2} \cup \dots$$

since  $\{A_n\}$  is an increasing sequence of sets. Now for  $x \in D_n$ , we have that  $x \in A_n$  but  $x \notin A_{n-1}$ . <sup>3</sup> In particular, we have

$$\frac{1}{n} \le d(x, E \cap U^C) < \frac{1}{n-1}.$$

<sup>4</sup> Let  $m \ge n + 2$ . Consider  $y \in D_m$ ,  $x \in D_n$  and  $z \in E \cap U^C$ . Then we know by the triangle inequality that

$$\frac{1}{n} \le d(x, z) \le d(x, y) + d(y, z).$$

 $^1$  We look at points that get increasingly closer to the edge of the set  $E \cap U$ , or in other words, increasingly closer to  $E \cap U^C$ .

<sup>2</sup> This part here requires an escape from where we already are. If your head is in the muddle, stop reading, go out, walk, and then come back.

Here, we ask ourselves: so what if we look at how much the  $A_k$ 's change as k increases?

<sup>3</sup> We see that the  $D_n$ 's form some kind of a ring-like partitioning of  $E \cup U$ .

 $^4$  Let's look at putting every odd  $D_n$ 's together, and see how far apart are they. Directly looking at  $D_n$ 's altogether is difficult because then their boundaries get muddled together.

We may then pick  $z_0 \in E \cap U^C$  such that

$$d(y,z_0)<\frac{1}{m-1}.$$

Then

$$\frac{1}{n} \le d(x, z_0) < d(x, y) + \frac{1}{m - 1},$$

and so

$$\frac{1}{n} - \frac{1}{m-1} < d(x, y).$$

Notice that

$$\frac{1}{n} - \frac{1}{m-1} \ge \frac{1}{n} - \frac{1}{n+2-1} = \frac{1}{n} - \frac{1}{n-1}.$$

Therefore,  $\forall x \in D_n$  and  $y \in D_m$ , we have

$$\frac{1}{n} - \frac{1}{n-1} < d(x,y).$$

In other words,

$$d(D_n, D_m) > 0$$

as long as  $m \ge n + 2$ .

Since  $\mu^*$  is a Metric Outer Measure, it follows that

$$\mu^* \left( \bigcup_{n \text{ odd}} D_n \right) = \sum_{n \text{ odd}} \mu^* (D_n).$$

Since  $D_1 \cup D_3 \cup \ldots \subseteq E \cap U$ , by subadditivity,

$$\mu^*(D_1 \cup D_3 \cup \ldots) \leq \mu^*(E \cap U) < \infty.$$

In particular,

$$\sum_{n \text{ odd}} \mu^*(D_n) < \infty.$$

Similarly, we can show that

$$\mu^* \left( \bigcup_{n \text{ even}} D_n \right) = \sum_{n \text{ even}} \mu^*(D_n) < \infty.$$

Putting the two together, we have

$$\mu^*\left(\bigcup_n D_n\right) = \sum_n \mu^*(D_n) < \infty.$$

Finally, since  $E \cap U = A_n \cup D_{n+1} \cup D_{n+2} \cup ...$ , by subadditivity,

$$\mu^*(E \cap U) \le \mu^*(A_n) + \mu^*\left(\bigcup_{m>n} D_m\right) = \mu^*(A_n) + \sum_{m=n+1}^{\infty} \mu^*(D_m).$$

Since  $\sum_n \mu^*(D_n) < \infty$ , the tail

$$\sum_{m=n+1}^{\infty} \mu^*(D_m) \to 0$$

as  $n \to \infty$ . Therefore,

$$\mu^*(E \cap U) \le \lim_{n \to \infty} \mu^*(A_n) + \lim_{n \to \infty} \sum_{m=n+1}^{\infty} \mu^*(D_m)$$
$$= \lim_{n \to \infty} \mu^*(A_n),$$

as required.

**♦** Proposition 10 (Lebesgue-Stieltjes Outer Measure on Halfopen Intervals)

Let  $\mu_F^*$  be the Lebesgue-Stieltjes Outer Measure. Then for  $a < b \in \mathbb{R}$ , we have

$$\mu_F^*((a,b]) = F(b) - F(a).$$

#### Proof

First, notice that  $(a, b] \subseteq (a, b]$ , and so

$$\mu_{\scriptscriptstyle F}^*((a,b]) \le F(b) - F(a)$$

by definition.

Let  $\varepsilon > 0$ . Pick a covering  $(a, b] \subseteq \bigcup_n (a_n, b_n]$  such that

$$\sum_n (F(b_n) - F(a_n)) \le \mu_F^*((a,b]) + \varepsilon.$$

By right-continuity of F, we may pick  $b'_n > b_n$  such that

$$F(b'_n) < F(b_n) + \frac{\varepsilon}{2}.$$

Similarly, we can pick a' > a such that  $F(a') < F(a) + \varepsilon$ . Then

$$[a,b]\subseteq (a,b]\subseteq \bigcup_n (a_n,b_n]\subseteq \bigcup_n (a_n,b'_n).$$

By **compactness**, there exists a finite subcover, i.e.  $\exists N \in \mathbb{N}$  such that

$$[a',b] \subseteq \bigcup_{k=1}^{N} (a_{n_k},b'_{n_k}).$$

The proof shall be completed in the next lecture.



# A.1 Proving that $\mathfrak M$ is closed under countable unions in Carathéodory's Theorem

This section is created in reference to the proof for Carathéodory's Theorem.

We have

$$\mathfrak{M} = \{ A \subseteq X : A \text{ is } \mu^*\text{-measurable } \}$$

where  $\mu^*$  is an outer measure. We wanted to show that  $\mathfrak{M}$  is a  $\sigma$ -algebra. In particular, the hard problem was to show that  $\mathfrak{M}$  is closed under countable unions.

Consider  $\{A_n\}_n \subseteq \mathfrak{M}$ . Thinking from behind, WTS  $\forall E \subseteq X$ ,

$$\mu^{*}(E) \geq \mu^{*} \left( E \cap \bigcup_{n} A_{n} \right) + \mu^{*} \left( E \cap \left( \bigcup_{n} A_{n} \right)^{C} \right)$$
$$= \mu^{*} \left( E \cap \bigcup_{n} A_{n} \right) + \mu^{*} \left( E \cap \left( \bigcap_{n} A_{n}^{C} \right) \right).$$

For simplicity, write  $B = \bigcup_n A_n$ . WTS

$$\mu^*(E) \ge \mu^*(E \cap B) + \mu^*(E \cap B^C). \tag{*}$$

Also thinking from behind, if  $\mathfrak M$  is a  $\sigma$ -algebra,  $^1$  then it must be an algebra (of sets). We showed that  $\mathfrak M$  is closed under complementation.

If  $\mathfrak{M}$  is closed under finite unions, <sup>2</sup> then for each  $N \in \mathbb{N}$ ,

<sup>2</sup> Unproved point 1

<sup>&</sup>lt;sup>1</sup> Useful links: Algebra of Sets, σ-Algebra of Sets.

$$\mu^*(E) = \mu^* \left( E \cap \bigcup_{n=1}^N A_n \right) + \mu^* \left( E \cap \left( \bigcup_{n=1}^N A_n \right)^C \right).$$

Let  $B_N := \bigcup_{n=1}^N A_n \in \mathfrak{M}$ . Then

$$\mu^*(E) = \mu^*(E \cap B_N) + \mu^*(E \cap B_N^C)$$
 (†)

for each  $N \in \mathbb{N}$ .

Notice that

$$B_N = \bigcup_{n=1}^N A_n \subseteq \bigcup_{n=1}^\infty A_n = B.$$

Consequently,

$$\implies B^C \supseteq B_N^C \implies \mu^*(B^C) \le \mu^*(B_N^C)$$

by the monotonicity of the outer measure.

As a result, looking at Equation (\*) and Equation (†), we see that

$$\mu^*(E) = \mu^*(E \cap B_N) + \mu^*(E \cap B_N^C)$$
  
  $\geq \mu^*(E \cap B_N) + \mu^*(E \cap B^C)$ 

for each  $N \in \mathbb{N}$ .

We are in quite the predicament at this point. We need to do something about  $\mu^*(E \cap B_N)$  and somehow relate it to  $\mu^*(E \cap B)$ . We can try and see that

$$\mu^*(E \cap B_N) \le \sum_{n=1}^N \mu^*(E \cap A_n).$$

Notice that in the case of equality, we would have

$$\mu^*(E) \geq \sum_{n=1}^N \mu^*(E \cap A_n) + \mu^*(E \cap B^C)$$

for all  $N \in \mathbb{N}$ . Since  $\{\sum_{n=1}^{N} \mu^*(E \cap A_n)\}_N$  is an increasing sequence in  $\mathbb{R}$ , we have

$$\mu^*(E) \ge \sum_{n=1}^{\infty} \mu^*(E \cap A_n) + \mu^*(E \cap B^C),$$

and

$$\sum_{n=1}^{\infty} \mu^*(E \cap A_n) \ge \mu^*(E \cap B)$$

by subadditivity since  $B = \bigcup_{n=1}^{\infty} A_n$ .

Unfortunately, the equality does not always hold. But, since  $\mu^*$  is an outer measure, we can make an educated guess  $^3$  that given  $\{A_n\}_n$  a disjoint collection of sets,

<sup>3</sup> Unproved point 2

$$\mu^*\left(\bigcup_{n=1}^N A_n\right) = \sum_{n=1}^N \mu^*(A_n).$$

Our work becomes even easier with the realization of Homework 4. Proving that all of our above argument works for the case of  $\{A_n\}_n \subseteq$  ${\mathfrak M}$  being disjoint, is sufficient to prove that  ${\mathfrak M}$  is indeed a  $\sigma\text{-algebra}.$ 



## B.1 Re-represent an arbitrary union using disjoint sets

A common trick in measure theory, especially when it comes to a collection of sets, is to represent its union as a disjoint union of sets. This is a useful trick because measures simply add over disjoint sets, instead of just having subadditivity.

#### Example B.1.1

Given a collection  $\{A_n\}_n$  of sets, we may define a collection of disjoint sets whose union is  $\bigcup_n A_n$  as such:

$$F_{1} = A_{1}$$

$$F_{2} = A_{2} \setminus A_{1}$$

$$F_{3} = A_{3} \setminus (A_{1} \cup A_{2})$$

$$\vdots$$

$$F_{n} = A_{n} \setminus \bigcup_{i=1}^{n-1} A_{i}$$

$$\vdots$$

#### Example B.1.2

Given an increasing collection  $\{A_n\}_n$  of sets, i.e.

$$A_1 \subseteq A_2 \subseteq A_3 \subseteq \ldots$$
,

we may represent the countable union of the  $A_n$ 's as such: let

$$F_1 = A_1$$

$$F_2 = A_2 \setminus A_1$$

$$F_3 = A_3 \setminus A_2$$

$$\vdots$$

$$F_n = A_n \setminus A_{n-1}$$

$$\vdots$$

#### Remark B.1.1

The reason why we simply consider  $F_n = A_n \setminus A_{n-1}$  instead of having to take a union up to the  $(n-1)^{th}$  set in Example B.1.2 is because

$$\bigcup_{i=1}^{n-1} A_i = A_{n-1}. (B.1)$$

The reader may also notice that Example B.1.2 is an application of Example B.1.1 just because of Equation (B.1).





$\mu$ -null set, 36	Lebesgue's Outer Measure, 41		
$\mu^*$ -measurability, $44$			
$\sigma$ -Algebra of Sets, 16	Measure, 25		
$\sigma$ -finite measure, 29	Measure Space, 27		
	Metric Outer Measure, 46		
Algebra of Sets, 15			
Almost always, 24	Null Set, 36		
Borel $\sigma$ -algebra, 21	Outer Measure, 41		
Carathéodory's Second Theorem,	partition, 11		
51	probability measure, 27		
Carathéodory's Theorem, 44	probability space, 27		
Characteristic Function, 25	Properties of Algebra of Sets, 15		
co-countable, 16			
cofinite, 16	random variable, 49		
complete measure, 36	Riemann integrable, 12		
Complete Measure Space, 36	Riemann Integration, 11		
Counting Measure, 27	Riemann lower integral, 12		
cumulative distribution function,	Riemann lower sum, 11		
49	Riemann upper integral, 12		
	Riemann upper sum, 11		
finite additivity, 29			
Finite measure, 29	Semi-finite measure, 29		
Generator of a $\sigma$ -algebra, 20			
Infinitely Often, 24			