


Student Info

Jingyi Zhuang

jz2907@columbia.edu

Homework 6

(A)



Yes, they match!

As mentioned in the instruction:

Note that the Matlab SVD algorithm may have obtained the negative of any one of the singular vectors in in Figure 2.

So here in my plot of Mode 1 to 5, the 1, 2 and 5 are the negative V_{mode_i} , just to make it look similar.

Script:

```
% initiate variables

clearvars;
load('MOR_EOF_228_Z.mat');

% A.

[U,S,V] = svd(Z,'econ');
x=linspace(-40,40,81);

figure(1);

for i = 1:5
    ax(i)=subplot('Position',[0.1 0.1+0.17*(5-i) 0.8 0.17]);

    % To match the paper Small 1998, the Mode 1, 2, 5 are -
    V(:,i), and the
    % Mode 3 and 4 are V(:,i), as mentioned in the problem
    instructions.

    if i == 1 || i == 2 || i == 5
        plot(x,-V(:,i),"k-","LineWidth",2);
    else
        plot(x,V(:,i),"k-","LineWidth",2);
    end

    % Make the ylim looks better:

    lim = max(abs(V(:,i)))/0.8;
```

```

ylim([-lim,lim])
legend(sprintf('Mode %g',i));


end

set(ax,'LineWidth',1,'FontSize',14,'YTickLabel','');
set(ax(1:4),'XTickLabel','');
xlabel('Distance(km)');

```

(B)

Reproduce the main plot in Figure 2 of Small (1998):



Script:

```

% B.

var = diag(S);
var_ttl = sum(var);
var_percentage = var/var_ttl;

figure(2);

```

```

hold on;

plot(1:length(var_percentage), var_percentage, "ko-",
"MarkerSize",8,"LineWidth",2);
plot(1:5, var_percentage(1:5),"ko-
", "MarkerSize",8,"LineWidth",2,'MarkerFaceColor','#000000');
text(20,0.1,sprintf('First 5 Sum is:
%.3f',sum(var_percentage(1:5))),'FontSize',16 );
set(gca,'LineWidth',1,'FontSize',14);
xlabel('Mode');
ylabel('% Variance');
hold off;

```

- The newline in `plot` function is caused by the `markdown` editor thus it's same line in `matlab`.


(C)

Why:

Because the singular value component refers to the same level as the data itself while the variance should be the square of that, the variance contribution of each mode is not calculated correctly by using

$$\%s_i = \frac{s_i}{\sum_1^N s_j}.$$

Instead, the variance contribution should be $\%s_i^2 = \frac{s_i^2}{\sum_1^N s_j^2}.$



So the corrected fraction of the total variance due to the first five singular vectors would be 0.844.

Script:

```
% C.

S2 = S^2;


var2 = diag(S2);
var2_ttl = sum(var2);
var2_percentage = var2/var2_ttl;

figure(3);
hold on;

plot(1:length(var2_percentage), var2_percentage, "ko-
", "MarkerSize",8,"LineWidth",2);
plot(1:5, var2_percentage(1:5),"ko-
", "MarkerSize",8,"LineWidth",2,'MarkerFaceColor','#000000');
text(20,0.2,sprintf('First 5 Sum is:
%.3f',sum(var2_percentage(1:5))), 'FontSize',16 );
set(gca, 'LineWidth',1, 'FontSize',14);
xlabel('Mode');
ylabel('% Variance');

hold off;
```

(D)



The calculated \mathbf{Z}_p file represents a smooth version of the original \mathbf{Z} file and keeps the main tendency and characteristics of \mathbf{Z} as well. It confirms our knowledge that the factors, contributing most of the variance, can already represent the data very good, by ignoring the small variance. Thus, the \mathbf{Z}_p shows better (= more obvious by eye) symmetric features around the ocean ridge ($x = 0\text{km}$).

```
% D.
% close all;

Sp = zeros(length(S),length(S));
Sp(1:5,1:5) = S(1:5,1:5);

Zp = U*Sp*V';

rows = [1 80 120];
```

```

figure(4);
for r = rows
    i = find(rows==r);
    axx(i)=subplot('Position',[0.1 0.1+0.3*(3-i) 0.8 0.3]);
    plot(x, Zp(i,:), "r-" , ...
        x, Z(i,:), "k-", ...
        'LineWidth',2);
    legend(sprintf('row %g of Z_p',r),sprintf('row %g of
Z',r))
end

set(axx, 'LineWidth',1, 'FontSize',14, 'YTickLabel', '');
set(axx(1:2), 'XTickLabel', '');
xlabel('Distance(km)');

```